

Appendix

M. Pitkänen,

August 16, 2023

Email: matpitka6@gmail.com.

http://tgdtheory.com/public_html/.

Recent postal address: Rinnekatu 2-4 A 8, 03620, Karkkila, Finland.

Contents

1	Introduction	3
2	Embedding space $M^4 \times CP_2$	3
2.1	Basic facts about CP_2	4
2.1.1	CP_2 as a manifold	4
2.1.2	Metric and Kähler structure of CP_2	4
2.1.3	Spinors In CP_2	7
2.1.4	Geodesic sub-manifolds of CP_2	7
2.2	CP_2 geometry and Standard Model symmetries	8
2.2.1	Identification of the electro-weak couplings	8
2.2.2	Electroweak symmetry breaking	12
2.2.3	Discrete symmetries	13
3	Induction procedure and many-sheeted space-time	14
3.1	Induction procedure for gauge fields and spinor connection	14
3.2	Induced gauge fields for space-times for which CP_2 projection is a geodesic sphere	15
3.3	Many-sheeted space-time	15
3.3.1	Superposition of effects instead of superposition of fields	15
3.3.2	Wormhole contacts	15
3.3.3	The relationship between the many-sheeted space-time of TGD and of GRT space-time	16
3.3.4	Topological field quantization and the notion of magnetic body	16
3.4	Embedding space spinors and induced spinors	16
3.5	About induced gauge fields	17
3.5.1	Space-times with vanishing em, Z^0 , or Kähler fields	18
3.5.2	The effective form of CP_2 metric for surfaces with 2-dimensional CP_2 projection	19
3.5.3	Topological quantum numbers	19

4	The relationship of TGD to QFT and string models	20
4.1	TGD as a generalization of wave mechanism obtained by replacing point-like particles with 3-surfaces	20
4.2	Extension of superconformal invariance	20
4.3	String-like objects and strings	21
4.4	TGD view of elementary particles	21
5	About the selection of the action defining the Kähler function of the "world of classical worlds" (WCW)	22
5.1	Could twistor lift fix the choice of the action uniquely?	22
5.2	Two paradoxes	23
5.2.1	The hierarchy subalgebras of supersymplectic algebra implies the decomposition of WCW into sectors with different actions	23
5.2.2	Number theoretic vision implies the decomposition of WCW into sectors with different actions	24
5.2.3	Number theoretic discretization of WCW and maxima of WCW Kähler function	26
6	Number theoretic vision of TGD	26
6.1	p-Adic numbers and TGD	27
6.1.1	p-Adic number fields	27
6.1.2	Canonical correspondence between p-adic and real numbers	28
6.1.3	The notion of p-adic manifold	30
6.2	Hierarchy of Planck constants and dark matter hierarchy	31
6.3	$M^8 - H$ duality as it is towards the end of 2021	31
7	Zero energy ontology (ZEO)	32
7.1	Basic motivations and ideas of ZEO	32
7.2	Some implications of ZEO	33
8	Some notions relevant to TGD inspired consciousness and quantum biology	34
8.1	The notion of magnetic body	34
8.2	Number theoretic entropy and negentropic entanglement	34
8.3	Life as something residing in the intersection of reality and p-adicities	34
8.4	Sharing of mental images	35
8.5	Time mirror mechanism	35

1 Introduction

Originally this appendix was meant to be a purely technical summary of basic facts but in its recent form it tries to briefly summarize those basic visions about TGD which I dare to regard as stabilized. I have added illustrations making it easier to build mental images about what is involved and represented briefly the key arguments. This chapter is hoped to help the reader to get fast grasp about the concepts of TGD.

The basic properties of embedding space and related spaces are discussed and the relationship of CP_2 to the standard model is summarized. The basic vision is simple: the geometry of the embedding space $H = M^4 \times CP_2$ geometrizes standard model symmetries and quantum numbers. The assumption that space-time surfaces are basic objects, brings in dynamics as dynamics of 3-D surfaces based on the induced geometry. Second quantization of free spinor fields of H induces quantization at the level of H , which means a dramatic simplification.

The notions of induction of metric and spinor connection, and of spinor structure are discussed. Many-sheeted space-time and related notions such as topological field quantization and the relationship many-sheeted space-time to that of GRT space-time are discussed as well as the recent view about induced spinor fields and the emergence of fermionic strings. Also the relationship to string models is discussed briefly.

Various topics related to p-adic numbers are summarized with a brief definition of p-adic manifold and the idea about generalization of the number concept by gluing real and p-adic number fields to a larger book like structure analogous to adèle [L1, L2]. In the recent view of quantum TGD [L17], both notions reduce to physics as number theory vision, which relies on $M^8 - H$ duality [L8, L9] and is complementary to the physics as geometry vision.

Zero energy ontology (ZEO) [L7] [K13] has become a central part of quantum TGD and leads to a TGD inspired theory of consciousness as a generalization of quantum measurement theory having quantum biology as an application. Also these aspects of TGD are briefly discussed.

2 Embedding space $M^4 \times CP_2$

Space-times are regarded as 4-surfaces in $H = M^4 \times CP_2$ the Cartesian product of empty Minkowski space - the space-time of special relativity - and compact 4-D space CP_2 with size scale of order 10^4 Planck lengths. One can say that embedding space is obtained by replacing each point m of empty Minkowski space with 4-D tiny CP_2 . The space-time of general relativity is replaced by a 4-D surface in H which has very complex topology. The notion of many-sheeted space-time gives an idea about what is involved.

Fig. 1. Embedding space $H = M^4 \times CP_2$ as Cartesian product of Minkowski space M^4 and complex projective space CP_2 . <http://tgdtheory.fi/appfigures/Hoo.jpg>

Denote by M_+^4 and M_-^4 the future and past directed lightcones of M^4 . Denote their intersection, which is not unique, by CD. In zero energy ontology (ZEO) [L7, L11] [K13] causal diamond (CD) is defined as cartesian product $CD \times CP_2$. Often I use CD to refer just to $CD \times CP_2$ since CP_2 factor is relevant from the point of view of ZEO.

Fig. 2. Future and past light-cones M_+^4 and M_-^4 . Causal diamonds (CD) are defined as their intersections. <http://tgdtheory.fi/appfigures/futurepast.jpg>

Fig. 3. Causal diamond (CD) is highly analogous to Penrose diagram but simpler. <http://tgdtheory.fi/appfigures/penrose.jpg>

A rather recent discovery was that CP_2 is the only compact 4-manifold with Euclidian signature of metric allowing twistor space with Kähler structure. M^4 is in turn is the only 4-D space with Minkowskian signature of metric allowing twistor space with Kähler structure [A4] so that $H = M^4 \times CP_2$ is twistorially unique.

One can loosely say that quantum states in a given sector of “world of classical worlds” (WCW) are superpositions of space-time surfaces inside CDs and that positive and negative energy parts of zero energy states are localized and past and future boundaries of CDs. CDs form a hierarchy. One can have CDs within CDs and CDs can also overlap. The size of CD is characterized by the proper time distance between its two tips. One can perform both translations and also Lorentz

boosts of CD leaving either boundary invariant. Therefore one can assign to CDs a moduli space and speak about wave function in this moduli space.

In number theoretic approach it is natural to restrict the allowed Lorentz boosts to some discrete subgroup of Lorentz group and also the distances between the tips of CDs to multiples of CP_2 radius defined by the length of its geodesic. Therefore the moduli space of CDs discretizes. The quantization of cosmic recession velocities for which there are indications, could relate to this quantization.

2.1 Basic facts about CP_2

CP_2 as a four-manifold is very special. The following arguments demonstrate that it codes for the symmetries of standard models via its isometries and holonomies.

2.1.1 CP_2 as a manifold

CP_2 , the complex projective space of two complex dimensions, is obtained by identifying the points of complex 3-space C^3 under the projective equivalence

$$(z^1, z^2, z^3) \equiv \lambda(z^1, z^2, z^3) . \quad (2.1)$$

Here λ is any non-zero complex number. Note that CP_2 can be also regarded as the coset space $SU(3)/U(2)$. The pair z^i/z^j for fixed j and $z^i \neq 0$ defines a complex coordinate chart for CP_2 . As j runs from 1 to 3 one obtains an atlas of three coordinate charts covering CP_2 , the charts being holomorphically related to each other (e.g. CP_2 is a complex manifold). The points $z^3 \neq 0$ form a subset of CP_2 homeomorphic to R^4 and the points with $z^3 = 0$ a set homeomorphic to S^2 . Therefore CP_2 is obtained by “adding the 2-sphere at infinity to R^4 ”.

Besides the standard complex coordinates $\xi^i = z^i/z^3$, $i = 1, 2$ the coordinates of Eguchi and Freund [A3] will be used and their relation to the complex coordinates is given by

$$\begin{aligned} \xi^1 &= z + it , \\ \xi^2 &= x + iy . \end{aligned} \quad (2.2)$$

These are related to the “spherical coordinates” via the equations

$$\begin{aligned} \xi^1 &= r \exp(i \frac{(\Psi + \Phi)}{2}) \cos(\frac{\Theta}{2}) , \\ \xi^2 &= r \exp(i \frac{(\Psi - \Phi)}{2}) \sin(\frac{\Theta}{2}) . \end{aligned} \quad (2.3)$$

The ranges of the variables r, Θ, Φ, Ψ are $[0, \infty], [0, \pi], [0, 4\pi], [0, 2\pi]$ respectively.

Considered as a real four-manifold CP_2 is compact and simply connected, with Euler number Euler number 3, Pontryagin number 3 and second $b = 1$.

Fig. 4. CP_2 as manifold. <http://tgdtheory.fi/appfigures/cp2.jpg>

2.1.2 Metric and Kähler structure of CP_2

In order to obtain a natural metric for CP_2 , observe that CP_2 can be thought of as a set of the orbits of the isometries $z^i \rightarrow \exp(i\alpha)z^i$ on the sphere S^5 : $\sum z^i \bar{z}^i = R^2$. The metric of CP_2 is obtained by projecting the metric of S^5 orthogonally to the orbits of the isometries. Therefore the distance between the points of CP_2 is that between the representative orbits on S^5 .

The line element has the following form in the complex coordinates

$$ds^2 = g_{a\bar{b}} d\xi^a d\bar{\xi}^b , \quad (2.4)$$

where the Hermitian, in fact Kähler metric $g_{a\bar{b}}$ is defined by

$$g_{a\bar{b}} = R^2 \partial_a \partial_{\bar{b}} K , \quad (2.5)$$

where the function K , Kähler function, is defined as

$$\begin{aligned} K &= \log(F) , \\ F &= 1 + r^2 . \end{aligned} \quad (2.6)$$

The Kähler function for S^2 has the same form. It gives the S^2 metric $dzd\bar{z}/(1+r^2)^2$ related to its standard form in spherical coordinates by the coordinate transformation $(r, \phi) = (\tan(\theta/2), \phi)$.

The representation of the CP_2 metric is deducible from S^5 metric is obtained by putting the angle coordinate of a geodesic sphere constant in it and is given

$$\frac{ds^2}{R^2} = \frac{(dr^2 + r^2 \sigma_3^2)}{F^2} + \frac{r^2(\sigma_1^2 + \sigma_2^2)}{F} , \quad (2.7)$$

where the quantities σ_i are defined as

$$\begin{aligned} r^2 \sigma_1 &= \text{Im}(\xi^1 d\xi^2 - \xi^2 d\xi^1) , \\ r^2 \sigma_2 &= -\text{Re}(\xi^1 d\xi^2 - \xi^2 d\xi^1) , \\ r^2 \sigma_3 &= -\text{Im}(\xi^1 d\bar{\xi}^1 + \xi^2 d\bar{\xi}^2) . \end{aligned} \quad (2.8)$$

R denotes the radius of the geodesic circle of CP_2 . The vierbein forms, which satisfy the defining relation

$$s_{kl} = R^2 \sum_A e_k^A e_l^A , \quad (2.9)$$

are given by

$$\begin{aligned} e^0 &= \frac{dr}{F} , & e^1 &= \frac{r\sigma_1}{\sqrt{F}} , \\ e^2 &= \frac{r\sigma_2}{\sqrt{F}} , & e^3 &= \frac{r\sigma_3}{F} . \end{aligned} \quad (2.10)$$

The explicit representations of vierbein vectors are given by

$$\begin{aligned} e^0 &= \frac{dr}{F} , & e^1 &= \frac{r(\sin\Theta \cos\Psi d\Phi + \sin\Psi d\Theta)}{2\sqrt{F}} , \\ e^2 &= \frac{r(\sin\Theta \sin\Psi d\Phi - \cos\Psi d\Theta)}{2\sqrt{F}} , & e^3 &= \frac{r(d\Psi + \cos\Theta d\Phi)}{2F} . \end{aligned} \quad (2.11)$$

The explicit representation of the line element is given by the expression

$$ds^2/R^2 = \frac{dr^2}{F^2} + \frac{r^2}{4F^2} (d\Psi + \cos\Theta d\Phi)^2 + \frac{r^2}{4F} (d\Theta^2 + \sin^2\Theta d\Phi^2) . \quad (2.12)$$

From this expression one finds that at coordinate infinity $r = \infty$ line element reduces to $\frac{r^2}{4F} (d\Theta^2 + \sin^2\Theta d\Phi^2)$ of S^2 meaning that 3-sphere degenerates metrically to 2-sphere and one can say that CP_2 is obtained by adding to R^4 a 2-sphere at infinity.

The vierbein connection satisfying the defining relation

$$de^A = -V_B^A \wedge e^B , \quad (2.13)$$

is given by

$$\begin{aligned} V_{01} &= -\frac{e^1}{r} , & V_{23} &= \frac{e^1}{r} , \\ V_{02} &= -\frac{e^2}{r} , & V_{31} &= \frac{e^2}{r} , \\ V_{03} &= (r - \frac{1}{r})e^3 , & V_{12} &= (2r + \frac{1}{r})e^3 . \end{aligned} \quad (2.14)$$

The representation of the covariantly constant curvature tensor is given by

$$\begin{aligned} R_{01} &= e^0 \wedge e^1 - e^2 \wedge e^3 , & R_{23} &= e^0 \wedge e^1 - e^2 \wedge e^3 , \\ R_{02} &= e^0 \wedge e^2 - e^3 \wedge e^1 , & R_{31} &= -e^0 \wedge e^2 + e^3 \wedge e^1 , \\ R_{03} &= 4e^0 \wedge e^3 + 2e^1 \wedge e^2 , & R_{12} &= 2e^0 \wedge e^3 + 4e^1 \wedge e^2 . \end{aligned} \quad (2.15)$$

Metric defines a real, covariantly constant, and therefore closed 2-form J

$$J = -is_{a\bar{b}}d\xi^a d\bar{\xi}^b , \quad (2.16)$$

the so called Kähler form. Kähler form J defines in CP_2 a symplectic structure because it satisfies the condition

$$J^k_r J^{rl} = -s^{kl} . \quad (2.17)$$

The condition states that J and g give representations of real unit and imaginary units related by the formula $i^2 = -1$.

Kähler form is expressible locally in terms of Kähler gauge potential

$$J = dB , \quad (2.18)$$

where B is the so called Kähler potential, which is not defined globally since J describes homological magnetic monopole.

$dJ = ddB = 0$ gives the topological half of Maxwell equations (vanishing of magnetic charges and Faraday's induction law) and self-duality $*J = J$ reduces the remaining equations to $dJ = 0$. Hence the Kähler form can be regarded as a curvature form of a $U(1)$ gauge potential B carrying a magnetic charge of unit $1/2g$ (g denotes the gauge coupling).

The magnetic flux of J through a 2-surface in CP_2 is proportional to its homology equivalence class, which is integer valued. The explicit representations of J and B are given by

$$\begin{aligned} B &= 2re^3 , \\ J &= 2(e^0 \wedge e^3 + e^1 \wedge e^2) = \frac{r}{F^2} dr \wedge (d\Psi + \cos\Theta d\Phi) + \frac{r^2}{2F} \sin\Theta d\Theta \wedge d\Phi . \end{aligned} \quad (2.19)$$

The vierbein curvature form and Kähler form are covariantly constant and have in the complex coordinates only components of type $(1, 1)$.

Useful coordinates for CP_2 are the so called canonical (or symplectic or Darboux) coordinates in which the Kähler potential and Kähler form have very simple expressions

$$\begin{aligned} B &= \sum_{k=1,2} P_k dQ_k , \\ J &= \sum_{k=1,2} dP_k \wedge dQ_k . \end{aligned} \quad (2.20)$$

The relationship of the canonical coordinates to the "spherical" coordinates is given by the equations

$$\begin{aligned}
P_1 &= -\frac{1}{1+r^2} , \\
P_2 &= -\frac{r^2 \cos \Theta}{2(1+r^2)} , \\
Q_1 &= \Psi , \\
Q_2 &= \Phi .
\end{aligned} \tag{2.21}$$

2.1.3 Spinors In CP_2

CP_2 doesn't allow spinor structure in the conventional sense [A2]. However, the coupling of the spinors to a half odd multiple of the Kähler potential leads to a respectable spinor structure. Because the delicacies associated with the spinor structure of CP_2 play a fundamental role in TGD, the arguments of Hawking are repeated here.

To see how the space can fail to have an ordinary spinor structure consider the parallel transport of the vierbein in a simply connected space M . The parallel propagation around a closed curve with a base point x leads to a rotated vierbein at x : $e^A = R_B^A e^B$ and one can associate to each closed path an element of $SO(4)$.

Consider now a one-parameter family of closed curves $\gamma(v) : v \in (0, 1)$ with the same base point x and $\gamma(0)$ and $\gamma(1)$ trivial paths. Clearly these paths define a sphere S^2 in M and the element $R_B^A(v)$ defines a closed path in $SO(4)$. When the sphere S^2 is contractible to a point e.g., homologically trivial, the path in $SO(4)$ is also contractible to a point and therefore represents a trivial element of the homotopy group $\Pi_1(SO(4)) = Z_2$.

For a homologically nontrivial 2-surface S^2 the associated path in $SO(4)$ can be homotopically nontrivial and therefore corresponds to a nonclosed path in the covering group $Spin(4)$ (leading from the matrix 1 to -1 in the matrix representation). Assume this is the case.

Assume now that the space allows spinor structure. Then one can parallel propagate also spinors and by the above construction associate a closed path of $Spin(4)$ to the surface S^2 . Now, however this path corresponds to a lift of the corresponding $SO(4)$ path and cannot be closed. Thus one ends up with a contradiction.

From the preceding argument it is clear that one could compensate the non-allowed -1 - factor associated with the parallel transport of the spinor around the sphere S^2 by coupling it to a gauge potential in such a way that in the parallel transport the gauge potential introduces a compensating -1 -factor. For a $U(1)$ gauge potential this factor is given by the exponential $\exp(i2\Phi)$, where Φ is the magnetic flux through the surface. This factor has the value -1 provided the $U(1)$ potential carries half odd multiple of Dirac charge $1/2g$. In case of CP_2 the required gauge potential is half odd multiple of the Kähler potential B defined previously. In the case of $M^4 \times CP_2$ one can in addition couple the spinor components with different chiralities independently to an odd multiple of $B/2$.

2.1.4 Geodesic sub-manifolds of CP_2

Geodesic sub-manifolds are defined as sub-manifolds having common geodesic lines with the embedding space. As a consequence the second fundamental form of the geodesic manifold vanishes, which means that the tangent vectors h_α^k (understood as vectors of H) are covariantly constant quantities with respect to the covariant derivative taking into account that the tangent vectors are vectors both with respect to H and X^4 .

In [A5] a general characterization of the geodesic sub-manifolds for an arbitrary symmetric space G/H is given. Geodesic sub-manifolds are in 1-1-correspondence with the so called Lie triple systems of the Lie-algebra g of the group G . The Lie triple system t is defined as a subspace of g characterized by the closedness property with respect to double commutation

$$[X, [Y, Z]] \in t \text{ for } X, Y, Z \in t . \tag{2.22}$$

$SU(3)$ allows, besides geodesic lines, two nonequivalent (not isometry related) geodesic spheres. This is understood by observing that $SU(3)$ allows two nonequivalent $SU(2)$ algebras corresponding

to subgroups $SO(3)$ (orthogonal 3×3 matrices) and the usual isospin group $SU(2)$. By taking any subset of two generators from these algebras, one obtains a Lie triple system and by exponentiating this system, one obtains a 2-dimensional geodesic sub-manifold of CP_2 .

Standard representatives for the geodesic spheres of CP_2 are given by the equations

$$S_I^2 : \xi^1 = \bar{\xi}^2 \text{ or equivalently } (\Theta = \pi/2, \Psi = 0) ,$$

$$S_{II}^2 : \xi^1 = \xi^2 \text{ or equivalently } (\Theta = \pi/2, \Phi = 0) .$$

The non-equivalence of these sub-manifolds is clear from the fact that isometries act as holomorphic transformations in CP_2 . The vanishing of the second fundamental form is also easy to verify. The first geodesic manifold is homologically trivial: in fact, the induced Kähler form vanishes identically for S_I^2 . S_{II}^2 is homologically nontrivial and the flux of the Kähler form gives its homology equivalence class.

2.2 CP_2 geometry and Standard Model symmetries

2.2.1 Identification of the electro-weak couplings

The delicacies of the spinor structure of CP_2 make it a unique candidate for space S . First, the coupling of the spinors to the $U(1)$ gauge potential defined by the Kähler structure provides the missing $U(1)$ factor in the gauge group. Secondly, it is possible to couple different H -chiralities independently to a half odd multiple of the Kähler potential. Thus the hopes of obtaining a correct spectrum for the electromagnetic charge are considerable. In the following it will be demonstrated that the couplings of the induced spinor connection are indeed those of the GWS model [B5] and in particular that the right handed neutrinos decouple completely from the electro-weak interactions.

To begin with, recall that the space H allows to define three different chiralities for spinors. Spinors with fixed H -chirality $e = \pm 1$, CP_2 -chirality l, r and M^4 -chirality L, R are defined by the condition

$$\begin{aligned} \Gamma\Psi &= e\Psi , \\ e &= \pm 1 , \end{aligned} \tag{2.23}$$

where Γ denotes the matrix $\Gamma_9 = \gamma_5 \otimes \gamma_5$, $1 \otimes \gamma_5$ and $\gamma_5 \otimes 1$ respectively. Clearly, for a fixed H -chirality CP_2 - and M^4 -chiralities are correlated.

The spinors with H -chirality $e = \pm 1$ can be identified as quark and lepton like spinors respectively. The separate conservation of baryon and lepton numbers can be understood as a consequence of generalized chiral invariance if this identification is accepted. For the spinors with a definite H -chirality one can identify the vielbein group of CP_2 as the electro-weak group: $SO(4)$ having as its covering group $SU(2)_L \times SU(2)_R$.

The covariant derivatives are defined by the spinorial connection

$$A = V + \frac{B}{2}(n_+ 1_+ + n_- 1_-) . \tag{2.24}$$

Here V and B denote the projections of the vielbein and Kähler gauge potentials respectively and $1_{+(-)}$ projects to the spinor H -chirality $+(-)$. The integers n_{\pm} are odd from the requirement of a respectable spinor structure.

The explicit representation of the vielbein connection V and of B are given by the equations

$$\begin{aligned} V_{01} &= -\frac{e^1}{r_2} , & V_{23} &= \frac{e^1}{r_2} , \\ V_{02} &= -\frac{e^2}{r} , & V_{31} &= \frac{e^2}{r} , \\ V_{03} &= (r - \frac{1}{r})e^3 , & V_{12} &= (2r + \frac{1}{r})e^3 , \end{aligned} \tag{2.25}$$

and

$$B = 2re^3 , \quad (2.26)$$

respectively. The explicit representation of the vielbein is not needed here.

Let us first show that the charged part of the spinor connection couples purely left handedly. Identifying Σ_3^0 and Σ_2^1 as the diagonal (neutral) Lie-algebra generators of $SO(4)$, one finds that the charged part of the spinor connection is given by

$$A_{ch} = 2V_{23}I_L^1 + 2V_{13}I_L^2 , \quad (2.27)$$

where one have defined

$$\begin{aligned} I_L^1 &= \frac{(\Sigma_{01} - \Sigma_{23})}{2} , \\ I_L^2 &= \frac{(\Sigma_{02} - \Sigma_{13})}{2} . \end{aligned} \quad (2.28)$$

A_{ch} is clearly left handed so that one can perform the identification of the gauge potential as

$$W^\pm = \frac{2(e^1 \pm ie^2)}{r} , \quad (2.29)$$

where W^\pm denotes the charged intermediate vector boson.

The covariantly constant curvature tensor is given by

$$\begin{aligned} R_{01} &= -R_{23} = e^0 \wedge e^1 - e^2 \wedge e^3 , \\ R_{02} &= -R_{31} = e^0 \wedge e^2 - e^3 \wedge e^1 , \\ R_{03} &= 4e^0 \wedge e^3 + 2e^1 \wedge e^2 , \\ R_{12} &= 2e^0 \wedge e^3 + 4e^1 \wedge e^2 . \end{aligned} \quad (2.30)$$

The charged part of the curvature tensor is left handed.

This is to be compared with the Weyl tensor, which defines a representation of quaternionic imaginary units.

$$\begin{aligned} W_{03} &= W_{12} \equiv 2I_3 = 2(e^0 \wedge e^3 + e^1 \wedge e^2) , \\ W_{01} &= W_{23} \equiv I_1 = -e^0 \wedge e^1 - e^2 \wedge e^3 , \\ W_{02} &= W_{31} \equiv I_2 = -e^0 \wedge e^2 - e^3 \wedge e^1 . \end{aligned} \quad (2.31)$$

The charged part of the Weyl tensor is right-handed and that the relative sign of the two terms in the curvature tensor and Weyl tensor are opposite.

Consider next the identification of the neutral gauge bosons γ and Z^0 as appropriate linear combinations of the two functionally independent quantities

$$\begin{aligned} X &= re^3 , \\ Y &= \frac{e^3}{r} , \end{aligned} \quad (2.32)$$

appearing in the neutral part of the spinor connection. We show first that the mere requirement that photon couples vectorially implies the basic coupling structure of the GWS model leaving only the value of Weinberg angle undetermined.

To begin with let us define

$$\begin{aligned} \bar{\gamma} &= aX + bY , \\ \bar{Z}^0 &= cX + dY , \end{aligned} \quad (2.33)$$

where the normalization condition

$$ad - bc = 1 \quad ,$$

is satisfied. The physical fields γ and Z^0 are related to $\bar{\gamma}$ and \bar{Z}^0 by simple normalization factors.

Expressing the neutral part of the spinor connection in term of these fields one obtains

$$\begin{aligned} A_{nc} &= [(c+d)2\Sigma_{03} + (2d-c)2\Sigma_{12} + d(n_+1_+ + n_-1_-)]\bar{\gamma} \\ &+ [(a-b)2\Sigma_{03} + (a-2b)2\Sigma_{12} - b(n_+1_+ + n_-1_-)]\bar{Z}^0 \quad . \end{aligned} \quad (2.34)$$

Identifying Σ_{12} and $\Sigma_{03} = 1 \times \gamma_5 \Sigma_{12}$ as vectorial and axial Lie-algebra generators, respectively, the requirement that γ couples vectorially leads to the condition

$$c = -d \quad . \quad (2.35)$$

Using this result plus previous equations, one obtains for the neutral part of the connection the expression

$$A_{nc} = \gamma Q_{em} + Z^0 (I_L^3 - \sin^2 \theta_W Q_{em}) \quad . \quad (2.36)$$

Here the electromagnetic charge Q_{em} and the weak isospin are defined by

$$\begin{aligned} Q_{em} &= \Sigma^{12} + \frac{(n_+1_+ + n_-1_-)}{6} \quad , \\ I_L^3 &= \frac{(\Sigma^{12} - \Sigma^{03})}{2} \quad . \end{aligned} \quad (2.37)$$

The fields γ and Z^0 are defined via the relations

$$\begin{aligned} \gamma &= 6d\bar{\gamma} = \frac{6}{(a+b)}(aX + bY) \quad , \\ Z^0 &= 4(a+b)\bar{Z}^0 = 4(X - Y) \quad . \end{aligned} \quad (2.38)$$

The value of the Weinberg angle is given by

$$\sin^2 \theta_W = \frac{3b}{2(a+b)} \quad , \quad (2.39)$$

and is not fixed completely. Observe that right handed neutrinos decouple completely from the electro-weak interactions.

The determination of the value of the Weinberg angle is a dynamical problem. The original approach was based on the assumption that it makes sense to talk about electroweak action defined at fundamental level and introduce a symmetry breaking by adding an additional term proportional to Kähler action. The recent view is that Kähler action plus volume term defines the fundamental action.

The Weinberg angle is completely fixed if one requires that the electroweak action contains no cross term of type γZ^0 . This leads to a definite value for the Weinberg angle.

One can however add a symmetry breaking term proportional to Kähler action and this changes the value of the Weinberg angle. As a matter fact, color gauge action identifying color gauge field as proportional to $H^A J_{\alpha\beta}$ is proportional to Kähler action. A possible interpretation would be as a sum of electroweak and color gauge interactions.

To evaluate the value of the Weinberg angle one can express the neutral part F_{nc} of the induced gauge field as

$$F_{nc} = 2R_{03}\Sigma^{03} + 2R_{12}\Sigma^{12} + J(n_+1_+ + n_-1_-) , \quad (2.40)$$

where one has

$$\begin{aligned} R_{03} &= 2(2e^0 \wedge e^3 + e^1 \wedge e^2) , \\ R_{12} &= 2(e^0 \wedge e^3 + 2e^1 \wedge e^2) , \\ J &= 2(e^0 \wedge e^3 + e^1 \wedge e^2) , \end{aligned} \quad (2.41)$$

in terms of the fields γ and Z^0 (photon and Z - boson)

$$F_{nc} = \gamma Q_{em} + Z^0(I_L^3 - \sin^2\theta_W Q_{em}) . \quad (2.42)$$

Evaluating the expressions above, one obtains for γ and Z^0 the expressions

$$\begin{aligned} \gamma &= 3J - \sin^2\theta_W R_{12} , \\ Z^0 &= 2R_{03} . \end{aligned} \quad (2.43)$$

For the Kähler field one obtains

$$J = \frac{1}{3}(\gamma + \sin^2\theta_W Z^0) . \quad (2.44)$$

Expressing the neutral part of the symmetry broken YM action

$$\begin{aligned} L_{ew} &= L_{sym} + f J^{\alpha\beta} J_{\alpha\beta} , \\ L_{sym} &= \frac{1}{4g^2} Tr(F^{\alpha\beta} F_{\alpha\beta}) , \end{aligned} \quad (2.45)$$

where the trace is taken in spinor representation, in terms of γ and Z^0 one obtains for the coefficient X of the γZ^0 cross term (this coefficient must vanish) the expression

$$\begin{aligned} X &= -\frac{K}{2g^2} + \frac{fp}{18} , \\ K &= Tr [Q_{em}(I_L^3 - \sin^2\theta_W Q_{em})] , \end{aligned} \quad (2.46)$$

This parameter can be calculated by substituting the values of quark and lepton charges and weak isospins.

In the general case the value of the coefficient K is given by

$$K = \sum_i \left[-\frac{(18 + 2n_i^2)\sin^2\theta_W}{9} \right] , \quad (2.47)$$

where the sum is over the spinor chiralities, which appear as elementary fermions and n_i is the integer describing the coupling of the spinor field to the Kähler potential. The cross term vanishes provided the value of the Weinberg angle is given by

$$\sin^2\theta_W = \frac{9 \sum_i 1}{(fg^2 + 2 \sum_i (18 + n_i^2))} . \quad (2.48)$$

In the scenario where both leptons and quarks are elementary fermions the value of the Weinberg angle is given by

$$\sin^2\theta_W = \frac{9}{\left(\frac{fg^2}{2} + 28\right)}. \quad (2.49)$$

The bare value of the Weinberg angle is $9/28$ in this scenario, which is not far from the typical value $9/24$ of GUTs at high energies [B1]. The experimental value at the scale length scale of the electron can be deduced from the ratio of W and Z boson masses as $\sin^2\theta_W = 1 - (m_W/m_Z)^2 \simeq .22290$. This ratio and also the weak boson masses depend on the length scale.

If one interprets the additional term proportional to J as color action, one could perhaps interpret the value of Weinberg angle as expressing a connection between strong and weak coupling constant evolution. The limit $f \rightarrow 0$ should correspond to an infinite value of color coupling strength and at this limit one would have $\sin^2\theta_W = \frac{9}{28}$ for $f/g^2 \rightarrow 0$. This does not make sense since the Weinberg angle is in the standard model much smaller in QCD scale Λ corresponding roughly to pion mass scale. The Weinberg angle is in principle predicted by the p-adic coupling constant evolution fixed by the number theoretical vision of TGD.

One could however have a sum of electroweak action, correction terms changing the value of Weinberg angle, and color action and coupling constant evolution could be understood in terms of the coupling parameters involved.

2.2.2 Electroweak symmetry breaking

One of the hardest challenges in the development of the TGD based view of weak symmetry breaking was the fact that classical field equations allow space-time surfaces with finite but arbitrarily large size. For a fixed space-time surface, the induced gauge fields, including classical weak fields, are long ranged. On the other hand, the large mass for weak bosons would require a short correlation length. How can one understand this together with the fact that a photon has a long correlation length?

In zero energy ontology quantum states are superpositions of space-time surfaces as analogs of almost unique Bohr orbits of particles identified as 3-D surfaces. For some reason the superposition should be such that the quantum averages of weak gauge boson fields vanish below the weak scale whereas the quantum average of electromagnetic fields is non-vanishing.

This is indeed the case.

1. The supersymplectic symmetries form isometries of the world of classical worlds (WCW) and they act in CP_2 degrees of freedom as symplectic transformations leaving the CP_2 symplectic form J invariant and therefore also its contribution to the electromagnetic field since this part is the same for all space-time surfaces in the superposition of space-time surfaces as a representation of supersymplectic isometry group (as a special case a representation of color group).
2. In TGD, color and electroweak symmetries acting as holonomies are not independent and for the $SU(2)_L$ part of induced spinor connection the symplectic transformations induces $SU(2)_L \times U(1)_R$ gauge transformation. This suggests that the quantum expectations of the induced weak fields over the space-time surfaces vanish above the quantum coherence scale. The averages of W and of the left handed part of Z^0 should therefore vanish.
3. $\langle Z^0 \rangle$ should vanish. For $U(1)_R$ part of Z^0 , the action of gauge transformation is trivial in gauge theory. Now however the space-time surface changes under symplectic transformations and this could make the average of the right-handed part of Z^0 vanishing. The vanishing of the average of the axial part of the Z^0 is suggested by the partially conserved axial current hypothesis.

One can formulate this picture quantitatively.

1. The electromagnetic field [L19] contains, besides the induced Kähler form, also the induced curvature form R_{12} , which couples vectorially. Conserved vector current hypothesis suggests that the average of R_{12} is non-vanishing. One can express the neutral part of the induced gauge field in terms of induced spinor curvature and Kähler form J as

$$\begin{aligned}
R_{03} &= 2(2e^0 \wedge e^3 + e^1 \wedge e^2) = J + 2e^0 \wedge e^3 , \\
J &= 2(e^0 \wedge e^3 + e^1 \wedge e^2) , \\
R_{12} &= 2(e^0 \wedge e^3 + 2e^1 \wedge e^2) = 3J - 2e^0 \wedge e^3 ,
\end{aligned} \tag{2.50}$$

2. The induced fields γ and Z^0 (photon and Z - boson) can be expressed as

$$\begin{aligned}
\gamma &= 3J - \sin^2\theta_W R_{12} , \\
Z^0 &= 2R_{03} = 2(J + 2e^0 \wedge e^3)
\end{aligned} \tag{2.51}$$

$$per. \tag{2.52}$$

The condition $\langle Z^0 \rangle = 0$ gives $2\langle e^0 \wedge e^3 \rangle = -2J$ and this in turn gives $\langle R_{12} \rangle = 4J$. The average over γ would be

$$\langle \gamma \rangle = (3 - 4\sin^2\theta_W)J .$$

For $\sin^2\theta_W = 3/4$ $\langle \gamma \rangle$ would vanish.

The quantum averages of classical weak fields quite generally vanish. What about correlation functions?

1. One expects that the correlators of classical weak fields as color invariants, and perhaps even symplectic invariants, are non-vanishing below the Compton length since in this kind of situation the points in the correlation function belong to the same 3-surface representing particle, such as hadron.
2. The intuitive picture is that in longer length scales one has disjoint 3-surfaces with a size scale of Compton length. If the states associated with two disjoint 3-surfaces are separately color invariant there are no correlations in color degrees of freedom and correlators reduce to the products of expectations of classical weak fields and vanish. This could also hold when the 3-surfaces are connected by flux tube bonds.

Below the Compton length weak bosons would thus behave as correlated massless fields. The Compton lengths of weak bosons are proportional to the value of effective Planck constant h_{eff} and in living systems the Compton lengths are proposed to be even of the order of cell size. This would explain the mysterious chiral selection in living systems requiring large parity violation.

3. What about the averages and correlators of color gauge fields? Classical color gauge fields are proportional to the products of Hamiltonians of color isometries induced Kähler form and the expectations of color Hamiltonians give vanishing average above Compton length and therefore vanishing average. Correlators are non-vanishing below the hadron scale. Gluons do not propagate in long scales for the same reason as weak bosons. This is implied by color confinement, which has also classical description in the sense that 3-surfaces have necessarily a finite size.

A large value of h_{eff} allows colored states even in biological scales below the Compton length since in this kind of situation the points in the correlation function belong to the same 3-surface representing particle, such as dark hadron.

2.2.3 Discrete symmetries

The treatment of discrete symmetries C, P, and T is based on the following requirements:

1. Symmetries must be realized as purely geometric transformations.
2. Transformation properties of the field variables should be essentially the same as in the conventional quantum field theories [B2].

The action of the reflection P on spinors of is given by

$$\Psi \rightarrow P\Psi = \gamma^0 \otimes \gamma^0 \Psi . \quad (2.53)$$

in the representation of the gamma matrices for which γ^0 is diagonal. It should be noticed that W and Z^0 bosons break parity symmetry as they should since their charge matrices do not commute with the matrix of P .

The guess that a complex conjugation in CP_2 is associated with T transformation of the physicist turns out to be correct. One can verify by a direct calculation that pure Dirac action is invariant under T realized according to

$$\begin{aligned} m^k &\rightarrow T(M^k) , \\ \xi^k &\rightarrow \bar{\xi}^k , \\ \Psi &\rightarrow \gamma^1 \gamma^3 \otimes 1 \Psi . \end{aligned} \quad (2.54)$$

The operation bearing closest resemblance to the ordinary charge conjugation corresponds geometrically to complex conjugation in CP_2 :

$$\begin{aligned} \xi^k &\rightarrow \bar{\xi}^k , \\ \Psi &\rightarrow \Psi^\dagger \gamma^2 \gamma^0 \otimes 1 . \end{aligned} \quad (2.55)$$

As one might have expected symmetries CP and T are exact symmetries of the pure Dirac action.

3 Induction procedure and many-sheeted space-time

Since the classical gauge fields are closely related in TGD framework, it is not possible to have space-time sheets carrying only single kind of gauge field. For instance, em fields are accompanied by Z^0 fields for extremals of Kähler action.

Classical em fields are always accompanied by Z^0 field and some components of color gauge field. For extremals having homologically non-trivial sphere as a CP_2 projection em and Z^0 fields are the only non-vanishing electroweak gauge fields. For homologically trivial sphere only W fields are non-vanishing. Color rotations does not affect the situation.

For vacuum extremals all electro-weak gauge fields are in general non-vanishing although the net gauge field has U(1) holonomy by 2-dimensionality of the CP_2 projection. Color gauge field has U(1) holonomy for all space-time surfaces and quantum classical correspondence suggest a weak form of color confinement meaning that physical states correspond to color neutral members of color multiplets.

3.1 Induction procedure for gauge fields and spinor connection

Induction procedure for gauge potentials and spinor structure is a standard procedure of bundle theory. If one has embedding of some manifold to the base space of a bundle, the bundle structure can be induced so that it has as a base space the imbedded manifold, whose points have as fiber the fiber if embedding space at their image points. In the recent case the embedding of space-time surface to embedding space defines the induction procedure. The induced gauge potentials and gauge fields are projections of the spinor connection of the embedding space to the space-time surface (see <http://tgdtheory.fi/appfigures/induct.jpg>).

Induction procedure makes sense also for the spinor fields of embedding space and one obtains geometrization of both electroweak gauge potentials and of spinors. The new element is induction of gamma matrices which gives their projections at space-time surface.

As a matter fact, the induced gamma matrices cannot appear in the counterpart of massless Dirac equation. To achieve super-symmetry, Dirac action must be replaced with Kähler-Dirac action for which gamma matrices are contractions of the canonical momentum currents of Kähler

action with embedding space gamma matrices. Induced gamma matrices in Dirac action would correspond to 4-volume as action.

Fig. 9. Induction of spinor connection and metric as projection to the space-time surface. <http://tgdtheory.fi/appfigures/induct.jpg>.

3.2 Induced gauge fields for space-times for which CP_2 projection is a geodesic sphere

If one requires that space-time surface is an extremal of Kähler action and has a 2-dimensional CP_2 projection, only vacuum extremals and space-time surfaces for which CP_2 projection is a geodesic sphere, are allowed. Homologically non-trivial geodesic sphere correspond to vanishing W fields and homologically non-trivial sphere to non-vanishing W fields but vanishing γ and Z^0 . This can be verified by explicit examples.

$r = \infty$ surface gives rise to a homologically non-trivial geodesic sphere for which e_0 and e_3 vanish imply the vanishing of W field. For space-time sheets for which CP_2 projection is $r = \infty$ homologically non-trivial geodesic sphere of CP_2 one has

$$\gamma = \left(\frac{3}{4} - \frac{\sin^2(\theta_W)}{2}\right)Z^0 \simeq \frac{5Z^0}{8} .$$

The induced W fields vanish in this case and they vanish also for all geodesic sphere obtained by $SU(3)$ rotation.

$Im(\xi^1) = Im(\xi^2) = 0$ corresponds to homologically trivial geodesic sphere. A more general representative is obtained by using for the phase angles of standard complex CP_2 coordinates constant values. In this case e^1 and e^3 vanish so that the induced em, Z^0 , and Kähler fields vanish but induced W fields are non-vanishing. This holds also for surfaces obtained by color rotation. Hence one can say that for non-vacuum extremals with 2-D CP_2 projection color rotations and weak symmetries commute.

3.3 Many-sheeted space-time

TGD space-time is many-sheeted: in other words, there are in general several space-sheets which have projection to the same M^4 region. Second manner to say this is that CP_2 coordinates are many-valued functions of M^4 coordinates. The original physical interpretation of many-sheeted space-time time was not correct: it was assumed that single sheet corresponds to GRT space-time and this obviously leads to difficulties since the induced gauge fields are expressible in terms of only four embedding space coordinates.

Fig. 10. Illustration of many-sheeted space-time of TGD. <http://tgdtheory.fi/appfigures/manysheeted.jpg>

3.3.1 Superposition of effects instead of superposition of fields

The first objection against TGD is that superposition is not possible for induced gauge fields and induced metric. The resolution of the problem is that it is effects which need to superpose, not the fields.

Test particle topologically condenses simultaneously to all space-time sheets having a projection to same region of M^4 (that is touches them). The superposition of effects of fields at various space-time sheets replaces the superposition of fields. This is crucial for the understanding also how GRT space-time relates to TGD space-time, which is also in the appendix of this book).

3.3.2 Wormhole contacts

Wormhole contacts are key element of many-sheeted space-time. One does not expect them to be stable unless there is non-trivial Kähler magnetic flux flowing through them so that the throats look like Kähler magnetic monopoles.

Fig. 11. Wormhole contact. <http://tgdtheory.fi/appfigures/wormholecontact.jpg>

Since the flow lines of Kähler magnetic field must be closed this requires the presence of another wormhole contact so that one obtains closed monopole flux tube decomposing to two Minkowskian

pieces at the two space-time sheets involved and two wormhole contacts with Euclidian signature of the induced metric. These objects are identified as space-time correlates of elementary particles and are clearly analogous to string like objects.

3.3.3 The relationship between the many-sheeted space-time of TGD and of GRT space-time

The space-time of general relativity is single-sheeted and there is no need to regard it as surface in H although the assumption about representability as vacuum extremal gives very powerful constraints in cosmology and astrophysics and might make sense in simple situations.

The space-time of GRT can be regarded as a long length scale approximation obtained by lumping together the sheets of the many-sheeted space-time to a region of M^4 and providing it with an effective metric obtained as sum of M^4 metric and deviations of the induced metrics of various space-time sheets from M^4 metric. Also induced gauge potentials sum up in the similar manner so that also the gauge fields of gauge theories would not be fundamental fields.

Fig. 12. The superposition of fields is replaced with the superposition of their effects in many-sheeted space-time. <http://tgdtheory.fi/appfigures/fieldsuperpose.jpg>

Space-time surfaces of TGD are considerably simpler objects than the space-times of general relativity and relate to GRT space-time like elementary particles to systems of condensed matter physics. Same can be said about fields since all fields are expressible in terms of embedding space coordinates and their gradients, and general coordinate invariance means that the number of bosonic field degrees is reduced locally to 4. TGD space-time can be said to be a microscopic description whereas GRT space-time a macroscopic description. In TGD complexity of space-time topology replaces the complexity due to large number of fields in quantum field theory.

3.3.4 Topological field quantization and the notion of magnetic body

Topological field quantization also TGD from Maxwell's theory. TGD predicts topological light rays ("massless extremals (MEs)") as space-time sheets carrying waves or arbitrary shape propagating with maximal signal velocity in single direction only and analogous to laser beams and carrying light-like gauge currents in the general case. There are also magnetic flux quanta and electric flux quanta. The deformations of cosmic strings with 2-D string orbit as M^4 projection gives rise to magnetic flux tubes carrying monopole flux made possible by CP_2 topology allowing homological Kähler magnetic monopoles.

Fig. 13. Topological quantization for magnetic fields replaces magnetic fields with bundles of them defining flux tubes as topological field quanta. <http://tgdtheory.fi/appfigures/field.jpg>

The imbeddability condition for say magnetic field means that the region containing constant magnetic field splits into flux quanta, say tubes and sheets carrying constant magnetic field. Unless one assumes a separate boundary term in Kähler action, boundaries in the usual sense are forbidden except as ends of space-time surfaces at the boundaries of causal diamonds. One obtains typically pairs of sheets glued together along their boundaries giving rise to flux tubes with closed cross section possibly carrying monopole flux.

These kind of flux tubes might make possible magnetic fields in cosmic scales already during primordial period of cosmology since no currents are needed to generate these magnetic fields: cosmic string would be indeed this kind of objects and would dominate during the primordial period. Even superconductors and maybe even ferromagnets could involve this kind of monopole flux tubes.

3.4 Embedding space spinors and induced spinors

One can geometrize also fermionic degrees of freedom by inducing the spinor structure of $M^4 \times CP_2$.

CP_2 does not allow spinor structure in the ordinary sense but one can couple the opposite H -chiralities of H -spinors to an $n = 1$ ($n = 3$) integer multiple of Kähler gauge potential to obtain a respectable modified spinor structure. The em charges of resulting spinors are fractional (integer valued) and the interpretation as quarks (leptons) makes sense since the couplings to the induced

spinor connection having interpretation in terms electro-weak gauge potential are identical to those assumed in standard model.

The notion of quark color differs from that of standard model.

1. Spinors do not couple to color gauge potential although the identification of color gauge potential as projection of $SU(3)$ Killing vector fields is possible. This coupling must emerge only at the effective gauge theory limit of TGD.
2. Spinor harmonics of embedding space correspond to triality $t = 1$ ($t = 0$) partial waves. The detailed correspondence between color and electroweak quantum numbers is however not correct as such and the interpretation of spinor harmonics of embedding space is as representations for ground states of super-conformal representations. The wormhole pairs associated with physical quarks and leptons must carry also neutrino pair to neutralize weak quantum numbers above the length scale of flux tube (weak scale or Compton length). The total color quantum numbers of these states must be those of standard model. For instance, the color quantum numbers of fundamental left-hand neutrino and lepton can compensate each other for the physical lepton. For fundamental quark-lepton pair they could sum up to those of physical quark.

The well-definedness of em charge is crucial condition.

1. Although the embedding space spinor connection carries W gauge potentials one can say that the embedding space spinor modes have well-defined em charge. One expects that this is true for induced spinor fields inside wormhole contacts with 4-D CP_2 projection and Euclidian signature of the induced metric.
2. The situation is not the same for the modes of induced spinor fields inside Minkowskian region and one must require that the CP_2 projection of the regions carrying induced spinor field is such that the induced W fields and above weak scale also the induced Z^0 fields vanish in order to avoid large parity breaking effects. This condition forces the CP_2 projection to be 2-dimensional. For a generic Minkowskian space-time region this is achieved only if the spinor modes are localized at 2-D surfaces of space-time surface - string world sheets and possibly also partonic 2-surfaces.
3. Also the Kähler-Dirac gamma matrices appearing in the modified Dirac equation must vanish in the directions normal to the 2-D surface in order that Kähler-Dirac equation can be satisfied. This does not seem plausible for space-time regions with 4-D CP_2 projection.
4. One can thus say that strings emerge from TGD in Minkowskian space-time regions. In particular, elementary particles are accompanied by a pair of fermionic strings at the opposite space-time sheets and connecting wormhole contacts. Quite generally, fundamental fermions would propagate at the boundaries of string world sheets as massless particles and wormhole contacts would define the stringy vertices of generalized Feynman diagrams. One obtains geometrized diagrammatics, which brings looks like a combination of stringy and Feynman diagrammatics.
5. This is what happens in the the generic situation. Cosmic strings could serve as examples about surfaces with 2-D CP_2 projection and carrying only em fields and allowing delocalization of spinor modes to the entire space-time surfaces.

3.5 About induced gauge fields

In the following the induced gauge fields are studied for general space-time surface without assuming the preferred extremal property (Bohr orbit property). Therefore the following arguments are somewhat obsolete in their generality.

3.5.1 Space-times with vanishing em, Z^0 , or Kähler fields

The following considerations apply to a more general situation in which the homologically trivial geodesic sphere and extremal property are not assumed. It must be emphasized that this case is possible in TGD framework only for a vanishing Kähler field.

Using spherical coordinates (r, Θ, Ψ, Φ) for CP_2 , the expression of Kähler form reads as

$$\begin{aligned} J &= \frac{r}{F^2} dr \wedge (d\Psi + \cos(\Theta)d\Phi) + \frac{r^2}{2F} \sin(\Theta) d\Theta \wedge d\Phi , \\ F &= 1 + r^2 . \end{aligned} \quad (3.1)$$

The general expression of electromagnetic field reads as

$$\begin{aligned} F_{em} &= (3 + 2p) \frac{r}{F^2} dr \wedge (d\Psi + \cos(\Theta)d\Phi) + (3 + p) \frac{r^2}{2F} \sin(\Theta) d\Theta \wedge d\Phi , \\ p &= \sin^2(\Theta_W) , \end{aligned} \quad (3.2)$$

where Θ_W denotes Weinberg angle.

1. The vanishing of the electromagnetic fields is guaranteed, when the conditions

$$\begin{aligned} \Psi &= k\Phi , \\ (3 + 2p) \frac{1}{r^2 F} (d(r^2)/d\Theta)(k + \cos(\Theta)) + (3 + p) \sin(\Theta) &= 0 , \end{aligned} \quad (3.3)$$

hold true. The conditions imply that CP_2 projection of the electromagnetically neutral space-time is 2-dimensional. Solving the differential equation one obtains

$$\begin{aligned} r &= \sqrt{\frac{X}{1-X}} , \\ X &= D \left[\left| \frac{k+u}{C} \right| \right]^\epsilon , \\ u &\equiv \cos(\Theta) , \quad C = k + \cos(\Theta_0) , \quad D = \frac{r_0^2}{1+r_0^2} , \quad \epsilon = \frac{3+p}{3+2p} , \end{aligned} \quad (3.4)$$

where C and D are integration constants. $0 \leq X \leq 1$ is required by the reality of r . $r = 0$ would correspond to $X = 0$ giving $u = -k$ achieved only for $|k| \leq 1$ and $r = \infty$ to $X = 1$ giving $|u+k| = [(1+r_0^2)/r_0^2]^{(3+2p)/(3+p)}$ achieved only for

$$\text{sign}(u+k) \times \left[\frac{1+r_0^2}{r_0^2} \right]^{\frac{3+2p}{3+p}} \leq k+1 ,$$

where $\text{sign}(x)$ denotes the sign of x .

The expressions for Kähler form and Z^0 field are given by

$$\begin{aligned} J &= -\frac{p}{3+2p} X du \wedge d\Phi , \\ Z^0 &= -\frac{6}{p} J . \end{aligned} \quad (3.5)$$

The components of the electromagnetic field generated by varying vacuum parameters are proportional to the components of the Kähler field: in particular, the magnetic field is parallel to the Kähler magnetic field. The generation of a long range Z^0 vacuum field is a purely TGD based feature not encountered in the standard gauge theories.

2. The vanishing of Z^0 fields is achieved by the replacement of the parameter ϵ with $\epsilon = 1/2$ as becomes clear by considering the condition stating that Z^0 field vanishes identically. Also the relationship $F_{em} = 3J = -\frac{3}{4}\frac{r^2}{F}du \wedge d\Phi$ is useful.
3. The vanishing Kähler field corresponds to $\epsilon = 1, p = 0$ in the formula for em neutral space-times. In this case classical em and Z^0 fields are proportional to each other:

$$\begin{aligned}
Z^0 &= 2e^0 \wedge e^3 = \frac{r}{F^2}(k+u)\frac{\partial r}{\partial u}du \wedge d\Phi = (k+u)du \wedge d\Phi , \\
r &= \sqrt{\frac{X}{1-X}} , \quad X = D|k+u| , \\
\gamma &= -\frac{p}{2}Z^0 .
\end{aligned} \tag{3.6}$$

For a vanishing value of Weinberg angle ($p = 0$) em field vanishes and only Z^0 field remains as a long range gauge field. Vacuum extremals for which long range Z^0 field vanishes but em field is non-vanishing are not possible.

3.5.2 The effective form of CP_2 metric for surfaces with 2-dimensional CP_2 projection

The effective form of the CP_2 metric for a space-time having vanishing em, Z^0 , or Kähler field is of practical value in the case of vacuum extremals and is given by

$$\begin{aligned}
ds_{eff}^2 &= (s_{rr}(\frac{dr}{d\Theta})^2 + s_{\Theta\Theta})d\Theta^2 + (s_{\Phi\Phi} + 2ks_{\Phi\Psi})d\Phi^2 = \frac{R^2}{4}[s_{\Theta\Theta}^{eff}d\Theta^2 + s_{\Phi\Phi}^{eff}d\Phi^2] , \\
s_{\Theta\Theta}^{eff} &= X \times \left[\frac{\epsilon^2(1-u^2)}{(k+u)^2} \times \frac{1}{1-X} + 1 - X \right] , \\
s_{\Phi\Phi}^{eff} &= X \times [(1-X)(k+u)^2 + 1 - u^2] ,
\end{aligned} \tag{3.7}$$

and is useful in the construction of vacuum embedding of, say Schwarzschild metric.

3.5.3 Topological quantum numbers

Space-times for which either em, Z^0 , or Kähler field vanishes decompose into regions characterized by six vacuum parameters: two of these quantum numbers (ω_1 and ω_2) are frequency type parameters, two (k_1 and k_2) are wave vector like quantum numbers, two of the quantum numbers (n_1 and n_2) are integers. The parameters ω_i and n_i will be referred as electric and magnetic quantum numbers. The existence of these quantum numbers is not a feature of these solutions alone but represents a much more general phenomenon differentiating in a clear cut manner between TGD and Maxwell's electrodynamics.

The simplest manner to avoid surface Kähler charges and discontinuities or infinities in the derivatives of CP_2 coordinates on the common boundary of two neighboring regions with different vacuum quantum numbers is topological field quantization, 3-space decomposes into disjoint topological field quanta, 3-surfaces having outer boundaries with possibly macroscopic size.

Under rather general conditions the coordinates Ψ and Φ can be written in the form

$$\begin{aligned}
\Psi &= \omega_2 m^0 + k_2 m^3 + n_2 \phi + \text{Fourier expansion} , \\
\Phi &= \omega_1 m^0 + k_1 m^3 + n_1 \phi + \text{Fourier expansion} .
\end{aligned} \tag{3.8}$$

m^0, m^3 and ϕ denote the coordinate variables of the cylindrical M^4 coordinates) so that one has $k = \omega_2/\omega_1 = n_2/n_1 = k_2/k_1$. The regions of the space-time surface with given values of the vacuum parameters ω_i, k_i and n_i and m and C are bounded by the surfaces at which space-time surface becomes ill-defined, say by $r > 0$ or $r < \infty$ surfaces.

The space-time surface decomposes into regions characterized by different values of the vacuum parameters r_0 and Θ_0 . At $r = \infty$ surfaces n_2, ω_2 and m can change since all values of Ψ correspond

to the same point of CP_2 : at $r = 0$ surfaces also n_1 and ω_1 can change since all values of Φ correspond to same point of CP_2 , too. If $r = 0$ or $r = \infty$ is not in the allowed range space-time surface develops a boundary.

This implies what might be called topological quantization since in general it is not possible to find a smooth global embedding for, say a constant magnetic field. Although global embedding exists it decomposes into regions with different values of the vacuum parameters and the coordinate u in general possesses discontinuous derivative at $r = 0$ and $r = \infty$ surfaces. A possible manner to avoid edges of space-time is to allow field quantization so that 3-space (and field) decomposes into disjoint quanta, which can be regarded as structurally stable units a 3-space (and of the gauge field). This doesn't exclude partial join along boundaries for neighboring field quanta provided some additional conditions guaranteeing the absence of edges are satisfied.

For instance, the vanishing of the electromagnetic fields implies that the condition

$$\Omega \equiv \frac{\omega_2}{n_2} - \frac{\omega_1}{n_1} = 0 \quad , \quad (3.9)$$

is satisfied. In particular, the ratio ω_2/ω_1 is rational number for the electromagnetically neutral regions of space-time surface. The change of the parameter n_1 and n_2 (ω_1 and ω_2) in general generates magnetic field and therefore these integers will be referred to as magnetic (electric) quantum numbers.

4 The relationship of TGD to QFT and string models

The recent view of the relationship of TGD to QFT and string models has developed slowly during years and it seems that in a certain sense TGD means a return to roots: instead of QFT like description involving path integral one would have wave mechanics for 3-surfaces.

4.1 TGD as a generalization of wave mechanism obtained by replacing point-like particles with 3-surfaces

The first vision of TGD was as a generalization of quantum field theory (string models) obtained by replacing pointlike particles (strings) as fundamental objects with 3-surfaces.

The later work has revealed that TGD could be seen as a generalization of the wave mechanism based on the replacement of a point-like particle with 3-D surface. This is due to holography implied by general coordinate invariance. The definition of the metric of the "world of classical worlds" (WCW) must assign a unique or at least almost unique space-time surface to a given 3-surface. This 4-surface is analogous to Bohr orbit so that also Bohr orbitology becomes an exact part of quantum physics. The failure of strict determinism forces to replace 3-surfaces with 4-surfaces and this leads to zero energy ontology (ZEO) in which quantum states are superpositions of space-time surfaces [K3, K2, K8] [L12, L17].

Fig. 5. TGD replaces point-like particles with 3-surfaces. <http://tgdtheory.fi/appfigures/particletgd.jpg>

4.2 Extension of superconformal invariance

The fact that light-like 3-surfaces are effectively metrically 2-dimensional and thus possess generalization of 2-dimensional conformal symmetries with light-like radial coordinate defining the analog of second complex coordinate suggests that this generalization could work and extend the super-conformal symmetries to their 4-D analogs.

The boundary $\delta M_+^4 = S^2 \times R_+$ of 4-D light-cone M_+^4 is also metrically 2-dimensional and allows extended conformal invariance. Also the group of isometries of light-cone boundary and of light-like 3-surfaces is infinite-dimensional since the conformal scalings of S^2 can be compensated by S^2 -local scaling of the light-like radial coordinate of R_+ . These simple facts mean that 4-dimensional Minkowski space and 4-dimensional space-time surfaces are in a completely unique position as far as symmetries are considered.

In fact, this leads to a generalization of the Kac-Moody type symmetries of string models. $\delta M_+^4 \times CP_2$ allows huge supersymplectic symmetries for which the radial light-like coordinate of

δM_+^4 plays the role of complex string coordinate in string models. These symmetries are assumed to act as isometries of WCW.

4.3 String-like objects and strings

String like objects obtained as deformations of cosmic strings $X^2 \times Y^2$, where X^2 is minimal surface in M^4 and Y^2 a holomorphic surface of CP_2 are fundamental extremals of Kähler action having string world sheet as M^4 projections. Cosmic strings dominate the primordial cosmology of the TGD Universe and the inflationary period corresponds to the transition to radiation dominated cosmology for which space-time sheets with 4-D M^4 projection dominate.

Also genuine string-like objects emerge from TGD. The conditions that the em charge of modes of induced spinor fields is well-defined requires in the generic case the localization of the modes at 2-D surfaces -string world sheets and possibly also partonic 2-surfaces. This in Minkowskian space-time regions.

Fig. 6. Well-definedness of em charge forces the localization of induced spinor modes to 2-D surfaces in generic situations in Minkowskian regions of space-time surface. <http://tgdtheory.fi/appfigures/fermistring.jpg>

4.4 TGD view of elementary particles

The TGD based view about elementary particles has two key aspects.

1. The space-time correlates of elementary particles are identified as pairs of wormhole contacts with Euclidean signature of metric and having 4-D CP_2 projection. Their throats behave effectively as Kähler magnetic monopoles so that wormhole throats must be connected by Kähler magnetic flux tubes with monopole flux so that closed flux tubes are obtained.
2. At the level of H Fermion number is carried by the modes of the induced spinor field. In space-time regions with Minkowski signature the modes are localized at string world sheets connecting the wormhole contacts.

Fig. 7. TGD view about elementary particles. a) Particle orbit corresponds to a 4-D generalization of a world line or b) with its light-like 3-D boundary (holography). c) Particle world lines have Euclidean signature of the induced metric. d) They can be identified as wormhole contacts. e) The throats of wormhole contacts carry effective Kähler magnetic charges so that wormhole contacts must appear as pairs in order to obtain closed flux tubes. f) Wormhole contacts are accompanied by fermionic strings connecting the throats at the same sheet: the strings do not extend inside the wormhole contacts. <http://tgdtheory.fi/appfigures/elparticletgd.jpg>

Particle interactions involve both stringy and QFT aspects.

1. The boundaries of string world sheets correspond to fundamental fermions. This gives rise to massless propagator lines in generalized Feynman diagrammatics. One can speak of “long” string connecting wormhole contacts and having a hadronic string as a physical counterpart. Long strings should be distinguished from wormhole contacts which due to their superconformal invariance behave like “short” strings with length scale given by CP_2 size, which is 10^4 times longer than Planck scale characterizing strings in string models.
2. Wormhole contact defines basic stringy interaction vertex for fermion-fermion scattering. The propagator is essentially the inverse of the superconformal scaling generator L_0 . Wormhole contacts containing fermion and antifermion at its opposite throats behave like virtual bosons so that one has BFF type vertices typically.
3. In topological sense one has 3-vertices serving as generalizations of 3-vertices of Feynman diagrams. In these vertices 4-D “lines” of generalized Feynman diagrams meet along their 3-D ends. One obtains also the analogs of stringy diagrams but stringy vertices do not have the usual interpretation in terms of particle decays but in terms of propagation of particles along two different routes.

Fig. 8. a) TGD analogs of Feynman and string diagrammatics at the level of space-time topology. b) The 4-D analogs of both string diagrams and QFT diagrams appear but the interpretation of the analogs stringy diagrams is different. <http://tgdtheory.fi/appfigures/tgdgraphs.jpg>

5 About the selection of the action defining the Kähler function of the "world of classical worlds" (WCW)

The proposal is that space-time surfaces correspond to preferred extremals of some action principle, being analogous to Bohr orbits, so that they are almost deterministic. The action for the preferred extremal would define the Kähler function of WCW [K3, K8].

How unique is the choice of the action defining WCW Kähler metric? The problem is that twistor lift strongly suggests the identification of the preferred extremals as 4-D surfaces having 4-D generalization of complex structure and that a large number of general coordinate invariant actions constructible in terms of the induced geometry have the same preferred extremals.

5.1 Could twistor lift fix the choice of the action uniquely?

The twistor lift of TGD [K12] [L12, L13, L14] generalizes the notion of induction to the level of twistor fields and leads to a proposal that the action is obtained by dimensional reduction of the action having as its preferred extremals the counterpart of twistor space of the space-time surface identified as 6-D surface in the product $T(M^4) \times T(CP_2)$ twistor spaces of $T(M^4)$ and $T(CP_2)$ of M^4 and CP_2 . Only M^4 and CP_2 allow a twistor space with Kähler structure [A4] so that TGD would be unique. Dimensional reduction is forced by the condition that the 6-surface has S^2 -bundle structure characterizing twistor spaces and the base space would be the space-time surface.

1. Dimensional reduction of 6-D Kähler action implies that at the space-time level the fundamental action can be identified as the sum of Kähler action and volume term (cosmological constant). Other choices of the action do not look natural in this picture although they would have the same preferred extremals.
2. Preferred extremals are proposed to correspond to minimal surfaces with singularities such that they are also extremals of 4-D Kähler action outside the singularities. The physical analogue are soap films spanned by frames and one can localize the violation of the strict determinism and of strict holography to the frames.
3. The preferred extremal property is realized as the holomorphicity characterizing string world sheets, which generalizes to the 4-D situation. This in turn implies that the preferred extremals are the same for any general coordinate invariant action defined on the induced gauge fields and induced metric apart from possible extremals with vanishing CP_2 Kähler action.

For instance, 4-D Kähler action and Weyl action as the sum of the tensor squares of the components of the Weyl tensor of CP_2 representing quaternionic imaginary units constructed from the Weyl tensor of CP_2 as an analog of gauge field would have the same preferred extremals and only the definition of Kähler function and therefore Kähler metric of WCW would change. One can even consider the possibility that the volume term in the 4-D action could be assigned to the tensor square of the induced metric representing a quaternionic or octonionic real unit.

Action principle does not seem to be unique. On the other hand, the WCW Kähler form and metric should be unique since its existence requires maximal isometries.

Unique action is not the only way to achieve this. One cannot exclude the possibility that the Kähler gauge potential of WCW in the complex coordinates of WCW differs only by a complex gradient of a holomorphic function for different actions so that they would give the same Kähler form for WCW. This gradient is induced by a symplectic transformation of WCW inducing a $U(1)$ gauge transformation. The Kähler metric is the same if the symplectic transformation is an isometry.

Symplectic transformations of WCW could give rise to inequivalent representations of the theory in terms of action at space-time level. Maybe the length scale dependent coupling parameters of an effective action could be interpreted in terms of a choice of WCW Kähler function, which maximally simplifies the computations at a given scale.

1. The 6-D analogues of electroweak action and color action reducing to Kähler action in 4-D case exist. The 6-D analog of Weyl action based on the tensor representation of quaternionic imaginary units does not however exist. One could however consider the possibility that only the base space of twistor space $T(M^4)$ and $T(CP_2)$ have quaternionic structure.
2. Kähler action has a huge vacuum degeneracy, which clearly distinguishes it from other actions. The presence of the volume term removes this degeneracy. However, for minimal surfaces having CP_2 projections, which are Lagrangian manifolds and therefore have a vanishing induced Kähler form, would be preferred extremals according to the proposed definition. For these 4-surfaces, the existence of the generalized complex structure is dubious.

For the electroweak action, the terms corresponding to charged weak bosons eliminate these extremals and one could argue that electroweak action or its sum with the analogue of color action, also proportional Kähler action, defines the more plausible choice. Interestingly, also the neutral part of electroweak action is proportional to Kähler action.

Twistor lift strongly suggests that also M^4 has the analog of Kähler structure. M^8 must be complexified by adding a commuting imaginary unit i . In the E^8 subspace, the Kähler structure of E^4 is defined in the standard sense and it is proposed that this generalizes to M^4 allowing also generalization of the quaternionic structure. M^4 Kähler structure violates Lorentz invariance but could be realized at the level of moduli space of these structures.

The minimal possibility is that the M^4 Kähler form vanishes: one can have a different representation of the Kähler gauge potential for it obtained as generalization of symplectic transformations acting non-trivially in M^4 . The recent picture about the second quantization of spinors of $M^4 \times CP_2$ assumes however non-trivial Kähler structure in M^4 .

5.2 Two paradoxes

TGD view leads to two apparent paradoxes.

1. If the preferred extremals satisfy 4-D generalization of holomorphicity, a very large set of actions gives rise to the same preferred extremals unless there are some additional conditions restricting the number of preferred extremals for a given action.
2. WCW metric has an infinite number of zero modes, which appear as parameters of the metric but do not contribute to the line element. The induced Kähler form depends on these degrees of freedom. The existence of the Kähler metric requires maximal isometries, which suggests that the Kähler metric is uniquely fixed apart from a conformal scaling factor Ω depending on zero modes. This cannot be true: galaxy and elementary particle cannot correspond to the same Kähler metric.

Number theoretical vision and the hierarchy of inclusions of HFFs associated with supersymplectic algebra acting as isometries of WCW provide equivalent realizations of the measurement resolution. This solves these paradoxes and predicts that WCW decomposes into sectors for which Kähler metrics of WCW differ in a natural way.

5.2.1 The hierarchy subalgebras of supersymplectic algebra implies the decomposition of WCW into sectors with different actions

Supersymplectic algebra of $\delta M_+^4 \times CP_2$ is assumed to act as isometries of WCW [L17]. There are also other important algebras but these will not be discussed now.

1. The symplectic algebra A of $\delta M_+^4 \times CP_2$ has the structure of a conformal algebra in the sense that the radial conformal weights with non-negative real part, which is half integer, label the elements of the algebra have an interpretation as conformal weights.

The super symplectic algebra A has an infinite hierarchy of sub-algebras [L17] such that the conformal weights of sub-algebras $A_{n(SS)}$ are integer multiples of the conformal weights of the entire algebra. The superconformal gauge conditions are weakened. Only the subalgebra

$A_{n(SS)}$ and the commutator $[A_{n(SS)}, A]$ annihilate the physical states. Also the corresponding classical Noether charges vanish for allowed space-time surfaces.

This weakening makes sense also for ordinary superconformal algebras and associated Kac-Moody algebras. This hierarchy can be interpreted as a hierarchy symmetry breakings, meaning that sub-algebra $A_{n(SS)}$ acts as genuine dynamical symmetries rather than mere gauge symmetries. It is natural to assume that the super-symplectic algebra A does not affect the coupling parameters of the action.

2. The generators of A correspond to the dynamical quantum degrees of freedom and leave the induced Kähler form invariant. They affect the induced space-time metric but this effect is gravitational and very small for Einsteinian space-time surfaces with 4-D M^4 projection.

The number of dynamical degrees of freedom increases with $n(SS)$. Therefore WCW decomposes into sectors labelled by $n(SS)$ with different numbers of dynamical degrees of freedom so that their Kähler metrics cannot be equivalent and cannot be related by a symplectic isometry. They can correspond to different actions.

5.2.2 Number theoretic vision implies the decomposition of WCW into sectors with different actions

The number theoretical vision leads to the same conclusion as the hierarchy of HFFs. The number theoretic vision of TGD based on $M^8 - H$ duality [L17] predicts a hierarchy with levels labelled by the degrees $n(P)$ of rational polynomials P and corresponding extensions of rationals characterized by Galois groups and by ramified primes defining p-adic length scales.

These sequences allow us to imagine several discrete coupling constant evolutions realized at the level H in terms of action whose coupling parameters depend on the number theoretic parameters.

1. Coupling constant evolution with respect to $n(P)$

The first coupling constant evolution would be with respect to $n(P)$.

1. The coupling constants characterizing action could depend on the degree $n(P)$ of the polynomial defining the space-time region by $M^8 - H$ duality. The complexity of the space-time surface would increase with $n(P)$ and new degrees of freedom would emerge as the number of the rational coefficients of P .
2. This coupling constant evolution could naturally correspond to that assignable to the inclusion hierarchy of hyperfinite factors of type II_1 (HFFs). I have indeed proposed [L17] that the degree $n(P)$ equals to the number $n(\text{braid})$ of braids assignable to HFF for which super symplectic algebra subalgebra $A_{n(SS)}$ with radial conformal weights coming as $n(SS)$ -multiples of those of entire algebra A . One would have $n(P) = n(\text{braid}) = n(SS)$. The number of dynamical degrees of freedom increases with n which just as it increases with $n(P)$ and $n(SS)$.
3. The actions related to different values of $n(P) = n(\text{braid}) = n(SS)$ cannot define the same Kähler metric since the number of allowed space-time surfaces depends on $n(SS)$.

WCW could decompose to sub-WCWs corresponding to different actions, a kind of theory space. These theories would not be equivalent. A possible interpretation would be as a hierarchy of effective field theories.

4. Hierarchies of composite polynomials define sequences of polynomials with increasing values of $n(P)$ such that the order of a polynomial at a given level is divided by those at the lower levels. The proposal is that the inclusion sequences of extensions are realized at quantum level as inclusion hierarchies of hyperfinite factors of type II_1 .

A given inclusion hierarchy corresponds to a sequence $n(SS)_i$ such that $n(SS)_i$ divides $n(SS)_{i+1}$. Therefore the degree of the composite polynomials increases very rapidly. The values of $n(SS)_i$ can be chosen to be primes and these primes correspond to the degrees of so called prime polynomials [L15] so that the decompositions correspond to prime factorizations of integers. The "densest" sequence of this kind would come in powers of 2 as

$n(SS)_i = 2^i$. The corresponding p-adic length scales (assignable to maximal ramified primes for given $n(SS)_i$) are expected to increase roughly exponentially, say as 2^{r2^i} . $r = 1/2$ would give a subset of scales $2^{r/2}$ allowed by the p-adic length scale hypothesis. These transitions would be very rare.

A theory corresponding to a given composite polynomial would contain as sub-theories the theories corresponding to lower polynomial composites. The evolution with respect to $n(SS)$ would correspond to a sequence of phase transitions in which the action genuinely changes. For instance, color confinement could be seen as an example of this phase transition.

5. A subset of p-adic primes allowed by the p-adic length scale hypothesis $p \simeq 2^k$ defining the proposed p-adic length scale hierarchy could relate to n_S changing phase transition. TGD suggests a hierarchy of hadron physics corresponding to a scale hierarchy defined by Mersenne primes and their Gaussian counterparts [K5, K6]). Each of them would be characterized by a confinement phase transition in which n_S and therefore also the action changes.

2. Coupling constant evolutions with respect to ramified primes for a given value of $n(P)$

For a given value of $n(P)$, one could have coupling constant sub-evolutions with respect to the set of ramified primes of P and dimensions $n = h_{eff}/h_0$ of algebraic extensions. The action would only change by U(1) gauge transformation induced by a symplectic isometry of WCW. Coupling parameters could change but the actions would be equivalent.

The choice of the action in an optimal manner in a given scale could be seen as a choice of the most appropriate effective field theory in which radiative corrections would be taken into account. One can interpret the possibility to use a single choice of coupling parameters in terms of quantum criticality.

The range of the p-adic length scales labelled by ramified primes and effective Planck constants h_{eff}/h_0 is finite for a given value of $n(SS)$.

The first coupling constant evolution of this kind corresponds to ramified primes defining p-adic length scales for given $n(SS)$.

1. Ramified primes are factors of the discriminant $D(P)$ of P , which is expressible as a product of non-vanishing root differentials and reduces to a polynomial of the n coefficients of P . Ramified primes define p-adic length scales assignable to the particles in the amplitudes scattering amplitudes defined by zero energy states.

P would represent the space-time surface defining an interaction region in N -particle scattering. The N ramified primes dividing $D(P)$ would characterize the p-adic length scales assignable to these particles. If $D(P)$ reduces to a single ramified prime, one has elementary particle [L15], and the forward scattering amplitude corresponds to the propagator.

This would give rise to a multi-scale p-adic length scale evolution of the amplitudes analogous to the ordinary continuous coupling constant evolution of n-point scattering amplitudes with respect to momentum scales of the particles. This kind of evolutions extend also to evolutions with respect to $n(SS)$.

2. According to [L15], physical constraints require that $n(P)$ and the maximum size of the ramified prime of P correlate.

A given rational polynomial of degree $n(P)$ can be always transformed to a polynomial with integer coefficients. If the integer coefficients are smaller than $n(P)$, there is an upper bound for the ramified primes. This assumption also implies that finite fields become fundamental number fields in number theoretical vision [L15].

3. p-Adic length scale hypothesis [L18] in its basic form states that there exist preferred primes $p \simeq 2^k$ near some powers of 2. A more general hypothesis states that also primes near some powers of 3 possibly also other small primes are preferred physically. The challenge is to understand the origin of these preferred scales.

For polynomials P with a given degree $n(P)$ for which discriminant $D(P)$ is prime, there exists a maximal ramified prime. Numerical calculations suggest that the upper bound depends exponentially on $n(P)$.

Could these maximal ramified primes satisfy the p-adic length scale hypothesis or its generalization? The maximal prime defines a fixed point of coupling constant evolution in accordance with the earlier proposal. For instance, could one think that one has $p \simeq 2^k$, $k = n(SS)$? Each p-adic prime would correspond to a p-adic coupling constant sub-evolution representable in terms of symplectic isometries.

Also the dimension n of the algebraic extension associated with P , which is identified in terms of effective Planck constant $h_{eff}/h_0 = n$ labelling different phases of the ordinary matter behaving like dark matter, could give rise to coupling constant evolution for given $n(SS)$. The range of allowed values of n is finite. Note however that several polynomials of a given degree can correspond to the same dimension of extension.

5.2.3 Number theoretic discretization of WCW and maxima of WCW Kähler function

Number theoretic approach involves a unique discretization of space-time surface and also of WCW. The question is how the points of the discretized WCW correspond to the preferred extremals.

1. The exponents of Kähler function for the maxima of Kähler function, which correspond to the universal preferred extremals, appear in the scattering amplitudes. The number theoretical approach involves a unique discretization of space-time surfaces defining the WCW coordinates of the space-time surface regarded as a point of WCW.

In [L17] it is assumed that these WCW points appearing in the number theoretical discretization correspond to the maxima of the Kähler function. The maxima would depend on the action and would differ for ghd maxima associated with different actions unless they are not related by symplectic WCW isometry.

2. The symplectic transformations of WCW acting as isometries are assumed to be induced by the symplectic transformations of $\delta M_+^4 \times CP_2$ [K3, K2]. As isometries they would naturally permute the maxima with each other.

6 Number theoretic vision of TGD

Physics as number theory vision is complementary to the physics as geometry vision and has developed gradually since 1993. Langlands program is the counterpart of this vision in mathematics [L16].

The notion of p-adic number fields emerged with the motivation coming from the observation that elementary particle mass scales and mass ratios could be understood in terms of the so-called p-adic length scale hypothesis [K7, K4, K1]. The fusion of the various p-adic physics leads to what I call adelic physics [L1, L2]. Later the hypothesis about hierarchy of Planck constants labelling phases of ordinary matter behaving like dark matter emerged [K9, K10, K11, K11].

Eventually this led to that the values of effective Planck constant could be identified as the dimension of an algebraic extension of rationals assignable to polynomials with rational coefficients. This led to the number theoretic vision in which so-called $M^8 - H$ duality [L8, L9] plays a key role. M^8 (actually a complexification of real M^8) is analogous to momentum space so that the duality generalizes momentum position duality for point-like particles. M^8 has an interpretation as complexified octonions.

The dynamics of 4-surfaces in M^8 is coded by polynomials with rational coefficients, whose roots define mass shells H^3 of $M^4 \subset M^8$. It has turned out that the polynomials satisfy stringent additional conditions and one can speak of number theoretic holography [L15, L16]. Also the ordinary $3 \rightarrow 4$ holography is needed to assign 4-surfaces with these 3-D mass shells. The number theoretic dynamics is based on the condition that the normal space of the 4-surface in M^8 is associative (quaternionic) and contains a commutative complex sub-space. This makes it possible to assign to this surface space-time surface in $H = M^4 \times CP_2$.

At the level of H the space-time surfaces are by holography preferred extremals and are assumed to be determined by the twistor lift of TGD [K12] giving rise to an action which is sum

of the Kähler action and volume term. The preferred extremals would be minimal surfaces analogous to soap films spanned by frames. Outside frames they would be simultaneous extremals of the Kähler action, which requires a generalization of the holomorphy characterizing string world sheets.

In the following only p-adic numbers and hierarchy of Planck constants will be discussed.

6.1 p-Adic numbers and TGD

6.1.1 p-Adic number fields

p-Adic numbers (p is prime: 2, 3, 5, ...) can be regarded as a completion of the rational numbers using a norm, which is different from the ordinary norm of real numbers [A1]. p-Adic numbers are representable as power expansion of the prime number p of form

$$x = \sum_{k \geq k_0} x(k)p^k, \quad x(k) = 0, \dots, p-1. \quad (6.1)$$

The norm of a p-adic number is given by

$$|x| = p^{-k_0(x)}. \quad (6.2)$$

Here $k_0(x)$ is the lowest power in the expansion of the p-adic number. The norm differs drastically from the norm of the ordinary real numbers since it depends on the lowest binary digit of the p-adic number only. Arbitrarily high powers in the expansion are possible since the norm of the p-adic number is finite also for numbers, which are infinite with respect to the ordinary norm. A convenient representation for p-adic numbers is in the form

$$x = p^{k_0} \varepsilon(x), \quad (6.3)$$

where $\varepsilon(x) = k + \dots$ with $0 < k < p$, is p-adic number with unit norm and analogous to the phase factor $\exp(i\phi)$ of a complex number.

The distance function $d(x, y) = |x - y|_p$ defined by the p-adic norm possesses a very general property called ultra-metricity:

$$d(x, z) \leq \max\{d(x, y), d(y, z)\}. \quad (6.4)$$

The properties of the distance function make it possible to decompose R_p into a union of disjoint sets using the criterion that x and y belong to same class if the distance between x and y satisfies the condition

$$d(x, y) \leq D. \quad (6.5)$$

This division of the metric space into classes has following properties:

1. Distances between the members of two different classes X and Y do not depend on the choice of points x and y inside classes. One can therefore speak about distance function between classes.
2. Distances of points x and y inside single class are smaller than distances between different classes.
3. Classes form a hierarchical tree.

Notice that the concept of the ultra-metricity emerged in physics from the models for spin glasses and is believed to have also applications in biology [B4]. The emergence of p-adic topology as the topology of the effective space-time would make ultra-metricity property basic feature of physics.

6.1.2 Canonical correspondence between p-adic and real numbers

The basic challenge encountered by p-adic physicist is how to map the predictions of the p-adic physics to real numbers. p-Adic probabilities provide a basic example in this respect. Identification via common rationals and canonical identification and its variants have turned out to play a key role in this respect.

1. Basic form of the canonical identification

There exists a natural continuous map $I : R_p \rightarrow R_+$ from p-adic numbers to non-negative real numbers given by the “binary” expansion of the real number for $x \in R$ and $y \in R_p$ this correspondence reads

$$\begin{aligned} y &= \sum_{k>N} y_k p^k \rightarrow x = \sum_{k<N} y_k p^{-k} , \\ y_k &\in \{0, 1, \dots, p-1\} . \end{aligned} \quad (6.6)$$

This map is continuous as one easily finds out. There is however a little difficulty associated with the definition of the inverse map since the binary expansion like also decimal expansion is not unique ($1 = 0.999\dots$) for the real numbers x , which allow binary expansion with finite number of binary digits

$$\begin{aligned} x &= \sum_{k=N_0}^N x_k p^{-k} , \\ x &= \sum_{k=N_0}^{N-1} x_k p^{-k} + (x_N - 1)p^{-N} + (p-1)p^{-N-1} \sum_{k=0,\dots} p^{-k} . \end{aligned} \quad (6.7)$$

The p-adic images associated with these expansions are different

$$\begin{aligned} y_1 &= \sum_{k=N_0}^N x_k p^k , \\ y_2 &= \sum_{k=N_0}^{N-1} x_k p^k + (x_N - 1)p^N + (p-1)p^{N+1} \sum_{k=0,\dots} p^k \\ &= y_1 + (x_N - 1)p^N - p^{N+1} , \end{aligned} \quad (6.8)$$

so that the inverse map is either two-valued for p-adic numbers having expansion with finite binary digits or single valued and discontinuous and non-surjective if one makes binary expansion unique by choosing the one with finite binary digits. The finite binary digit expansion is a natural choice since in the numerical work one always must use a binary cutoff on the real axis.

2. The topology induced by canonical identification

The topology induced by the canonical identification in the set of positive real numbers differs from the ordinary topology. The difference is easily understood by interpreting the p-adic norm as a norm in the set of the real numbers. The norm is constant in each interval $[p^k, p^{k+1})$ (see **Fig. 6.1.2**) and is equal to the usual real norm at the points $x = p^k$: the usual linear norm is replaced with a piecewise constant norm. This means that p-adic topology is coarser than the usual real topology and the higher the value of p is, the coarser the resulting topology is above a given length scale. This hierarchical ordering of the p-adic topologies will be a central feature as far as the proposed applications of the p-adic numbers are considered.

Ordinary continuity implies p-adic continuity since the norm induced from the p-adic topology is rougher than the ordinary norm. p-Adic continuity implies ordinary continuity from right as is

clear already from the properties of the p-adic norm (the graph of the norm is indeed continuous from right). This feature is one clear signature of the p-adic topology.

Fig. 14. The real norm induced by canonical identification from 2-adic norm. <http://tgdtheory.fi/appfigures/norm.png>

The linear structure of the p-adic numbers induces a corresponding structure in the set of the non-negative real numbers and p-adic linearity in general differs from the ordinary concept of linearity. For example, p-adic sum is equal to real sum only provided the summands have no common binary digits. Furthermore, the condition $x +_p y < \max\{x, y\}$ holds in general for the p-adic sum of the real numbers. p-Adic multiplication is equivalent with the ordinary multiplication only provided that either of the members of the product is power of p . Moreover one has $x \times_p y < x \times y$ in general. The p-Adic negative -1_p associated with p-adic unit 1 is given by $(-1)_p = \sum_k (p-1)p^k$ and defines p-adic negative for each real number x . An interesting possibility is that p-adic linearity might replace the ordinary linearity in some strongly nonlinear systems so these systems would look simple in the p-adic topology.

These results suggest that canonical identification is involved with some deeper mathematical structure. The following inequalities hold true:

$$\begin{aligned} (x + y)_R &\leq x_R + y_R , \\ |x|_p |y|_R &\leq (xy)_R \leq x_R y_R , \end{aligned} \quad (6.9)$$

where $|x|_p$ denotes p-adic norm. These inequalities can be generalized to the case of $(R_p)^n$ (a linear vector space over the p-adic numbers).

$$\begin{aligned} (x + y)_R &\leq x_R + y_R , \\ |\lambda|_p |y|_R &\leq (\lambda y)_R \leq \lambda_R y_R , \end{aligned} \quad (6.10)$$

where the norm of the vector $x \in T_p^n$ is defined in some manner. The case of Euclidian space suggests the definition

$$(x_R)^2 = \left(\sum_n x_n^2 \right)_R . \quad (6.11)$$

These inequalities resemble those satisfied by the vector norm. The only difference is the failure of linearity in the sense that the norm of a scaled vector is not obtained by scaling the norm of the original vector. Ordinary situation prevails only if the scaling corresponds to a power of p .

These observations suggests that the concept of a normed space or Banach space might have a generalization and physically the generalization might apply to the description of some non-linear systems. The nonlinearity would be concentrated in the nonlinear behavior of the norm under scaling.

3. Modified form of the canonical identification

The original form of the canonical identification is continuous but does not respect symmetries even approximately. This led to a search of variants which would do better in this respect. The modification of the canonical identification applying to rationals only and given by

$$I_Q(q = p^k \times \frac{r}{s}) = p^k \times \frac{I(r)}{I(s)} \quad (6.12)$$

is uniquely defined for rationals, maps rationals to rationals, has also a symmetry under exchange of target and domain. This map reduces to a direct identification of rationals for $0 \leq r < p$ and $0 \leq s < p$. It has turned out that it is this map which most naturally appears in the applications. The map is obviously continuous locally since p-adically small modifications of r and s mean small modifications of the real counterparts.

Canonical identification is in a key role in the successful predictions of the elementary particle masses. The predictions for the light elementary particle masses are within extreme accuracy same for I and I_Q but I_Q is theoretically preferred since the real probabilities obtained from p-adic ones by I_Q sum up to one in p-adic thermodynamics.

4. Generalization of number concept and notion of embedding space

TGD forces an extension of number concept: roughly a fusion of reals and various p-adic number fields along common rationals is in question. This induces a similar fusion of real and p-adic embedding spaces. Since finite p-adic numbers correspond always to non-negative reals n -dimensional space R^n must be covered by 2^n copies of the p-adic variant R_p^n of R^n each of which projects to a copy of R_+^n (four quadrants in the case of plane). The common points of p-adic and real embedding spaces are rational points and most p-adic points are at real infinity.

Real numbers and various algebraic extensions of p-adic number fields are thus glued together along common rationals and also numbers in algebraic extension of rationals whose number belong to the algebraic extension of p-adic numbers. This gives rise to a book like structure with rationals and various algebraic extensions of rationals taking the role of the back of the book. Note that Neper number is exceptional in the sense that it is algebraic number in p-adic number field Q_p satisfying $e^p \bmod p = 1$.

Fig. 15. Various number fields combine to form a book like structure. <http://tgdtheory.fi/appfigures/book.jpg>

For a given p-adic space-time sheet most points are literally infinite as real points and the projection to the real embedding space consists of a discrete set of rational points: the interpretation in terms of the unavoidable discreteness of the physical representations of cognition is natural. Purely local p-adic physics implies real p-adic fractality and thus long range correlations for the real space-time surfaces having enough common points with this projection.

p-Adic fractality means that M^4 projections for the rational points of space-time surface X^4 are related by a direct identification whereas CP_2 coordinates of X^4 at these points are related by I , I_Q or some of its variants implying long range correlates for CP_2 coordinates. Since only a discrete set of points are related in this manner, both real and p-adic field equations can be satisfied and there are no problems with symmetries. p-Adic effective topology is expected to be a good approximation only within some length scale range which means infrared and UV cutoffs. Also multi-p-fractality is possible.

6.1.3 The notion of p-adic manifold

The notion of p-adic manifold is needed in order to fuse real physics and various p-adic physics to a larger structure which suggests that real and p-adic number fields should be glued together along common rationals bringing in mind adeles. The notion is problematic because p-adic topology is totally disconnected implying that p-adic balls are either disjoint or nested so that ordinary definition of manifold using p-adic chart maps fails. A cure is suggested to be based on chart maps from p-adics to reals rather than to p-adics (see the appendix of the book)

The chart maps are interpreted as cognitive maps, "thought bubbles".

Fig. 16. The basic idea between p-adic manifold. <http://tgdtheory.fi/appfigures/padmanifold.jpg>

There are some problems.

1. Canonical identification does not respect symmetries since it does not commute with second pinary cutoff so that only a discrete set of rational points is mapped to their real counterparts by chart map arithmetic operations which requires pinary cutoff below which chart map takes rationals to rationals so that commutativity with arithmetics and symmetries is achieved in finite resolution: above the cutoff canonical identification is used
2. Canonical identification is continuous but does not map smooth p-adic surfaces to smooth real surfaces requiring second pinary cutoff so that only a discrete set of rational points is mapped to their real counterparts by chart map requiring completion of the image to smooth preferred extremal of Kähler action so that chart map is not unique in accordance with finite measurement resolution

3. Canonical identification violates general coordinate invariance of chart map: (cognition-induced symmetry breaking) minimized if p-adic manifold structure is induced from that for p-adic embedding space with chart maps to real embedding space and assuming preferred coordinates made possible by isometries of embedding space: one however obtains several inequivalent p-adic manifold structures depending on the choice of coordinates: these cognitive representations are not equivalent.

6.2 Hierarchy of Planck constants and dark matter hierarchy

Hierarchy of Planck constants was motivated by the “impossible” quantal effects of ELF em fields on vertebrate cyclotron energies $E = hf = \hbar \times eB/m$ are above thermal energy is possible only if \hbar has value much larger than its standard value. Also Nottale’s finding that planetary orbits might be understood as Bohr orbits for a gigantic gravitational Planck constant.

Hierarchy of Planck constant would mean that the values of Planck constant come as integer multiples of ordinary Planck constant: $h_{eff} = n \times h$. The particles at magnetic flux tubes characterized by h_{eff} would correspond to dark matter which would be invisible in the sense that only particle with same value of h_{eff} appear in the same vertex of Feynman diagram.

Hierarchy of Planck constants would be due to the non-determinism of the Kähler action predicting huge vacuum degeneracy allowing all space-time surfaces which are sub-manifolds of any $M^4 \times Y^2$, where Y^2 is Lagrangian sub-manifold of CP_2 . For a given Y^2 one obtains new manifolds Y^2 by applying symplectic transformations of CP_2 .

Non-determinism would mean that the 3-surface at the ends of causal diamond (CD) can be connected by several space-time surfaces carrying same conserved Kähler charges and having same values of Kähler action. Conformal symmetries defined by Kac-Moody algebra associated with the embedding space isometries could act as gauge transformations and respect the light-likeness property of partonic orbits at which the signature of the induced metric changes from Minkowskian to Euclidian (Minkowskian space-time region transforms to wormhole contact say). The number of conformal equivalence classes of these surfaces could be finite number n and define discrete physical degree of freedom and one would have $h_{eff} = n \times h$. This degeneracy would mean “second quantization” for the sheets of n-furcation: not only one but several sheets can be realized.

This relates also to quantum criticality postulated to be the basic characteristics of the dynamics of quantum TGD. Quantum criticalities would correspond to an infinite fractal hierarchy of broken conformal symmetries defined by sub-algebras of conformal algebra with conformal weights coming as integer multiples of n . This leads also to connections with quantum criticality and hierarchy of broken conformal symmetries, p-adicity, and negentropic entanglement which by consistency with standard quantum measurement theory would be described in terms of density matrix proportional $n \times n$ identity matrix and being due to unitary entanglement coefficients (typical for quantum computing systems).

Formally the situation could be described by regarding space-time surfaces as surfaces in singular n-fold singular coverings of embedding space. A stronger assumption would be that they are expressible as products of n_1 -fold covering of M^4 and n_2 -fold covering of CP_2 meaning analogy with multi-sheeted Riemann surfaces and that M^4 coordinates are n_1 -valued functions and CP_2 coordinates n_2 -valued functions of space-time coordinates for $n = n_1 \times n_2$. These singular coverings of embedding space form a book like structure with singularities of the coverings localizable at the boundaries of causal diamonds defining the back of the book like structure.

Fig. 17. Hierarchy of Planck constants. <http://tgdtheory.fi/appfigures/planckhierarchy.jpg>

6.3 $M^8 - H$ duality as it is towards the end of 2021

The view of $M^8 - H$ duality (see Appendix ??) has changed considerably towards the end 2021 [L12] after the realization that this duality is the TGD counterpart of momentum position duality of wave mechanics, which is lost in QFTs. Therefore M^8 and also space-time surface is analogous to momentum space. This forced us to give up the original simple identification of the points $M^4 \subset M^4 \times E^4 = M^8$ and of $M^4 \times CP_2$ so that it respects Uncertainty Principle (UP).

The first improved guess for the duality map was the replacement with the inversion $p^k \rightarrow m^k = \hbar_{eff} p^k / p^2$ conforming in spirit with UP but turned out to be too naive.

The improved form [L12] of the $M^8 - H$ duality map takes mass shells $p^2 = m^2$ of $M^4 \subset M^8$ to cds with size $L(m) = \hbar_{eff}/m$ with a common center. The slicing by mass shells is mapped to a Russian doll like slicing by cds. Therefore would be no CDs in M^8 contrary to what I believed first.

Quantum classical correspondence (QCC) inspires the proposal that the point $p^k \in M^8$ is mapped to a geodesic line corresponding to momentum p^k starting from the common center of cds. Its intersection with the opposite boundary of cd with size $L(m)$ defines the image point. This is not yet quite enough to satisfy UP but the additional details [L12] are not needed in the sequel.

The 6-D brane-like special solutions in M^8 are of special interest in the TGD inspired theory of consciousness. They have an M^4 projection which is $E = E_n$ 3-ball. Here E_n is a root of the real polynomial P defining $X^4 \subset M_c^8$ (M^8 is complexified to M_c^8) as a "root" of its octonionic continuation [L8, L9]. E_n has an interpretation as energy, which can be complex. The original interpretation was as moment of time. For this interpretation, $M^8 - H$ duality would be a linear identification and these hyper planes would be mapped to hyperplanes in $M^4 \subset H$. This motivated the term "very special moment in the life of self" for the image of the $E = E_n$ section of $X^4 \subset M^8$ [L5]. This notion does not make sense at the level M^8 anymore.

The modified $M^8 - H$ duality forces us to modify the original interpretation [L12]. The point $(E_n, p = 0)$ is mapped $(t_n = \hbar_{eff}/E_n, 0)$. The momenta (E_n, p) in $E = E_n$ plane are mapped to the boundary of cd and correspond to a continuous time interval at the boundary of CD: "very special moment" becomes a "very special time interval".

The quantum state however corresponds to a set of points corresponding to quark momenta, which belong to a cognitive representation and are therefore algebraic integers in the extension determined by the polynomial. These active points in E_n are mapped to a discrete set at the boundary of cd(m). A "very special moment" is replaced with a sequence of "very special moments".

So called Galois confinement [L10] forces the total momenta for bound states of quarks and antiquarks to be rational integers invariant under Galois group of extension of rationals determined by the polynomial P [L12]. These states correspond to states at boundaries of sub-CDs so that one obtains a hierarchy. Galois confinement provides a universal number theoretic mechanism for the formation of bound states.

7 Zero energy ontology (ZEO)

ZEO is implied by the holography forced in the TGD framework by general coordinate invariance.

7.1 Basic motivations and ideas of ZEO

The following gives a brief summary of ZEO [L7] [K13].

1. In ZEO quantum states are not 3-dimensional but superpositions of 4-dimensional deterministic time evolutions connecting ordinary initial 3-dimensional states. By holography they are equivalent to pairs of ordinary 3-D states identified as initial and final states of time evolution. One can say that in the TGD framework general coordinate invariance implies holography and the slight failure of its determinism in turn forces ZEO.

Quantum jumps replace this state with a new one: a superposition of deterministic time evolutions is replaced with a new superposition. Classical determinism of individual time evolution is not violated and this solves the basic paradox of quantum measurement theory. There are two kinds of quantum jumps: ordinary ("big") state function reductions (BSFRs) changing the arrow of time and "small" state function reductions (SSFRs) (weak measurements) preserving it and giving rise to the analog of Zeno effect [L7].

2. To avoid getting totally confused it is good to emphasize some aspects of ZEO.
 - (a) ZEO does not mean that physical states in the usual 3-D sense as snapshots of time evolution would have zero energy state pairs defining zero energy states as initial and

final states have same conserved quantities such as energy. Conservation implies that one can adopt the conventions that the values of conserved quantities are opposite for these states so that their sum vanishes: one can think that incoming and outgoing particles come from geometric past and future is the picture used in quantum field theories.

- (b) ZEO means two times: subjective time as sequence of quantum jumps and geometric time as space-time coordinate. These times are identifiable but are strongly correlated.
3. In BSFRs the arrow of time is changed and the time evolution in the final state occurs backwards with respect to the time of the external observer. BSFRs can occur in all scales since TGD predicts a hierarchy of effective Planck constants with arbitrarily large values. There is empirical support for BSFRs.
- (a) The findings of Mineev et al [L3] in atomic scale can be explained by the same mechanism [L3]. In BSFR a final zero energy state as a superposition of classical deterministic time evolutions emerges and for an observer with a standard arrow of time looks like a superposition of deterministic smooth time evolutions leading to the final state. Interestingly, once this evolution has started, it cannot be stopped unless one changes the stimulus signal inducing the evolution in which case the process does not lead to anywhere: the interpretation would be that BSFR back to the initial state occurs!
 - (b) Libets' experiments about active aspects of consciousness [J1] can be understood. Subject person raises his finger and neural activity starts before the conscious decision to do so. In the physicalistic framework it is thought to lead to raising of the finger. The problem with the explanation is that the activity beginning .5 seconds earlier seems to be dissipation with a reversed arrow of time: from chaotic and disordered to ordered at around .15 seconds. ZEO explanation is that macroscopic quantum jump occurred and generated a signal proceeding backwards in time and generated neural activity and dissipated to randomness.
 - (c) Earthquakes involve a strange anomaly: they are preceded by ELF radiation. One would expect that they generate ELF radiation. The identification as BSFR would explain the anomaly [L4]. In biology the reversal of the arrow of time would occur routinely and be a central element of biological self-organization, in particular self-organized quantum criticality (see [L6, L20]).

7.2 Some implications of ZEO

ZEO has profound implications for understanding self-organization and self-organized quantum criticality in terms of dissipation with non-standard arrow of time looking like generation of structures [L6, L20]. ZEO could also allow understanding of what planned actions - like realizing the experiment under consideration - could be.

1. Second law in the standard sense does not favor - perhaps even not allow - realization of planned actions. ZEO forces a generalization of thermodynamics: dissipation with a non-standard arrow of time for a subsystem would look like self-organization and planned action and its realization.

Could most if not all planned action be like this - induced by BSFR in the geometric future and only apparently planned? There would be however the experience of planning and realizing induced by the signals from geometric future by a higher level in the hierarchy of conscious entities predicted by TGD! In long time scales we would be realizing our fates or wishes of higher level conscious entities rather than agents with completely free will.

2. The notion of magnetic body (MB) serving as a boss of ordinary matter would be central. MB carries dark matter as $h_{eff} = nh_0$ phases of ordinary matter with n serving as a measure for algebraic complexity of extension of rationals as its dimension and defining a kind of universal IQ. There is a hierarchy of these phases and MBs labelled by extension of rationals and the value of n .

MBs would form a hierarchy of bosses - a realization for master slave hierarchy. Ordinary matter would be at the bottom and its coherent behavior would be induced from quantum

coherence at higher levels. BSFR for higher level MB would give rise to what looks like planned actions and experienced as planned action at the lower levels of hierarchy. One could speak of planned actions inducing a cascade of planned actions in shorter time scales and eventually proceeding to atomic level.

8 Some notions relevant to TGD inspired consciousness and quantum biology

Below some notions relevant to TGD inspired theory of consciousness and quantum biology.

8.1 The notion of magnetic body

Topological field quantization inspires the notion of field body about which magnetic body is especially important example and plays key role in TGD inspired quantum biology and consciousness theory. This is a crucial departure from the Maxwellian view. Magnetic body brings in third level to the description of living system as a system interacting strongly with environment. Magnetic body would serve as an intentional agent using biological body as a motor instrument and sensory receptor. EEG would communicate the information from biological body to magnetic body and Libet's findings from time delays of consciousness support this view.

The following pictures illustrate the notion of magnetic body and its dynamics relevant for quantum biology in TGD Universe.

Fig. 18. Magnetic body associated with dipole field. <http://tgdtheory.fi/appfigures/fluxquant.jpg>

Fig. 19. Illustration of the reconnection by magnetic flux loops. <http://tgdtheory.fi/appfigures/reconnect1.jpg>

Fig. 20. Illustration of the reconnection by flux tubes connecting pairs of molecules. <http://tgdtheory.fi/appfigures/reconnect2.jpg>

Fig. 21. Flux tube dynamics. a) Reconnection making possible magnetic body to "recognize" the presence of another magnetic body, b) braiding, knotting and linking of flux tubes making possible topological quantum computation, c) contraction of flux tube in phase transition reducing the value of h_{eff} allowing two molecules to find each other in dense molecular soup. <http://tgdtheory.fi/appfigures/fluxtubedynamics.jpg>

8.2 Number theoretic entropy and negentropic entanglement

TGD inspired theory of consciousness relies heavily p-Adic norm allows an to define the notion of Shannon entropy for rational probabilities (and even those in algebraic extension of rationals) by replacing the argument of logarithm of probability with its p-adic norm. The resulting entropy can be negative and the interpretation is that number theoretic entanglement entropy defined by this formula for the p-adic prime minimizing its value serves as a measure for conscious information. This negentropy characterizes two-particle system and has nothing to do with the formal negative negentropy assignable to thermodynamic entropy characterizing single particle. Negentropy Maximization Principle (NMP) implies that number theoretic negentropy increases during evolution by quantum jumps. The condition that NMP is consistent with the standard quantum measurement theory requires that negentropic entanglement has a density matrix proportional to unit matrix so that in 2-particle case the entanglement matrix is unitary.

Fig. 22. Schrödinger cat is neither dead or alive. For negentropic entanglement this state would be stable. <http://tgdtheory.fi/appfigures/cat.jpg>

8.3 Life as something residing in the intersection of reality and p-adicities

In TGD inspired theory of consciousness p-adic space-time sheets correspond to space-time correlates for thoughts and intentions. The intersections of real and p-adic preferred extremals consist

of points whose coordinates are rational or belong to some extension of rational numbers in preferred embedding space coordinates. They would correspond to the intersection of reality and various p -adicities representing the “mind stuff” of Descartes. There is temptation to assign life to the intersection of realities and p -adicities. The discretization of the chart map assigning to real space-time surface its p -adic counterpart would reflect finite cognitive resolution.

At the level of “world of classical worlds” (WCW) the intersection of reality and various p -adicities would correspond to space-time surfaces (or possibly partonic 2-surfaces) representable in terms of rational functions with polynomial coefficients with are rational or belong to algebraic extension of rationals.

The quantum jump replacing real space-time sheet with p -adic one (vice versa) would correspond to a buildup of cognitive representation (realization of intentional action).

Fig. 23. The quantum jump replacing real space-time surface with corresponding p -adic manifold can be interpreted as formation of thought, cognitive representation. Its reversal would correspond to a transformation of intention to action. <http://tgdtheory.fi/appfigures/padictoreal.jpg>

8.4 Sharing of mental images

The 3-surfaces serving as correlates for sub-selves can topologically condense to disjoint large space-time sheets representing selves. These 3-surfaces can also have flux tube connections and this makes possible entanglement of sub-selves, which unentangled in the resolution defined by the size of sub-selves. The interpretation for this negentropic entanglement would be in terms of sharing of mental images. This would mean that contents of consciousness are not completely private as assumed in neuroscience.

Fig. 24. Sharing of mental images by entanglement of subselves made possible by flux tube connections between topologically condensed space-time sheets associated with mental images. <http://tgdtheory.fi/appfigures/sharing.jpg>

8.5 Time mirror mechanism

Zero energy ontology (ZEO) is crucial part of both TGD and TGD inspired consciousness and leads to the understanding of the relationship between geometric time and experience time and how the arrow of psychological time emerges. One of the basic predictions is the possibility of negative energy signals propagating backwards in geometric time and having the property that entropy basically associated with subjective time grows in reversed direction of geometric time. Negative energy signals inspire time mirror mechanism (see **Fig.** <http://tgdtheory.fi/appfigures/timemirror.jpg> or **Fig. 24** in the appendix of this book) providing mechanisms of both memory recall, realization of intentional action initiating action already in geometric past, and remote metabolism. What happens that negative energy signal travels to past and is reflected as positive energy signal and returns to the sender. This process works also in the reverse time direction.

Fig. 25. Zero energy ontology allows time mirror mechanism as a mechanism of memory recall. Essentially “seeing” in time direction is in question. <http://tgdtheory.fi/appfigures/timemirror.jpg>

Acknowledgements: I am grateful for Dainis Zeps for enlightening discussions concerning CP_2 geometry.

REFERENCES

Mathematics

- [A1] Shafarevich IR Borevich ZI. *Number Theory*. Academic Press, 1966.
- [A2] Pope CN. Eigenfunctions and $Spin^c$ Structures on CP_2 , 1980.

- [A3] Hanson J Eguchi T, Gilkey B. *Phys Rep*, 66, 1980.
- [A4] N. Hitchin. Kählerian twistor spaces. *Proc London Math Soc*, 8(43):133–151, 1981.. Available at: <https://tinyurl.com/pb8zpqo>.
- [A5] Helgason S. *Differential Geometry and Symmetric Spaces*. Academic Press, New York, 1962.

Theoretical Physics

- [B1] Zee A. *The Unity of Forces in the Universe*. World Sci Press, Singapore, 1982.
- [B2] Drell S Björken J. *Relativistic Quantum Fields*. Mc Graw-Hill, New York, 1965.
- [B3] Mineev ZK et al. To catch and reverse a quantum jump mid-flight, 2019. Available at: <https://arxiv.org/abs/1803.00545>.
- [B4] Parisi G. *Field Theory, Disorder and Simulations*. World Scientific, 1992.
- [B5] Huang K. *Quarks, Leptons & Gauge Fields*. World Scientific, 1982.

Neuroscience and Consciousness

- [J1] Libet B. Readiness potentials preceding unrestricted spontaneous and preplanned voluntary acts, 1982. Available at: <https://tinyurl.com/jqp1>. See also the article *Libet's Research on Timing of Conscious Intention to Act: A Commentary* of Stanley Klein at <https://tinyurl.com/jqp1>.

Books related to TGD

- [K1] Pitkänen M. Construction of elementary particle vacuum functionals. In *p-Adic Physics*. Available at: <https://tgdtheory.fi/pdfpool/elvafu.pdf>, 2006.
- [K2] Pitkänen M. Construction of WCW Kähler Geometry from Symmetry Principles. In *Quantum Physics as Infinite-Dimensional Geometry*. Available at: <https://tgdtheory.fi/pdfpool/compl1.pdf>, 2006.
- [K3] Pitkänen M. Identification of the WCW Kähler Function. In *Quantum Physics as Infinite-Dimensional Geometry*. Available at: <https://tgdtheory.fi/pdfpool/kahler.pdf>, 2006.
- [K4] Pitkänen M. Massless states and particle massivation. In *p-Adic Physics*. Available at: <https://tgdtheory.fi/pdfpool/mless.pdf>, 2006.
- [K5] Pitkänen M. New Particle Physics Predicted by TGD: Part I. In *p-Adic Physics*. Available at: <https://tgdtheory.fi/pdfpool/mass4.pdf>, 2006.
- [K6] Pitkänen M. New Particle Physics Predicted by TGD: Part II. In *p-Adic Physics*. Available at: <https://tgdtheory.fi/pdfpool/mass5.pdf>, 2006.
- [K7] Pitkänen M. *p-Adic length Scale Hypothesis*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/padphys.html>, 2013.
- [K8] Pitkänen M. Recent View about Kähler Geometry and Spin Structure of WCW . In *Quantum Physics as Infinite-Dimensional Geometry*. Available at: <https://tgdtheory.fi/pdfpool/wcwnew.pdf>, 2014.
- [K9] Pitkänen M. Criticality and dark matter: part I. In *Hyper-finite Factors and Dark Matter Hierarchy: Part I*. Available at: <https://tgdtheory.fi/pdfpool/qcritdark1.pdf>, 2019.
- [K10] Pitkänen M. Criticality and dark matter: part II. In *Hyper-finite Factors and Dark Matter Hierarchy: Part I*. Available at: <https://tgdtheory.fi/pdfpool/qcritdark2.pdf>, 2019.

- [K11] Pitkänen M. Criticality and dark matter: part III. In *Hyper-finite Factors and Dark Matter Hierarchy: Part I*. Available at: <https://tgdtheory.fi/pdfpool/qcritdark3.pdf>, 2019.
- [K12] Pitkänen M. Some questions related to the twistor lift of TGD. In *Towards M-Matrix: Part II*. Available at: <https://tgdtheory.fi/pdfpool/twistquestions.pdf>, 2019.
- [K13] Pitkänen M. Zero Energy Ontology. In *Towards M-Matrix: part I*. Available at: <https://tgdtheory.fi/pdfpool/ZEO.pdf>, 2021.

Articles about TGD

- [L1] Pitkänen M. Philosophy of Adelic Physics. In *Trends and Mathematical Methods in Interdisciplinary Mathematical Sciences*, pages 241–319. Springer. Available at: https://link.springer.com/chapter/10.1007/978-3-319-55612-3_11, 2017.
- [L2] Pitkänen M. Philosophy of Adelic Physics. Available at: https://tgdtheory.fi/public_html/articles/adelephysics.pdf, 2017.
- [L3] Pitkänen M. Copenhagen interpretation dead: long live ZEO based quantum measurement theory! Available at: https://tgdtheory.fi/public_html/articles/Bohrdead.pdf, 2019.
- [L4] Pitkänen M. Earthquakes and volcanic eruptions as macroscopic quantum jumps in zero energy ontology. Available at: https://tgdtheory.fi/public_html/articles/earthquakes.pdf, 2019.
- [L5] Pitkänen M. $M^8 - H$ duality and consciousness. Available at: https://tgdtheory.fi/public_html/articles/M8Hconsc.pdf, 2019.
- [L6] Pitkänen M. Quantum self-organization by h_{eff} changing phase transitions. Available at: https://tgdtheory.fi/public_html/articles/heffselforg.pdf, 2019.
- [L7] Pitkänen M. Some comments related to Zero Energy Ontology (ZEO). Available at: https://tgdtheory.fi/public_html/articles/zeoquestions.pdf, 2019.
- [L8] Pitkänen M. A critical re-examination of $M^8 - H$ duality hypothesis: part I. Available at: https://tgdtheory.fi/public_html/articles/M8H1.pdf, 2020.
- [L9] Pitkänen M. A critical re-examination of $M^8 - H$ duality hypothesis: part II. Available at: https://tgdtheory.fi/public_html/articles/M8H2.pdf, 2020.
- [L10] Pitkänen M. About the role of Galois groups in TGD framework. https://tgdtheory.fi/public_html/articles/GaloisTGD.pdf, 2021.
- [L11] Pitkänen M. Some questions concerning zero energy ontology. https://tgdtheory.fi/public_html/articles/zeonew.pdf, 2021.
- [L12] Pitkänen M. TGD as it is towards the end of 2021. https://tgdtheory.fi/public_html/articles/TGD2021.pdf, 2021.
- [L13] Pitkänen M. About TGD counterparts of twistor amplitudes: part I. https://tgdtheory.fi/public_html/articles/twisttgd1.pdf, 2022.
- [L14] Pitkänen M. About TGD counterparts of twistor amplitudes: part II. https://tgdtheory.fi/public_html/articles/twisttgd2.pdf, 2022.
- [L15] Pitkänen M. Finite Fields and TGD. https://tgdtheory.fi/public_html/articles/finitefieldsTGD.pdf, 2022.
- [L16] Pitkänen M. Some New Ideas Related to Langlands Program *viz.* TGD. https://tgdtheory.fi/public_html/articles/Langlands2022.pdf, 2022.
- [L17] Pitkänen M. Trying to fuse the basic mathematical ideas of quantum TGD to a single coherent whole. https://tgdtheory.fi/public_html/articles/fusionTGD.pdf, 2022.

-
- [L18] Pitkänen M. Two objections against p-adic thermodynamics and their resolution. https://tgdtheory.fi/public_html/articles/padmass2022.pdf, 2022.
- [L19] Pitkänen M. Reduction of standard model structure to CP_2 geometry and other key ideas of TGD. https://tgdtheory.fi/public_html/articles/cp2etc.pdf, 2023.
- [L20] Pitkänen M and Rastmanesh R. Homeostasis as self-organized quantum criticality. Available at: https://tgdtheory.fi/public_html/articles/SP.pdf, 2020.