

Hierarchies of Conformal Symmetry Breakings, Quantum Criticalities, Planck Constants, and of Hyper-Finite factors

M. Pitkänen

Email: matpitka6@gmail.com.

<http://tgdtheory.com/>.

June 20, 2019

Contents

1	Introduction	1
2	Fractal hierarchy of breakings of super-symplectic symmetry as a gauge symmetry	2

Abstract

TGD is characterized by various hierarchies. There are fractal hierarchies of quantum criticalities, Planck constants and hyper-finite factors and these hierarchies relate to hierarchies of space-time sheets, and selves. These hierarchies are closely related and this article describes these connections. In this article the recent view about connections between various hierarchies associated with quantum TGD are described.

1 Introduction

TGD is characterized various hierarchies. There are fractal hierarchies of quantum criticalities, Planck constants and hyper-finite factors and these hierarchies relate to hierarchies of space-time sheets, and selves. These hierarchies are closely related and this note describes these connections.

TGD Universe is characterized by various hierarchies. At space-time level there is a hierarchy of space-time sheets labelled by a hierarchy of p-adic length scales coming as primes near powers of two and probably generalizing to primes near powers of prime [K6, K8]. In zero energy ontology (ZEO) and at imbedding space level there is a hierarchy of causal diamonds (CDs) labelled by their size scales coming as integer multiples of CP_2 scales. In TGD based generalization of quantum theory there is a hierarchy of Planck constants characterizing hierarchy of dark matters [K2], hierarchies of inclusions of hyper-finite factors [K4], hierarchies of breakings of super-symplectic gauge symmetry [?] associated with a hierarchy of quantum criticalities

qcritdark. There is also a number theoretic hierarchy of algebraic extensions of rationals accompanied by those of p-adic number fields [K8] allowing to see evolution as a gradual increase of the complexity for extensions of rationals assignable to the parameters characterizing string world sheets and partonic 2-surfaces. In TGD inspired theory of consciousness [K3] one has self hierarchy.

At the basic level the fundamental hierarchy seems to be the heirarchy of breakings of super-symplectic symmetry as gauge symmetry. Super-symplectic algebra and its Yangian generalization have the structure of conformal algebra and is naturally associated with critical systems which are now 4-dimensional. There are also other conformal algebras involved.

By the strong form of holography implied by the strong form of General Coordinate Invariance (GCI) the core of the mathematical description of quantum TGD - analogous to genes in biology - reduces to that for 2-D systems associated with partonic 2-surfaces and string world sheets. Although space-time is 4-D, all that can be said mathematically about quantum physics can be reduced to these 2-D “genes of space-time”. 4-D space-time surfaces are however necessary for the classical description of TGD necessary to interpret quantum measurements in terms of frequencies and wavelengths classical space-time picture about particles. This reduction implies that the representations of charges of super-symplectic Yangian [K9] are in terms of fermionic strings connecting partonic 2-surfaces, which means enormous simplification of the theory. This representation also involves a generalization of AdS/CFT duality to TGD framework as manifestation of strong form of GCI basically [K1].

This note discusses the fractal hierarchy of some of breakings of super-symplectic symmetry as a gauge symmetry and some these connections.

2 Fractal hierarchy of breakings of super-symplectic symmetry as a gauge symmetry

The basic almost prediction of TGD is a fractal hierarchy of breakings of symplectic symmetry as a gauge symmetry [K7, K5].

It is good to briefly summarize the basic facts about the symplectic algebra assigned with $\delta M_{\pm}^4 \times CP_2$ first.

1. Symplectic algebra has the structure of Virasoro algebra with respect to the light-like radial coordinate r_M of the light-cone boundary taking the role of complex coordinate for ordinary conformal symmetry. The Hamiltonians generating symplectic symmetries can be chosen to be proportional to functions $f_n(r_M)$. What is the natural choice for $f_n(r_M)$ is not quite clear. Ordinary conformal invariance would suggests $f_n(r_M) = r_M^n$. A more adventurous possibility is that the algebra is generated by Hamiltonians with $f_n(r_M) = r^{-s}$, where s is a root of Riemann Zeta so that one has either $s = 1/2 + iy$ (roots at critical line) or $s = -2n, n > 0$ (roots at negative real axis).
2. The set of conformal weights would be linear space spanned by combinations of all roots with integer coefficients $s = n - iy, s = \sum n_i y_i, n > -n_0$, where $-n_0 \geq 0$ is negative conformal weight. Mass squared is proportional to the total

conformal weight and must be real demanding $y = \sum y_i = 0$ for physical states: I call this conformal confinement analogous to color confinement. One could even consider introducing the analog of binding energy as “binding conformal weight”.

Mass squared must be also non-negative (no tachyons) giving $n_0 \geq 0$. The generating conformal weights however have negative real part $-1/2$ and are thus tachyonic. Rather remarkably, p-adic mass calculations force to assume negative half-integer valued ground state conformal weight. This plus the fact that the zeros of Riemann Zeta has been indeed assigned with critical systems forces to take the Riemannian variant of conformal weight spectrum with seriousness. The algebra allows also now infinite hierarchy of conformal sub-algebras with weights coming as n -ples of the conformal weights of the entire algebra.

3. The outcome would be an infinite number of hierarchies of symplectic conformal symmetry breakings. Only the generators of the sub-algebra of the symplectic algebra with radial conformal weight proportional to n would act as gauge symmetries at given level of the hierarchy. In the hierarchy n_i divides n_{i+1} . In the symmetry breaking $n_i \rightarrow n_{i+1}$ the conformal charges, which vanished earlier, would become non-vanishing. Gauge degrees of freedom would transform to physical degrees of freedom.
4. What about the conformal Kac-Moody algebras associated with spinor modes. It seems that in this case one can assume that the conformal gauge symmetry is exact just as in string models.

The natural interpretation of the conformal hierarchies $n_i \rightarrow n_{i+1}$ would be in terms of increasing measurement resolution.

1. Conformal degrees of freedom below measurement resolution would be gauge degrees of freedom and correspond to generators with conformal weight proportional to n_i . Conformal hierarchies and associated hierarchies of Planck constants and n -fold coverings of space-time surface connecting the 3-surfaces at the ends of causal diamond would give a concrete realization of the inclusion hierarchies for hyper-finite factors of type II_1 [?].

n_i could correspond to the integer labelling Jones inclusions and associating with them the quantum group phase factor $U_n = \exp(i2\pi/n)$, $n \geq 3$ and the index of inclusion given by $|M : N| = 4\cos^2(2\pi/n)$ defining the fractal dimension assignable to the degrees of freedom above the measurement resolution. The sub-algebra with weights coming as n -multiples of the basic conformal weights would act as gauge symmetries realizing the idea that these degrees of freedom are below measurement resolution.

2. If $h_{eff} = n \times h$ defines the conformal gauge sub-algebra, the improvement of the resolution would scale up the Compton scales and would quite concretely correspond to a zoom analogous to that done for Mandelbrot fractal to get new details visible. From the point of view of cognition the improving resolution

would fit nicely with the recent view about h_{eff}/h as a kind of intelligence quotient.

This interpretation might make sense for the symplectic algebra of $\delta M_{\pm}^4 \times CP_2$ for which the light-like radial coordinate r_M of light-cone boundary takes the role of complex coordinate. The reason is that symplectic algebra acts as isometries.

3. If Kähler action has vanishing total variation under deformations defined by the broken conformal symmetries, the corresponding conformal charges are conserved. The components of WCW Kähler metric expressible in terms of second derivatives of Kähler function can be however non-vanishing and have also components, which correspond to WCW coordinates associated with different partonic 2-surfaces. This conforms with the idea that conformal algebras extend to Yangian algebras generalizing the Yangian symmetry of $\mathcal{N} = 4$ symmetric gauge theories. The deformations defined by symplectic transformations acting gauge symmetries the second variation vanishes and there is not contribution to WCW Kähler metric.
4. One can interpret the situation also in terms of consciousness theory. The larger the value of h_{eff} , the lower the criticality, the more sensitive the measurement instrument since new degrees of freedom become physical, the better the resolution. In p-adic context large n means better resolution in angle degrees of freedom by introducing the phase $exp(i2\pi/n)$ to the algebraic extension and better cognitive resolution. Also the emergence of negentropic entanglement characterized by $n \times n$ unitary matrix with density matrix proportional to unit matrix means higher level conceptualization with more abstract concepts.

The extension of the super-conformal algebra to a larger Yangian algebra is highly suggestive and gives an additional aspect to the notion of measurement resolution.

1. Yangian would be generated from the algebra of super-conformal charges assigned with the points pairs belonging to two partonic 2-surfaces as stringy Noether charges assignable to strings connecting them. For super-conformal algebra associated with pair of partonic surface only single string associated with the partonic 2-surface. This measurement resolution is the almost the poorest possible (no strings at all would be no measurement resolution at all!).
2. Situation improves if one has a collection of strings connecting set of points of partonic 2-surface to other partonic 2-surface(s). This requires generalization of the super-conformal algebra in order to get the appropriate mathematics. Tensor powers of single string super-conformal charges spaces are obviously involved and the extended super-conformal generators must be multi-local and carry multi-stringy information about physics.
3. The generalization at the first step is simple and based on the idea that co-product is the "time inverse" of product assigning to single generator sum

of tensor products of generators giving via commutator rise to the generator. The outcome would be expressible using the structure constants of the super-conformal algebra schematically a $Q_A^1 = f_A^{BC} Q_B \otimes Q_C$. Here Q_B and Q_C are super-conformal charges associated with separate strings so that 2-local generators are obtained. One can iterate this construction and get a hierarchy of n -local generators involving products of n stringy super-conformal charges. The larger the value of n , the better the resolution, the more information is coded to the fermionic state about the partonic 2-surface and 3-surface. This affects the space-time surface and hence WCW metric but not the 3-surface so that the interpretation in terms of improved measurement resolution makes sense. This super-symplectic Yangian would be behind the quantum groups and Jones inclusions in TGD Universe.

4. n gives also the number of space-time sheets in the singular covering. One possible interpretation is in terms measurement resolution for counting the number of space-time sheets. Our recent quantum physics would only see single space-time sheet representing visible matter and dark matter would become visible only for $n > 1$.

It is not an accident that quantum phases are assignable to Yangian algebras, to quantum groups, and to inclusions of HFFs. The new deep notion added to this existing complex of high level mathematical concepts are hierarchy of Planck constants, dark matter hierarchy, hierarchy of criticalities, and negentropic entanglement representing physical notions. All these aspects represent new physics.

REFERENCES

Books related to TGD

- [K1] Pitkänen M. Construction of Quantum Theory: Symmetries. In *Towards M-Matrix*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/tgdquantum.html#quthe>, 2006.
- [K2] Pitkänen M. Does TGD Predict the Spectrum of Planck Constants? In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/neuplanck.html#Planck>, 2006.
- [K3] Pitkänen M. Matter, Mind, Quantum. In *TGD Inspired Theory of Consciousness*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/tgdconsc.html#conscic>, 2006.
- [K4] Pitkänen M. Was von Neumann Right After All? In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/padphys.html#vNeumann>, 2006.
- [K5] Pitkänen M. WCW Spinor Structure. In *Quantum Physics as Infinite-Dimensional Geometry*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/tgdgeom.html#cspin>, 2006.

- [K6] Pitkänen M. *p-Adic length Scale Hypothesis*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/padphys.html>, 2013.
- [K7] Pitkänen M. Recent View about Kähler Geometry and Spin Structure of WCW . In *Quantum Physics as Infinite-Dimensional Geometry*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/tgdgeom.html#wcwnew>, 2014.
- [K8] Pitkänen M. Unified Number Theoretical Vision. In *TGD as a Generalized Number Theory*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/tgdnumber.html#numbervision>, 2014.
- [K9] Pitkänen M. The classical part of the twistor story. In *Towards M-Matrix*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/tgdquantum.html#twistorstory>, 2016.