

# Does TGD Predict a Spectrum of Planck Constants?

M. Pitkänen,

November 30, 2016

Email: [matpitka@luukku.com](mailto:matpitka@luukku.com).

[http://tgdtheory.com/public\\_html/](http://tgdtheory.com/public_html/).

Recent postal address: Karkinkatu 3 I 3, 00360, Karkkila, Finland.

## Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
1.1	Evolution Of Mathematical Ideas . . . . .	5
1.2	The Evolution Of Physical Ideas . . . . .	6
1.3	Basic Physical Picture As It Is Now . . . . .	7
<b>2</b>	<b>Experimental Input</b>	<b>9</b>
2.1	Hints For The Existence Of Large $\hbar$ Phases . . . . .	9
2.2	Quantum Coherent Dark Matter And $\hbar$ . . . . .	9
2.3	The Phase Transition Changing The Value Of Planck Constant As A Transition To Non-Perturbative Phase . . . . .	10
2.3.1	A phase transition increasing $\hbar$ as a transition guaranteeing the convergence of perturbation theory . . . . .	10
2.3.2	The criterion for the occurrence of the phase transition increasing the value of $\hbar$ . . . . .	11
<b>3</b>	<b>A Generalization Of The Notion Of Imbedding Space As A Realization Of The Hierarchy Of Planck Constants</b>	<b>11</b>
3.1	Basic Ideas . . . . .	11
3.1.1	Scaling of Planck constant and scalings of CD and $CP_2$ metrics . . . . .	12
3.1.2	Kähler function codes for a perturbative expansion in powers of $\hbar(CD)/\hbar(CP_2)$ . . . . .	12
3.1.3	Jones inclusions and hierarchy of Planck constants . . . . .	12
3.2	The Vision . . . . .	13
3.3	Hierarchy Of Planck Constants And The Generalization Of The Notion Of Imbedding Space . . . . .	15
3.3.1	The evolution of physical ideas about hierarchy of Planck constants . . . . .	15

3.3.2	The most general option for the generalized imbedding space . . . . .	16
3.3.3	About the phase transitions changing Planck constant . . . . .	17
3.3.4	How one could fix the spectrum of Planck constants? . . . . .	18
3.3.5	Preferred values of Planck constants . . . . .	18
3.3.6	How Planck constants are visible in Kähler action? . . . . .	19
<b>4</b>	<b>Updated View About The Hierarchy Of Planck Constants</b>	<b>19</b>
4.1	Basic Physical Ideas . . . . .	19
4.2	Space-Time Correlates For The Hierarchy Of Planck Constants . . . . .	20
4.3	The Relationship To The Original View About The Hierarchy Of Planck Constants	21
4.4	Basic Phenomenological Rules Of Thumb In The New Framework . . . . .	22
4.5	Charge Fractionalization And Anyons . . . . .	23
4.6	Negentropic Entanglement Between Branches Of Multi-Furcations . . . . .	24
4.7	Dark Variants Of Nuclear And Atomic Physics . . . . .	25
4.8	What About The Relationship Of Gravitational Planck Constant To Ordinary Planck Constant? . . . . .	26
4.9	Hierarchy Of Planck Constants And Non-Determinism Of Kähler Action . . . . .	27
<b>5</b>	<b>Vision About Dark Matter As Phases With Non-Standard Value Of Planck Constant</b>	<b>28</b>
5.1	Dark Rules . . . . .	28
5.1.1	The notion of relative darkness . . . . .	28
5.1.2	Is dark matter anyonic? . . . . .	28
5.1.3	Field body as carrier of dark matter . . . . .	28
5.2	Phase Transitions Changing Planck Constant . . . . .	29
5.2.1	Transition to large $\hbar$ phase and failure of perturbation theory . . . . .	29
5.2.2	The mechanism of phase transition and selection rules . . . . .	30
5.3	Coupling Constant Evolution And Hierarchy Of Planck Constants . . . . .	30
5.3.1	Evolution with respect to phase resolution . . . . .	30
5.3.2	Is there a correlation between the values of p-adic prime and Planck constant? . . . . .	31
<b>6</b>	<b>Some Applications</b>	<b>31</b>
6.1	A Simple Model Of Fractional Quantum Hall Effect . . . . .	31
6.2	Gravitational Bohr Orbitology . . . . .	33
6.2.1	Dark matter as large $\hbar$ phase . . . . .	33
6.2.2	Prediction for the parameter $v_0$ . . . . .	33
6.2.3	Further predictions . . . . .	33
6.2.4	Comparison with Bohr quantization of planetary orbits . . . . .	34
6.2.5	Quantum Hall effect and dark anyonic systems in astrophysical scales . . . . .	36
6.2.6	Anyonic view about blackholes . . . . .	36
6.3	Accelerating Periods Of Cosmic Expansion As Phase Transitions Increasing The Value Of Planck Constant . . . . .	37
6.3.1	The four pieces of evidence for accelerated expansion . . . . .	37
6.3.2	Accelerated expansion in classical TGD . . . . .	38
6.3.3	Accelerated expansion and hierarchy of Planck constants . . . . .	38
6.3.4	Accelerated expansion and flatness of 3-cosmology . . . . .	38
6.3.5	The size of large voids is the characteristic scale . . . . .	38
6.3.6	Do cosmic strings with negative gravitational mass cause the phase transition inducing accelerated expansion . . . . .	39
6.4	Phase Transition Changing Planck Constant And Expanding Earth Theory . . . . .	39
6.4.1	The claims of Adams . . . . .	40
6.4.2	The critic of Adams of the subduction mechanism . . . . .	40
6.4.3	Expanding Earth theories are not new . . . . .	41
6.4.4	Summary of TGD based theory of Expanding Earth . . . . .	41
6.4.5	Did intra-terrestrial life burst to the surface of Earth during Cambrian ex- pansion? . . . . .	43
6.5	Allais Effect As Evidence For Large Values Of Gravitational Planck Constant? . . . . .	44

---

6.5.1	Experimental findings . . . . .	44
6.5.2	TGD based models for Allais effect . . . . .	44
6.6	Applications To Elementary Particle Physics, Nuclear Physics, And Condensed Matter Physics . . . . .	45
6.6.1	Leptohadron hypothesis . . . . .	45
6.6.2	Cold fusion, plasma electrolysis, and burning salt water . . . . .	45
6.7	Applications To Biology And Neuroscience . . . . .	46
6.7.1	Do molecular symmetries in living matter relate to non-standard values of Planck constant? . . . . .	46
6.7.2	High $T_c$ super-conductivity in living matter . . . . .	47
6.7.3	Magnetic body as a sensory perceiver and intentional agent . . . . .	48
6.7.4	DNA as topological quantum computer . . . . .	49
6.7.5	Quantum model of nerve pulse and EEG . . . . .	51
<b>7</b>	<b>Appendix</b>	<b>52</b>
7.1	About Inclusions Of Hyper-Finite Factors Of Type $II_1$ . . . . .	52
7.2	Generalization From $Su(2)$ To Arbitrary Compact Group . . . . .	53

### Abstract

The quantization of Planck constant has been the basic theme of TGD since 2005. The basic idea was stimulated by the suggestion of Nottale that planetary orbits could be seen as Bohr orbits with enormous value of Planck constant given by  $\hbar_{gr} = GM_1M_2/v_0$ , where the velocity parameter  $v_0$  has the approximate value  $v_0 \simeq 2^{-11}$  for the inner planets. This inspired the ideas that quantization is due to a condensation of ordinary matter around dark matter concentrated near Bohr orbits and that dark matter is in macroscopic quantum phase in astrophysical scales. The second crucial empirical input were the anomalies associated with living matter. The recent version of the chapter represents the evolution of ideas about quantization of Planck constants from a perspective given by seven years's work with the idea. A very concise summary about the situation is as follows.

#### 1. Basic physical ideas

The basic phenomenological rules are simple.

1. The phases with non-standard values of effective Planck constant are identified as dark matter. The motivation comes from the natural assumption that only the particles with the same value of effective Planck can appear in the same vertex. One can illustrate the situation in terms of the book metaphor. Effective imbedding spaces with different values of Planck constant form a book like structure and matter can be transferred between different pages only through the back of the book where the pages are glued together. One important implication is that light exotic charged particles lighter than weak bosons are possible if they have non-standard value of Planck constant. The standard argument excluding them is based on decay widths of weak bosons and has led to a neglect of large number of particle physics anomalies.
2. Large effective or real value of Planck constant scales up Compton length - or at least de Broglie wave length - and its geometric correlate at space-time level identified as size scale of the space-time sheet assignable to the particle. This could correspond to the Kähler magnetic flux tube for the particle forming consisting of two flux tubes at parallel space-time sheets and short flux tubes at ends with length of order  $CP_2$  size.

This rule has far reaching implications in quantum biology and neuroscience since macroscopic quantum phases become possible as the basic criterion stating that macroscopic quantum phase becomes possible if the density of particles is so high that particles as Compton length sized objects overlap. Dark matter therefore forms macroscopic quantum phases. One implication is the explanation of mysterious looking quantal effects of ELF radiation in EEG frequency range on vertebrate brain:  $E = hf$  implies that the energies for the ordinary value of Planck constant are much below the thermal threshold but large value of Planck constant changes the situation. Also the phase transitions modifying the value of Planck constant and changing the lengths of flux tubes (by quantum classical correspondence) are crucial as also reconnections of the flux tubes.

The hierarchy of Planck constants suggests also a new interpretation for FQHE (fractional quantum Hall effect) in terms of anyonic phases with non-standard value of effective Planck constant realized in terms of the effective multi-sheeted covering of imbedding space: multi-sheeted space-time is to be distinguished from many-sheeted space-time.

In astrophysics and cosmology the implications are even more dramatic. The interpretation of  $\hbar_{gr}$  introduced by Nottale in TGD framework is as an effective Planck constant associated with space-time sheets mediating gravitational interaction between masses  $M$  and  $m$ . The huge value of  $\hbar_{gr}$  means that the integer  $\hbar_{gr}/\hbar_0$  interpreted as the number of sheets of covering is gigantic and that Universe possesses gravitational quantum coherence in astronomical scales. The gravitational Compton length  $GM/v_0 = r_S/2v_0$  does not depend on  $m$  so that all particles around say Sun say same gravitational Compton length.

By the independence of gravitational acceleration and gravitational Compton length on particle mass, it is enough to assume that only microscopic particles couple to the dark gravitons propagating along flux tubes mediating gravitational interaction. Therefore  $\hbar_{gr} = \hbar_{eff}$  could be true in microscopic scales and would predict that cyclotron energies have no dependence on the mass of the charged particle meaning that the spectrum ordinary photons resulting in the transformation of dark photons to ordinary photons is universal. An attractive identification of these photons would be as bio-photons with energies in visible and UV range and thus inducing molecular transitions making control of biochemistry by dark photons. This changes the view about gravitons and suggests

that gravitational radiation is emitted as dark gravitons which decay to pulses of ordinary gravitons replacing continuous flow of gravitational radiation. The energy of the graviton is gigantic unless the emission is assumed to take place from a microscopic system with large but not gigantic  $h_{gr}$ .

3. Why Nature would like to have large - maybe even gigantic - value of effective value of Planck constant? A possible answer relies on the observation that in perturbation theory the expansion takes in powers of gauge coupling strengths  $\alpha = g^2/4\pi\hbar$ . If the effective value of  $\hbar$  replaces its real value as one might expect to happen for multi-sheeted particles behaving like single particle,  $\alpha$  is scaled down and perturbative expansion converges for the new particles. One could say that Mother Nature loves theoreticians and comes in rescue in their attempts to calculate. In quantum gravitation the problem is especially acute since the dimensionless parameter  $GMm/\hbar$  has gigantic value. Replacing  $\hbar$  with  $\hbar_{gr} = GMm/v_0$  the coupling strength becomes  $v_0 < 1$ .

### 2. Space-time correlates for the hierarchy of Planck constants

The hierarchy of Planck constants was introduced to TGD originally as an additional postulate and formulated as the existence of a hierarchy of imbedding spaces defined as Cartesian products of singular coverings of  $M^4$  and  $CP_2$  with numbers of sheets given by integers  $n_a$  and  $n_b$  and  $\hbar = n\hbar_0$ .  $n = n_a n_b$ .

With the advent of zero energy ontology (ZEO), it became clear that the notion of singular covering space of the imbedding space could be only a convenient auxiliary notion. Singular means that the sheets fuse together at the boundary of multi-sheeted region. In ZEO 3-surfaces are unions of space-like 3-surfaces at opposite boundaries of CD. The non-determinism of Kähler action due to the huge vacuum degeneracy would naturally explain the existence of several space-time sheets connecting the two 3-surfaces at the opposite boundaries of CD. Quantum criticality suggests strongly conformal invariance and the identification of  $n$  as the number of conformal equivalence classes of these space-time sheets. Also a connection with the notion of negentropic entanglement emerges.

## 1 Introduction

The quantization of Planck constant has been the basic theme of TGD since 2005 and the perspective in the earlier version of this chapter reflected the situation for about year and one half after the basic idea stimulated by the finding of Nottale [E6] that planetary orbits could be seen as Bohr orbits with enormous value of Planck constant given by  $\hbar_{gr} = GM_1 M_2 / v_0$ ,  $v_0 \simeq 2^{-11}$  for the inner planets. The general form of  $\hbar_{gr}$  is dictated by Equivalence Principle. This inspired the ideas that quantization is due to a condensation of ordinary matter around dark matter concentrated near Bohr orbits and that dark matter is in macroscopic quantum phase in astrophysical scales.

The second crucial empirical input were the anomalies associated with living matter. Mention only the effects of ELF radiation at EEG frequencies on vertebrate brain and anomalous behavior of the ionic currents through cell membrane. If the value of Planck constant is large, the energy of EEG photons is above thermal energy and one can understand the effects on both physiology and behavior. If ionic currents through cell membrane have large Planck constant the scale of quantum coherence is large and one can understand the observed low dissipation in terms of quantum coherence. This approach led to the formula  $h_{eff} = n \times h$ . Rather recently (2014) it became clear that for microscopic systems the identification  $h_{eff} = h_{gr}$  makes sense and predicts universal energy spectrum for cyclotron energies of dark photons identifiable as energy spectrum of bio-photons in TGD inspired quantum biology.

### 1.1 Evolution Of Mathematical Ideas

The original formulation for the hierarchy of Planck constants was in terms of  $h_{eff}/h = n$ -fold singular coverings of the imbedding space  $H = M^4 \times CP_2$ . Later it turned out that there is no need to postulate these covering spaces although they are a nice auxiliary tool allowing to understand why the phase of matter with different values of  $n$  behave like dark matter relative to each other: they are simply at different pages of the book-like structure formed by the covering spaces.

Few years ago it became clear that the hierarchy of Planck constants could be only effective but have the same practical implications. The basic observation was that the effective hierarchy need

not be postulated separately but follows as a prediction from the vacuum degeneracy of Kähler action. In this formulation Planck constant at fundamental level has its standard value and its effective values come as its integer multiples so that one should write  $h_{eff} = n \times h$  rather than  $\hbar = n\hbar_0$  as I have done. For most practical purposes the states in question would behave as if Planck constant were an integer multiple of the ordinary one. This reduces the understanding of the effective hierarchy of Planck constants to quantum variant of multi-furcations for the dynamics of preferred extremals of Kähler action. The number of branches of multi-furcation defines the integer  $n$  in  $\hbar_{eff} = n\hbar$ .

One of the latest steps in the progress was the realization that the hierarchy of Planck constants can be understood in terms of quantum criticality of TGD Universe postulated from the beginning as a manner to obtain a unique theory. In accordance with what is known about 2-D critical systems, quantum criticality should correspond to a generalization of conformal invariance. TGD indeed predicts several analogs of super-conformal algebras: so called super-symplectic algebra acting in  $\delta M_{\pm}^4 \times CP_2$  should act as isometries of WCW and its generators are labeled by conformal weights. Light-cone boundary  $\delta M_{\pm}^4$  has an extension of conformal symmetries as conformal symmetries and an algebra isomorphic to the ordinary conformal algebra acts as its isometries. The light-like orbits of partonic 2-surfaces allow similar algebra of conformal symmetries and string world sheets and partonic 2-surfaces allow conformal symmetries.

The proposal is that super-symplectic algebra (at least it) defines a hierarchy of broken super-conformal gauge symmetries in the sense that the sub-algebra for which the conformal weights are  $n$ -ples of those for the entire algebra acts as gauge conformal symmetries.  $n = h_{eff}/h$  giving a connection to the hierarchy of Planck constants would hold true. These sub-algebras are isomorphic to the full algebra and thus form a fractal hierarchy. One has infinite number of hierarchies of broken conformal symmetries defined by the sequences  $n(i+1) = m_i \times n(i)$ . In the phase transition increasing  $n$  conformal gauge symmetry is reduced and some gauge degrees of freedom transform to physical ones and criticality is reduced so that the transition takes place spontaneously. TGD Universe is like a ball at the top of hill at the top of hill at...

This view has far reaching implication for the understanding of living matter and leads to deep connections between different key ideas of TGD. The hierarchy has also a purely number theoretical interpretation in terms of hierarchy of algebraic extensions of rationals appearing naturally in the adelic formulation of quantum TGD.  $n = h_{eff}/h$  would naturally correspond to an integer, which is product of so called ramified primes (rational primes for which the decomposition to primes of extension contains higher powers of these primes).

In this framework it becomes obvious that - instead of coverings of imbedding space postulated in the original formulation - one has space-time surfaces representable as singular  $n$ -fold coverings. The non-determinism of Kähler action - key element of criticality - would be the basic reason for the appearance of singular coverings: two 3-surfaces at the opposite boundaries of CD are connected by  $n$ -sheeted space-time surfaces for which the sheets co-incide at the boundaries. Criticality must be accompanied by 4-D variant of conformal gauge invariance already described so that these space-time surfaces are replaced by conformal gauge equivalence classes.

These coverings are highly analogous to the covering space associated with the analytic function  $w(z) = z^{1/n}$ . If one uses  $w$  as a variable, the ordinary conformal symmetries generated by functions of  $z$  indeed correspond to the algebra generated by  $w^n$  and the sheets of covering correspond to conformal gauge equivalence classes not transformed to each other by conformal transformations.

## 1.2 The Evolution Of Physical Ideas

The evolution of physical ideas related to the hierarchy of Planck constants and dark matter as a hierarchy of phases of matter with non-standard value of Planck constants was much faster than the evolution of mathematical ideas and quite a number of applications have been developed during last five years.

1. The basic idea was that ordinary matter condenses around dark matter which is a phase of matter characterized by non-standard value of Planck constant.
2. The realization that non-standard values of Planck constant give rise to charge and spin fractionization and anyonization led to the precise identification of the prerequisites of anyonic

phase [K17] . If the partonic 2-surface, which can have even astrophysical size, surrounds the tip of CD, the matter at the surface is anyonic and particles are confined at this surface. Dark matter could be confined inside this kind of light-like 3-surfaces around which ordinary matter condenses. If the radii of the basic pieces of these nearly spherical anyonic surfaces - glued to a connected structure by flux tubes mediating gravitational interaction - are given by Bohr rules, the findings of Nottale [E6] can be understood. Dark matter would resemble to a high degree matter in black holes replaced in TGD framework by light-like partonic 2-surfaces with minimum size of order Schwarzschild radius  $r_S$  of order scaled up Planck length:  $r_S \sim \sqrt{\hbar G}$ . Black hole entropy being inversely proportional to  $\hbar$  is predicted to be of order unity so that dramatic modification of the picture about black holes is implied.

3. Darkness is a relative concept and due to the fact that particles at different pages of book cannot appear in the same vertex of the generalized Feynman diagram. The phase transitions in which partonic 2-surface  $X^2$  during its travel along  $X_i^3$  leaks to different page of book are however possible and change Planck constant so that particle exchanges of this kind allow particles at different pages to interact. The interactions are strongly constrained by charge fractionization and are essentially phase transitions involving many particles. Classical interactions are also possible. This allows to conclude that we are actually observing dark matter via classical fields all the time and perhaps have even photographed it [K20] , [I4] .
4. Perhaps the most fascinating applications are in biology. The anomalous behavior ionic currents through cell membrane (low dissipation, quantal character, no change when the membrane is replaced with artificial one) has a natural explanation in terms of dark supra currents. This leads to a vision about how dark matter and phase transitions changing the value of Planck constant could relate to the basic functions of cell, functioning of DNA and amino-acids, and to the mysteries of bio-catalysis. This leads also a model for EEG interpreted as a communication and control tool of magnetic body containing dark matter and using biological body as motor instrument and sensory receptor. One especially shocking outcome is the emergence of genetic code of vertebrates from the model of dark nuclei as nuclear strings [L1, K20] , [L1] .

### 1.3 Basic Physical Picture As It Is Now

The basic phenomenological rules are simple and remained roughly the same during years.

1. The phases with non-standard values of effective Planck constant are identified as dark matter. The motivation comes from the natural assumption that only the particles with the same value of effective Planck can appear in the same vertex. One can illustrate the situation in terms of the book metaphor. Imbedding spaces with different values of Planck constant form a book like structure and matter can be transferred between different pages only through the back of the book where the pages are glued together. One important implication is that light exotic charged particles lighter than weak bosons are possible if they have non-standard value of Planck constant. The standard argument excluding them is based on decay widths of weak bosons and has led to a neglect of large number of particle physics anomalies [K21].
2. Large effective or real value of Planck constant scales up Compton length - or at least de Broglie wave length - and its geometric correlate at space-time level identified as size scale of the space-time sheet assignable to the particle. This could correspond to the Kähler magnetic flux tube for the particle forming consisting of two flux tubes at parallel space-time sheets and short flux tubes at ends with length of order  $CP_2$  size.

This rule has far reaching implications in quantum biology and neuroscience since macroscopic quantum phases become possible as the basic criterion stating that macroscopic quantum phase becomes possible if the density of particles is so high that particles as Compton length sized objects overlap. Dark matter therefore forms macroscopic quantum phases. One implication is the explanation of mysterious looking quantal effects of ELF radiation in EEG frequency range on vertebrate brain:  $E = hf$  implies that the energies for the ordinary value of Planck constant are much below the thermal threshold but large value of Planck constant changes the situation. Also the phase transitions modifying the value of Planck constant and

changing the lengths of flux tubes (by quantum classical correspondence) are crucial as also reconnections of the flux tubes.

The hierarchy of Planck constants suggests also a new interpretation for FQHE (fractional quantum Hall effect) [K17] in terms of anyonic phases with non-standard value of effective Planck constant realized in terms of the effective multi-sheeted covering of imbedding space: multi-sheeted space-time is to be distinguished from many-sheeted space-time.

In astrophysics and cosmology the implications are even more dramatic. It was Nottale [E6] who first introduced the notion of gravitational Planck constant as  $\hbar_{gr} = GMm/v_0$ ,  $v_0 < 1$  has interpretation as velocity light parameter in units  $c = 1$ . This would be true for  $GMm/v_0 \geq 1$ . The interpretation of  $\hbar_{gr}$  in TGD framework is as an effective Planck constant associated with space-time sheets mediating gravitational interaction between masses  $M$  and  $m$ . The huge value of  $\hbar_{gr}$  means that the integer  $\hbar_{gr}/\hbar_0$  interpreted as the number of sheets of covering is gigantic and that Universe possesses gravitational quantum coherence in super-astronomical scales for masses which are large. This changes the view about gravitons and suggests that gravitational radiation is emitted as dark gravitons which decay to pulses of ordinary gravitons replacing continuous flow of gravitational radiation.

3. Why Nature would like to have large effective value of Planck constant? A possible answer relies on the observation that in perturbation theory the expansion takes in powers of gauge couplings strengths  $\alpha = g^2/4\pi\hbar$ . If the effective value of  $\hbar$  replaces its real value as one might expect to happen for multi-sheeted particles behaving like single particle,  $\alpha$  is scaled down and perturbative expansion converges for the new particles. One could say that Mother Nature loves theoreticians and comes in rescue in their attempts to calculate. In quantum gravitation the problem is especially acute since the dimensionless parameter  $GMm/\hbar$  has gigantic value. Replacing  $\hbar$  with  $\hbar_{gr} = GMm/v_0$  the coupling strength becomes  $v_0 < 1$ .
4. The interpretation of the hierarchy of Planck constants as labels for quantum critical systems is especially powerful in TGD inspired quantum biology and consciousness theory. The increase of Planck constant by integer factor occurs spontaneously and means an increase of complexity and sensory and cognitive resolution - in other words evolution. Living matter is however fighting to stay at the existing level of criticality. The reason is that the changes involves state function reduction at the opposite boundary of CD and means death of self followed by re-incarnation.

Negentropy Maximization Principle [K12] saves the system from this fate if it is able to generate negentropic entanglement by some other means. Metabolic energy suggested already earlier to be a carrier of negentropic entanglement makes this possible. Also other metabolites can carry negentropy. Therefore living systems are eating each other to satisfy the demands of NMP! Why this non-sensical looking Karma's cycle? The sub-systems of self defining sub-selves (mental images) are dying and re-incarnating and generating negentropy: self is a gardener and sub-selves are the fruit trees and the longer self lives, the more fruits are produced. Hence this process, which Buddhist would call attachment to ego is the manner to generate what I have called "Akashic records". Everything has its purpose.

In this chapter I try to summarize the evolution of the ideas related to Planck constant. I have worked hardly to to achieve internal consistency but the old theory layers are there and might cause confusion.

The appendix of the book gives a summary about basic concepts of TGD with illustrations. There are concept maps about topics related to the contents of the chapter prepared using CMAP realized as html files. Links to all CMAP files can be found at <http://tgdtheory.fi/cmaphtml.html> [L2]. Pdf representation of same files serving as a kind of glossary can be found at <http://tgdtheory.fi/tgdglossary.pdf> [L3]. The topics relevant to this chapter are given by the following list.

- Hierarchy of Planck constants [L5]
- Geometrization of fields [L4]
- Magnetic body [L6]



## 2 Experimental Input

In this section basic experimental inputs suggesting a hierarchy of Planck constants and the identification of dark matter as phases with non-standard value of Planck constant are discussed.

### 2.1 Hints For The Existence Of Large $\hbar$ Phases

Quantum classical correspondence suggests the identification of space-time sheets identifiable as quantum coherence regions. Since they can have arbitrarily large sizes, phases with arbitrarily large quantum coherence lengths and arbitrarily long de-coherence times seem to be possible in TGD Universe. In standard physics context this seems highly implausible. If Planck constant can have arbitrarily large values, the situation changes since Compton lengths and other quantum scales are proportional to  $\hbar$ . Dark matter is excellent candidate for large  $\hbar$  phases.

The expression for  $\hbar_{gr}$  in the model explaining the Bohr orbits for planets is of form  $\hbar_{gr} = GM_1M_2/v_0$  [K18]. This suggests that the interaction is associated with some kind of interface between the systems, perhaps join along boundaries bonds/flux tubes connecting the space-time sheets associated with systems possessing gravitational masses  $M_1$  and  $M_2$ . Also a large space-time sheet carrying the mutual classical gravitational field could be in question. This argument generalizes to the case  $\hbar/\hbar_0 = Q_1Q_2\alpha/v_0$  in case of generic phase transition to a strongly interacting phase with  $\alpha$  describing gauge coupling strength.

There exist indeed some experimental indications for the existence of phases with a large  $\hbar$ .

1. With inspiration coming from the finding of Nottale [E6] I have proposed an explanation of dark matter as a macroscopic quantum phase with a large value of  $\hbar$  [K18]. Any interaction, if sufficiently strong, can lead to this kind of phase. The increase of  $\hbar$  would make the fine structure constant  $\alpha$  in question small and guarantee the convergence of perturbation series.
2. Living matter could represent a basic example of large  $\hbar$  phase [K7, K3]. Even ordinary condensed matter could be “partially dark” in many-sheeted space-time [K9]. In fact, the realization of hierarchy of Planck constants leads to a considerably weaker notion of darkness stating that only the interaction vertices involving particles with different values of Planck constant are impossible and that the notion of darkness is relative notion. For instance, classical interactions and photon exchanges involving a phase transition changing the value of  $\hbar$  of photon are possible in this framework.
3. There is claim about a detection in RHIC (Relativistic Heavy Ion Collider in Brookhaven) of states behaving in some respects like mini black holes [C5]. These states could have explanation as color flux tubes at Hagedorn temperature forming a highly tangled state and identifiable as stringy black holes of strong gravitation. The strings would carry a quantum coherent color glass condensate, and would be characterized by a large value of  $\hbar$  naturally resulting in confinement phase with a large value of  $\alpha_s$  [K19]. The progress in hadronic mass calculations led to a concrete model of color glass condensate of single hadron as many-particle state of super-symplectic gluons [K15, K13] - something completely new from the point of QCD - responsible for non-perturbative aspects of hadron physics. In RHIC events these color glass condensate would fuse to single large condensate. This condensate would be present also in ordinary black-holes and the blackness of black-hole would be darkness.
4. I have also discussed a model for cold fusion based on the assumption that nucleons can be in large  $\hbar$  phase. In this case the relevant strong interaction strength is  $Q_1Q_2\alpha_{em}$  for two nucleon clusters inside nucleus which can increase  $\hbar$  so large that the Compton length of protons becomes of order atomic size and nuclear protons form a macroscopic quantum phase [K9, K7].

### 2.2 Quantum Coherent Dark Matter And $\hbar$

The argument based on gigantic value of  $\hbar_{gr}$  explaining darkness of dark matter is attractive but one should be very cautious.

Consider first ordinary QED  $e = \sqrt{\alpha 4\pi\hbar}$  appears in vertices so that perturbation expansion in powers of  $\sqrt{\hbar}$  basically. This would suggest that large  $\hbar$  leads to large effects. All predictions are

however in powers of alpha and large  $\hbar$  means small higher order corrections. What happens can be understood on basis of dimensional analysis. For instance, cross sections are proportional to  $(\hbar/m)^2$ , where  $m$  is the relevant mass and the remaining factor depends on  $\alpha = e^2/(4\pi\hbar)$  only. In the more general case tree amplitudes with  $n$  vertices are proportional to  $e^n$  and thus to  $\hbar^{n/2}$  and loop corrections give only powers of  $\alpha$  which get smaller when  $\hbar$  increases. This must relate to the powers of  $1/\hbar$  from the integration measure associated with the momentum loop integrals affected by the change of  $\alpha$ .

Consider now the effects of the scaling of  $\hbar$ . The scaling of Compton lengths and other quantum kinematical parameters is the most obvious effect. An obvious effect is due to the change of  $\hbar$  in the commutation relations and in the change of unit of various quantum numbers. In particular, the right hand side of oscillator operator commutation and anti-commutation relations is scaled. A further effect is due to the scaling of the eigenvalues of the Kähler-Dirac operator  $\hbar\Gamma^\alpha D_\alpha$ .

The exponent  $exp(K)$  of Kähler function  $K$  defining perturbation series in WCW degrees of freedom is proportional to  $1/g_K^2$  and does not depend on  $\hbar$  at all if there is only single Planck constant. The propagator is proportional to  $g_K^2$ . This can be achieved also in QED by absorbing  $e$  from vertices to  $e^2$  in photon propagator. Hence it would seem that the dependence on  $\alpha_K$  (and  $\hbar$ ) must come from vertices which indeed involve Jones inclusions of the  $II_1$  factors of the incoming and outgoing lines.

This however suggests that the dependence of the scattering amplitudes on  $\hbar$  is purely kinematical so that all higher radiative corrections would be absent. This seems to leave only one option: the scale factors of covariant CD and  $CP_2$  metrics can vary and might have discrete spectrum of values.

1. The invariance of Kähler action with respect to overall scaling of metric however allows to keep  $CP_2$  metric fixed and consider only a spectrum for the scale factors of  $M^4$  metric.
2. The first guess motivated by Schrödinger equation is that the scaling factor of covariant CD metric corresponds the ratio  $r^2 = (\hbar/\hbar_0)^2$ . This would mean that the value of Kähler action depends on  $r^2$ . The scaling of  $M^4$  coordinate by  $r$  the metric reduces to the standard form but if causal diamonds with quantized temporal distance between their tips are the basic building blocks of WCW geometry as zero energy ontology requires, this scaling of  $\hbar$  scales the size of CD by  $r$  so that genuine effect results since  $M^4$  scalings are not symmetries of Kähler action.
3. In this picture  $r$  would code for radiative corrections to Kähler function and thus space-time physics. Even in the case that the radiative corrections to WCW functional integral vanish, as suggested by quantum criticality, they would be actually taken into account.

This kind of dynamics is not consistent with the original view about imbedding space and forces to generalize the notion of imbedding spaces since it is clear that particles with different Planck constants cannot appear in the same vertex of Feynman diagram. Somehow different values of Planck constant must be analogous to different pages of book having almost copies of imbedding space as pages. A possible resolution of the problem comes from the realization that the fundamental structure might be the inclusion hierarchy of number theoretical Clifford algebras from which entire TGD could emerge including generalization of the imbedding space concept.

## **2.3 The Phase Transition Changing The Value Of Planck Constant As A Transition To Non-Perturbative Phase**

### **2.3.1 A phase transition increasing $\hbar$ as a transition guaranteeing the convergence of perturbation theory**

The general vision is that a phase transition increasing  $\hbar$  occurs when perturbation theory ceases to converge. Very roughly, this would occur when the parameter  $x = Q_1 Q_2 \alpha$  becomes larger than one. The net quantum numbers for “spontaneously magnetized” regions provide new natural units for quantum numbers. The assumption that standard quantization rules prevail poses very strong restrictions on allowed physical states and selects a subspace of the original configuration space. One can of course, consider the possibility of giving up these rules at least partially in which case

a spectrum of fractionally charged anyon like states would result with confinement guaranteed by the fractionization of charges.

The necessity of large  $\hbar$  phases has been actually highly suggestive since the first days of quantum mechanics. The classical looking behavior of macroscopic quantum systems remains still a poorly understood problem and large  $\hbar$  phases provide a natural solution of the problem.

In TGD framework quantum coherence regions correspond to space-time sheets. Since their sizes are arbitrarily large the conclusion is that macroscopic and macro-temporal quantum coherence are possible in all scales. Standard quantum theory definitely fails to predict this and the conclusion is that large  $\hbar$  phases for which quantum length and time scales are proportional to  $\hbar$  and long are needed.

Somewhat paradoxically, large  $\hbar$  phases explain the effective classical behavior in long length and time scales. Quantum perturbation theory is an expansion in terms of gauge coupling strengths inversely proportional to  $\hbar$  and thus at the limit of large  $\hbar$  classical approximation becomes exact. Also the Coulomb contribution to the binding energies of atoms vanishes at this limit. The fact that we experience world as a classical only tells that large  $\hbar$  phase is essential for our sensory perception. Of course, this is not the whole story and the full explanation requires a detailed anatomy of quantum jump.

### **2.3.2 The criterion for the occurrence of the phase transition increasing the value of $\hbar$**

In the case of planetary orbits the large value of  $\hbar_{gr} = 2GM/v_0$  makes possible to apply Bohr quantization to planetary orbits. This leads to a more general idea that the phase transition increasing  $\hbar$  occurs when the system consisting of interacting units with charges  $Q_i$  becomes non-perturbative in the sense that the perturbation series in the coupling strength  $\alpha Q_i Q_j$ , where  $\alpha$  is the appropriate coupling strength and  $Q_i Q_j$  represents the maximum value for products of gauge charges, ceases to converge. Thus Mother Nature would resolve the problems of theoretician. A primitive formulation for this criterion is the condition  $\alpha Q_i Q_j \geq 1$ .

The first working hypothesis was the existence of dark matter hierarchies with  $\hbar = \lambda^k \hbar_0$ ,  $k = 0, 1, \dots$ ,  $\lambda = n/v_0$  or  $\lambda = 1/nv_0$ ,  $v_0 \simeq 2^{-11}$ . This rule turned out to be quite too specific. The mathematically plausible formulation predicts that in principle any rational value for  $r = \hbar(M^4)/\hbar(CP_2)$  is possible but there are certain number theoretically preferred values of  $r$  such as those coming powers of 2.

## **3 A Generalization Of The Notion Of Imbedding Space As A Realization Of The Hierarchy Of Planck Constants**

In the following the basic ideas concerning the realization of the hierarchy of Planck constants are summarized and after that a summary about generalization of the imbedding space is given. In [K17] the important delicacies associated with the Kähler structure of generalized imbedding space are discussed. The background for the recent vision is quite different from that for half decade ago. Zero energy ontology and the notion of causal diamond, number theoretic compactification leading to the precise identification of number theoretic braids, the realization of number theoretic universality, and the understanding of the quantum dynamics at the level of Kähler-Dirac action fix to a high degree the vision about generalized imbedding space.

### **3.1 Basic Ideas**

The first key idea in the geometric realization of the hierarchy of Planck constants emerges from the study of Schrödinger equation and states that Planck constant appears a scaling factor of  $M^4$  metric. Second key idea is the connection with Jones inclusions inspiring an explicit formula for Planck constants. For a long time this idea remained heuristic must-be-true feeling but the recent view about quantum TGD provide a justification for it.

### 3.1.1 Scaling of Planck constant and scalings of CD and $CP_2$ metrics

The key property of Schrödinger equation is that kinetic energy term depends on  $\hbar$  whereas the potential energy term has no dependence on it. This makes the scaling of  $\hbar$  a non-trivial transformation. If the contravariant metric scales as  $r = \hbar/\hbar_0$  the effect of scaling of Planck constant is realized at the level of imbedding space geometry provided it is such that it is possible to compare the regions of generalized imbedding space having different value of Planck constant.

In the case of Dirac equation same conclusion applies and corresponds to the minimal substitution  $p - eA \rightarrow i\hbar\nabla - eA$ . Consider next the situation in TGD framework.

1. The minimal substitution  $p - eA \rightarrow i\hbar\nabla - eA$  does not make sense in the case of  $CP_2$  Dirac operator since, by the non-triviality of spinor connection, one cannot choose the value of  $\hbar$  freely. In fact, spinor connection of  $CP_2$  is defined in such a manner that spinor connection corresponds to the quantity  $\hbar eQA$ , where denotes  $A$  gauge potential, and there is no natural manner to separate  $\hbar e$  from it.
2. The contravariant CD metric scales like  $\hbar^2$ . In the case of Dirac operator in  $M^4 \times CP_2$  one can assign separate Planck constants to Poincare and color algebras and the scalings of CD and  $CP_2$  metrics induce scalings of corresponding values of  $\hbar^2$ . As far as Kähler action is considered,  $CP_2$  metric could be always thought of being scaled to its standard form.
3. Dirac equation gives the eigenvalues of wave vector squared  $k^2 = k^i k_i$  rather than four-momentum squared  $p^2 = p^i p_i$  in CD degrees of freedom and its analog in  $CP_2$  degrees of freedom. The values of  $k^2$  are proportional to  $1/r^2$  so that  $p^2$  does not depend on it for  $p^i = \hbar k^i$ : analogous conclusion applies in  $CP_2$  degrees of freedom. This gives rise to the invariance of mass squared and the desired scaling of wave vector when  $\hbar$  changes.

This consideration generalizes to the case of the induced gamma matrices and induced metric in  $X^4$ , Kähler-Dirac operator, and Kähler action which carry dynamical information about the ratio  $r = \hbar_{eff}/\hbar_0$ .

### 3.1.2 Kähler function codes for a perturbative expansion in powers of $\hbar(CD)/\hbar(CP_2)$

Suppose that one accepts that the spectrum of CD *resp.*  $CP_2$  Planck constants is accompanied by a hierarchy of overall scalings of covariant CD (causal diamond) metric by  $(\hbar(M^4)/\hbar_0)^2$  and  $CP_2$  metric by  $(\hbar(CP_2)/\hbar_0)^2$  followed by overall scaling by  $r^2 = (\hbar_0/\hbar(CP_2))^2$  so that  $CP_2$  metric suffers no scaling and difficulties with isometric gluing procedure of sectors are avoided.

The first implication of this picture is that the Kähler-Dirac operator determined by the induced metric and spinor structure depends on  $r$  in a highly nonlinear manner but there is no dependence on the overall scaling of the  $H$  metric. This in turn implies that the fermionic oscillator algebra used to define WCW spinor structure and metric depends on the value of  $r$ . Same is true also for Kähler action and configuration space Kähler function. Hence Kähler function is analogous to an effective action expressible as infinite series in powers of  $r$ .

This interpretation allows to overcome the paradox caused by the hypothesis that loop corrections to the functional integral over WCW defined by the exponent of Kähler function serving as vacuum functional vanish so that tree approximation is exact. This would imply that all higher order corrections usually interpreted in terms of perturbative series in powers of  $1/\hbar$  vanish. The paradox would result from the fact that scattering amplitudes would not receive higher order corrections and classical approximation would be exact.

The dependence of both states created by Super Kac-Moody algebra and the Kähler function and corresponding propagator identifiable as contravariant WCW metric would mean that the expressions for scattering amplitudes indeed allow an expression in powers of  $r$ . What is so remarkable is that the TGD approach would be non-perturbative from the beginning and “semi-classical” approximation, which might be actually exact, automatically would give a full expansion in powers of  $r$ . This is in a sharp contrast to the usual quantization approach.

### 3.1.3 Jones inclusions and hierarchy of Planck constants

From the beginning it was clear that Jones inclusions of hyper-finite factors of type  $II_1$  are somehow related to the hierarchy of Planck constants. The basic motivation for this belief has been that

WCW Clifford algebra provides a canonical example of hyper-finite factor of type  $II_1$  and that Jones inclusion of these Clifford algebras is excellent candidate for a first principle description of finite measurement resolution.

Consider the inclusion  $\mathcal{N} \subset \mathcal{M}$  of hyper-finite factors of type  $II_1$  [K24]. A deep result is that one can express  $\mathcal{M}$  as  $\mathcal{N} : \mathcal{M}$ -dimensional module over  $\mathcal{N}$  with fractal dimension  $\mathcal{N} : \mathcal{M} = B_n$ .  $\sqrt{b_n}$  represents the dimension of a space of spinor space renormalized from the value 2 corresponding to  $n = \infty$  down to  $\sqrt{b_n} = 2\cos(\pi/n)$  varying thus in the range  $[1, 2]$ .  $B_n$  in turn would represent the dimension of the corresponding Clifford algebra. The interpretation is that finite measurement resolution introduces correlations between components of quantum spinor implying effective reduction of the dimension of quantum spinors providing a description of the factor space  $\mathcal{N}/\mathcal{M}$ .

This would suggest that somehow the hierarchy of Planck constants must represent finite measurement resolution and since phase factors coming as roots of unity are naturally associated with Jones inclusions the natural guess was that angular resolution and coupling constant evolution associated with it is in question. This picture would suggest that the realization of the hierarchy of Planck constant in terms of a book like structure of generalized imbedding space provides also a geometric realization for a hierarchy of Jones inclusions.

The notion of number theoretic braid and realization that the modified Dirac operator has only finite number of generalized eigenmodes -thanks to the vacuum degeneracy of Kähler action- finally led to the understanding how the notion of finite measurement resolution is coded to the Kähler action and the realized in practice by second quantization of induced spinor fields and how these spinor fields endowed with q-anti-commutation relations give rise to a representations of finite-quantum dimensional factor spaces  $\mathcal{N}/\mathcal{M}$  associated with the hierarchy of Jones inclusions having generalized imbedding space as space-time correlate. This means enormous simplification since infinite-dimensional spinor fields in infinite-dimensional world of classical worlds are replaced with finite-quantum-dimensional spinor fields in discrete points sets provided by number theoretic braids.

The study of a concrete model for Jones inclusions in terms of finite subgroups  $G$  of  $SU(2)$  defining sub-algebras of infinite-dimensional Clifford algebra as fixed point sub-algebras leads to what looks like a correct track concerning the understanding of quantization of Planck constants.

The ADE diagrams of  $A_n$  and  $D_{2n}$  characterize cyclic and dihedral groups whereas those of  $E_6$  and  $E_8$  characterize tetrahedral and icosahedral groups. This approach leads to the hypothesis that the scaling factor of Planck constant assignable to Poincare (color) algebra corresponds to the order of the maximal cyclic subgroup of  $G_b \subset SU(2)$  ( $G_a \subset SL(2, C)$ ) acting as symmetry of space-time sheet in  $CP_2$  (CD) degrees of freedom. It predicts arbitrarily large CD and  $CP_2$  Planck constants in the case of  $A_n$  and  $D_{2n}$  under rather general assumptions.

There are two manners for how  $G_a$  and  $G_b$  can act as symmetries corresponding to  $G_i$  coverings and factors spaces. These coverings and factor spaces are singular and associated with spaces  $\mathcal{C}D \setminus M^2$  and  $CP_2 \setminus S_I^2$ , where  $S_I^2$  is homologically trivial geodesic sphere of  $CP_2$ . The physical interpretation is that  $M^2$  and  $S_I^2$  fix preferred quantization axes for energy and angular moment and color quantum numbers so that also a connection with quantum measurement theory emerges.

## 3.2 The Vision

A brief summary of the basic vision behind the generalization of the imbedding space concept needed to realize the hierarchy of Planck constants is in order before going to the detailed representation.

1. The hierarchy of Planck constants cannot be realized without generalizing the notions of imbedding space and space-time because particles with different values of Planck constant cannot appear in the same interaction vertex. Some kind of book like structure for the generalized imbedding space forced also by p-adicization but in different sense is suggestive. Both  $M^4$  and  $CP_2$  factors would have the book like structure so that a Cartesian product of books would be in question.
2. The study of Schrödinger equation suggests that Planck constant corresponds to a scaling factor of CD metric whose value labels different pages of the book. The scaling of  $M^4$

coordinate so that original metric results in CD factor is possible so that the interpretation for scaled up value of  $\hbar$  is as scaling of the size of causal diamond CD.

3. The light-like 3-surfaces having their 2-D and light-boundaries of CD are in a key role in the realization of zero energy states, and the infinite-D spaces of light-like 3-surfaces inside scaled variants of CD define the fundamental building brick of WCW (world of classical worlds). Since the scaling of CD does not simply scale space-time surfaces the effect of scaling on classical and quantum dynamics is non-trivial and a coupling constant evolution results and the coding of radiative corrections to the geometry of space-time sheets becomes possible. The basic geometry of CD suggests that the allowed sizes of CD come in the basic sector  $\hbar = \hbar_0$  as powers of two. This predicts p-adic length scale hypothesis and lead to number theoretically universal discretized p-adic coupling constant evolution. Since the scaling is accompanied by a formation of singular coverings and factor spaces, different scales are distinguished at the level of topology. p-Adic length scale hierarchy affords similar characterization of length scales in terms of effective topology.
4. The idea that TGD Universe is quantum critical in some sense is one of the key postulates of quantum TGD. The basic ensuing prediction is that Kähler coupling strength is analogous to critical temperature. Quantum criticality in principle fixes the p-adic evolution of various coupling constants also the value of gravitational constant. The exact realization of quantum criticality would be in terms of critical sub-manifolds of  $M^4$  and  $CP_2$  common to all sectors of the generalized imbedding space. Quantum criticality of TGD Universe means that the two kinds of number theoretic braids assignable to  $M^4$  and  $CP_2$  projections of the partonic 2-surface belong by the very definition of number theoretic braids to these critical sub-manifolds. At the boundaries of CD associated with positive and negative energy parts of zero energy state in a given time scale partonic two-surfaces belong to a fixed page of the Big Book whereas light-like 3-surface decomposes to regions corresponding to different values of Planck constant much like matter decomposes to several phases at criticality.

The connection with Jones inclusions was originally a purely heuristic guess, and it took half decade to really understand why and how they are involved. The notion of measurement resolution is the key concept.

1. The key observation is that Jones inclusions are characterized by a finite subgroup  $G \subset SU(2)$  and the this group also characterizes the singular covering or factor spaces associated with CD or  $CP_2$  so that the pages of generalized imbedding space could indeed serve as correlates for Jones inclusions.
2. The dynamics of Kähler action realizes finite measurement resolution in terms of finite number of modes of the induced spinor field automatically implying cutoffs to the representations of various super-conformal algebras typical for the representations of quantum groups associated with Jones inclusions. The interpretation of the Clifford algebra spanned by the fermionic oscillator operators is as a realization for the concept of the factor space  $\mathcal{N}/\mathcal{M}$  of hyper-finite factors of type II<sub>1</sub> identified as the infinite-dimensional Clifford algebra  $\mathcal{N}$  of the configuration space and included algebra  $\mathcal{M}$  determining the finite measurement resolution for angle measurement in the sense that the action of this algebra on zero energy state has no detectable physical effects.  $\mathcal{M}$  takes the role of complex numbers in quantum theory and makes physics non-commutative. The resulting quantum Clifford algebra has anti-commutation relations dictated by the fractionization of fermion number so that unit becomes  $r = \hbar/\hbar_0$ .  $SU(2)$  Lie algebra transforms to its quantum variant corresponding to the quantum phase  $q = \exp(i2\pi/r)$ .
3.  $G$  invariance for the elements of the included algebra can be interpreted in terms of finite measurement resolution in the sense that action by  $G$  invariant Clifford algebra element has no detectable effects. Quantum groups realize this view about measurement resolution for angle measurement. The  $G$ -invariance of the physical states created by fermionic oscillator operators which by definition are not  $G$  invariant guarantees that quantum states as a whole have non-fractional quantum numbers so that the leakage between different pages is possible

in principle. This hypothesis is consistent with the TGD inspired model of quantum Hall effect [K17].

4. Concerning the formula for Planck constant in terms of the integers  $n_a$  and  $n_b$  characterizing orders of the maximal cyclic subgroups of groups  $G_a$  and  $G_b$  defining coverings and factor spaces associated with CD and  $CP_2$  the basic constraint is that the overall scaling of  $H$  metric has no effect on physics. What matters is the ratio of Planck constants  $r = \hbar(M^4)/\hbar(CP_2)$  appearing as a scaling factor of  $M^4$  metric. This leaves two options if one requires that the Planck constant defines a homomorphism. The model for dark gravitons suggests a unique choice between these two options but one must keep still mind open for the alternative.
5. Jones inclusions appear as two variants corresponding to  $\mathcal{N} : \mathcal{M} < 4$  and  $\mathcal{N} : \mathcal{M} = 4$ . The tentative interpretation is in terms of singular  $G$ -factor spaces and  $G$ -coverings of  $M^4$  and  $CP_2$  in some sense. The alternative interpretation assigning the inclusions to the two different geodesic spheres of  $CP_2$  would mean asymmetry between  $M^4$  and  $CP_2$  degrees of freedom and is therefore not convincing.
6. The natural question is why the hierarchy of Planck constants is needed. Is it really necessary? Number theoretic Universality suggests that this is the case. One must be able to define the notion of angle -or at least the notion of phase and of trigonometric functions- also in the p-adic context. All that one can achieve naturally is the notion of phase defined as a root of unity and introduced by allowing algebraic extension of p-adic number field by introducing the phase. In the framework of TGD inspired theory of consciousness this inspires a vision about cognitive evolution as the gradual emergence of increasingly complex algebraic extensions of p-adic numbers and involving also the emergence of improved angle resolution expressible in terms of phases  $exp(i2\pi/n)$  up to some maximum value of  $n$ . The coverings and factor spaces would realize these phases purely geometrically and quantum phases  $q$  assignable to Jones inclusions would realize them algebraically. Besides p-adic coupling constant evolution based on the hierarchy of p-adic length scales there would be coupling constant evolution with respect to  $\hbar$  and associated with angular resolution.

### **3.3 Hierarchy Of Planck Constants And The Generalization Of The Notion Of Imbedding Space**

In the following the recent view about structure of imbedding space forced by the quantization of Planck constant is summarized. The question is whether it might be possible in some sense to replace  $H$  or its Cartesian factors by their necessarily singular multiple coverings and factor spaces. One can consider two options: either  $M^4$  or the causal diamond CD. The latter one is the more plausible option from the point of view of WCW geometry.

#### **3.3.1 The evolution of physical ideas about hierarchy of Planck constants**

The evolution of the physical ideas related to the hierarchy of Planck constants and dark matter as a hierarchy of phases of matter with non-standard value of Planck constants was much faster than the evolution of mathematical ideas and quite a number of applications have been developed during last five years.

1. The starting point was the proposal of Nottale [E6] that the orbits of inner planets correspond to Bohr orbits with Planck constant  $\hbar_{gr} = GMm/v_0$  and outer planets with Planck constant  $\hbar_{gr} = 5GMm/v_0$ ,  $v_0/c \simeq 2^{-11}$ . The basic proposal [K18] was that ordinary matter condenses around dark matter which is a phase of matter characterized by a non-standard value of Planck constant whose value is gigantic for the space-time sheets mediating gravitational interaction. The interpretation of these space-time sheets could be as magnetic flux quanta or as massless extremals assignable to gravitons.
2. Ordinary particles possibly residing at these space-time sheet have enormous value of Compton length meaning that the density of matter at these space-time sheets must be very slowly varying. The string tension of string like objects implies effective negative pressure characterizing dark energy so that the interpretation in terms of dark energy might make sense [K19].

TGD predicted a one-parameter family of Robertson-Walker cosmologies with critical or over-critical mass density and the “pressure” associated with these cosmologies is negative.

3. The quantization of Planck constant does not make sense unless one modifies the view about standard space-time is. Particles with different Planck constant must belong to different worlds in the sense local interactions of particles with different values of  $\hbar$  are not possible. This inspires the idea about the book like structure of the imbedding space obtained by gluing almost copies of  $H$  together along common “back” and partially labeled by different values of Planck constant.
4. Darkness is a relative notion in this framework and due to the fact that particles at different pages of the book like structure cannot appear in the same vertex of the generalized Feynman diagram. The phase transitions in which partonic 2-surface  $X^2$  during its travel along  $X_l^3$  leaks to another page of book are however possible and change Planck constant. Particle (say photon -) exchanges of this kind allow particles at different pages to interact. The interactions are strongly constrained by charge fractionization and are essentially phase transitions involving many particles. Classical interactions are also possible. It might be that we are actually observing dark matter via classical fields all the time and perhaps have even photographed it [K20].
5. The realization that non-standard values of Planck constant give rise to charge and spin fractionization and anyonization led to the precise identification of the prerequisites of anyonic phase [K17]. If the partonic 2-surface, which can have even astrophysical size, surrounds the tip of CD, the matter at the surface is anyonic and particles are confined at this surface. Dark matter could be confined inside this kind of light-like 3-surfaces around which ordinary matter condenses. If the radii of the basic pieces of these nearly spherical anyonic surfaces - glued to a connected structure by flux tubes mediating gravitational interaction - are given by Bohr rules, the findings of Nottale [E6] can be understood. Dark matter would resemble to a high degree matter in black holes replaced in TGD framework by light-like partonic 2-surfaces with a minimum size of order Schwarzschild radius  $r_S$  of order scaled up Planck length  $l_{Pl} = \sqrt{\hbar_{gr} G} = GM$ . Black hole entropy is inversely proportional to  $\hbar$  and predicted to be of order unity so that dramatic modification of the picture about black holes is implied.
6. Perhaps the most fascinating applications are in biology. The anomalous behavior ionic currents through cell membrane (low dissipation, quantal character, no change when the membrane is replaced with artificial one) has a natural explanation in terms of dark supra currents. This leads to a vision about how dark matter and phase transitions changing the value of Planck constant could relate to the basic functions of cell, functioning of DNA and amino-acids, and to the mysteries of bio-catalysis. This leads also a model for EEG interpreted as a communication and control tool of magnetic body containing dark matter and using biological body as motor instrument and sensory receptor. One especially amazing outcome is the emergence of genetic code of vertebrates from the model of dark nuclei as nuclear strings [L1, K20], [L1].

### 3.3.2 The most general option for the generalized imbedding space

Simple physical arguments pose constraints on the choice of the most general form of the imbedding space.

1. The fundamental group of the space for which one constructs a non-singular covering space or factor space should be non-trivial. This is certainly not possible for  $M^4$ , CD,  $CP_2$ , or  $H$ . One can however construct singular covering spaces. The fixing of the quantization axes implies a selection of the sub-space  $H_4 = M^2 \times S^2 \subset M^4 \times CP_2$ , where  $S^2$  is geodesic sphere of  $CP_2$ .  $\hat{M}^4 = M^4 \setminus M^2$  and  $\hat{CP}_2 = CP_2 \setminus S^2$  have fundamental group  $Z$  since the codimension of the excluded sub-manifold is equal to two and homotopically the situation is like that for a punctured plane. The exclusion of these sub-manifolds defined by the choice of quantization axes could naturally give rise to the desired situation.



2.  $CP_2$  allows two geodesic spheres which left invariant by  $U(2)$  resp.  $SO(3)$ . The first one is homologically non-trivial. For homologically non-trivial geodesic sphere  $H_4 = M^2 \times S^2$  represents a straight cosmic string which is non-vacuum extremal of Kähler action (not necessarily preferred extremal). One can argue that the many-valuedness of  $\hbar$  is unacceptable for non-vacuum extremals so that only homologically trivial geodesic sphere  $S^2$  would be acceptable. One could go even further. If the extremals in  $M^2 \times CP_2$  can be preferred non-vacuum extremals, the singular coverings of  $M^4$  are not possible. Therefore only the singular coverings and factor spaces of  $CP_2$  over the homologically trivial geodesic sphere  $S^2$  would be possible. This however looks a non-physical outcome.
  - (a) The situation changes if the extremals of type  $M^2 \times Y^2$ ,  $Y^2$  a holomorphic surface of  $CP_3$ , fail to be hyperquaternionic. The tangent space  $M^2$  represents hypercomplex sub-space and the product of the Kähler-Dirac gamma matrices associated with the tangent spaces of  $Y^2$  should belong to  $M^2$  algebra. This need not be the case in general.
  - (b) The situation changes also if one reinterprets the gluing procedure by introducing scaled up coordinates for  $M^4$  so that metric is continuous at  $M^2 \times CP_2$  but CDs with different size have different sizes differing by the ratio of Planck constants and would thus have only piece of lower or upper boundary in common.
3. For the more general option one would have four different options corresponding to the Cartesian products of singular coverings and factor spaces. These options can be denoted by  $C - C$ ,  $C - F$ ,  $F - C$ , and  $F - F$ , where  $C$  ( $F$ ) signifies for covering (factor space) and first (second) letter signifies for CD ( $CP_2$ ) and correspond to the spaces  $(\hat{C}D \hat{\times} G_a) \times (CP_2 \hat{\times} G_b)$ ,  $(\hat{C}D \hat{\times} G_a) \times CP_2/G_b$ ,  $\hat{C}D/G_a \times (CP_2 \hat{\times} G_b)$ , and  $\hat{C}D/G_a \times CP_2/G_b$ .
4. The groups  $G_i$  could correspond to cyclic groups  $Z_n$ . One can also consider an extension by replacing  $M^2$  and  $S^2$  with its orbit under more general group  $G$  (say tetrahedral, octahedral, or icosahedral group). One expects that the discrete subgroups of  $SU(2)$  emerge naturally in this framework if one allows the action of these groups on the singular sub-manifolds  $M^2$  or  $S^2$ . This would replace the singular manifold with a set of its rotated copies in the case that the subgroups have genuinely 3-dimensional action (the subgroups which corresponds to exceptional groups in the ADE correspondence). For instance, in the case of  $M^2$  the quantization axes for angular momentum would be replaced by the set of quantization axes going through the vertices of tetrahedron, octahedron, or icosahedron. This would bring non-commutative homotopy groups into the picture in a natural manner.

### 3.3.3 About the phase transitions changing Planck constant

There are several non-trivial questions related to the details of the gluing procedure and phase transition as motion of partonic 2-surface from one sector of the imbedding space to another one.

1. How the gluing of copies of imbedding space at  $M^2 \times CP_2$  takes place? It would seem that the covariant metric of CD factor proportional to  $\hbar^2$  must be discontinuous at the singular manifold since only in this manner the idea about different scaling factor of CD metric can make sense. On the other hand, one can always scale the  $M^4$  coordinates so that the metric is continuous but the sizes of CDs with different Planck constants differ by the ratio of the Planck constants.
2. One might worry whether the phase transition changing Planck constant means an instantaneous change of the size of partonic 2-surface in  $M^4$  degrees of freedom. This is not the case. Light-likeness in  $M^2 \times S^2$  makes sense only for surfaces  $X^1 \times D^2 \subset M^2 \times S^2$ , where  $X^1$  is light-like geodesic. The requirement that the partonic 2-surface  $X^2$  moving from one sector of  $H$  to another one is light-like at  $M^2 \times S^2$  irrespective of the value of Planck constant requires that  $X^2$  has single point of  $M^2$  as  $M^2$  projection. Hence no sudden change of the size  $X^2$  occurs.
3. A natural question is whether the phase transition changing the value of Planck constant can occur purely classically or whether it is analogous to quantum tunnelling. Classical

non-vacuum extremals of Chern-Simons action have two-dimensional  $CP_2$  projection to homologically non-trivial geodesic sphere  $S^2_I$ . The deformation of the entire  $S^2_I$  to homologically trivial geodesic sphere  $S^2_{II}$  is not possible so that only combinations of partonic 2-surfaces with vanishing total homology charge (Kähler magnetic charge) can in principle move from sector to another one, and this process involves fusion of these 2-surfaces such that  $CP_2$  projection becomes single homologically trivial 2-surface. A piece of a non-trivial geodesic sphere  $S^2_I$  of  $CP_2$  can be deformed to that of  $S^2_{II}$  using 2-dimensional homotopy flattening the piece of  $S^2$  to curve. If this homotopy cannot be chosen to be light-like, the phase transitions changing Planck constant take place only via quantum tunnelling. Obviously the notions of light-like homotopies (cobordisms) are very relevant for the understanding of phase transitions changing Planck constant.

**3.3.4 How one could fix the spectrum of Planck constants?**

The question how the observed Planck constant relates to the integers  $n_a$  and  $n_b$  defining the covering and factors spaces, is far from trivial and I have considered several options. The basic physical inputs are the condition that scaling of Planck constant must correspond to the scaling of the metric of CD (that is Compton lengths) on one hand and the scaling of the gauge coupling strength  $g^2/4\pi\hbar$  on the other hand.

1. One can assign to Planck constant to both CD and  $CP_2$  by assuming that it appears in the commutation relations of corresponding symmetry algebras. Algebraist would argue that Planck constants  $\hbar(CD)$  and  $\hbar(CP_2)$  must define a homomorphism respecting multiplication and division (when possible) by  $G_i$ . This requires  $r(X) = \hbar(X)\hbar_0 = n$  for covering and  $r(X) = 1/n$  for factor space or vice versa.
2. If one assumes that  $\hbar^2(X)$ ,  $X = M^4$ ,  $CP_2$  corresponds to the scaling of the covariant metric tensor  $g_{ij}$  and performs an over-all scaling of  $H$ -metric allowed by the Weyl invariance of Kähler action by dividing metric with  $\hbar^2(CP_2)$ , one obtains the scaling of  $M^4$  covariant metric by  $r^2 \equiv \hbar^2/\hbar_0^2 = \hbar^2(M^4)/\hbar^2(CP_2)$  whereas  $CP_2$  metric is not scaled at all.
3. The condition that  $\hbar$  scales as  $n_a$  is guaranteed if one has  $\hbar(CD) = n_a\hbar_0$ . This does not fix the dependence of  $\hbar(CP_2)$  on  $n_b$  and one could have  $\hbar(CP_2) = n_b\hbar_0$  or  $\hbar(CP_2) = \hbar_0/n_b$ . The intuitive picture is that  $n_b$ -fold covering gives in good approximation rise to  $n_a n_b$  sheets and multiplies YM action action by  $n_a n_b$  which is equivalent with the  $\hbar = n_a n_b \hbar_0$  if one effectively compresses the covering to  $CD \times CP_2$ . One would have  $\hbar(CP_2) = \hbar_0/n_b$  and  $\hbar = n_a n_b \hbar_0$ . Note that the descriptions using ordinary Planck constant and coverings and scaled Planck constant but contracting the covering would be alternative descriptions.

This gives the following formulas  $r \equiv \hbar/\hbar_0 = r(M^4)/r(CP_2)$  in various cases.

$$\begin{array}{ccccc} C - C & F - C & C - F & F - F & \\ \hline r & n_a n_b & \frac{n_a}{n_b} & \frac{n_b}{n_a} & \frac{1}{n_a n_b} \end{array}$$

**3.3.5 Preferred values of Planck constants**

Number theoretic considerations favor the hypothesis that the integers corresponding to Fermat polygons constructible using only ruler and compass and given as products  $n_F = 2^k \prod_s F_s$ , where  $F_s = 2^{2^s} + 1$  are distinct Fermat primes, are favored. The reason would be that quantum phase  $q = \exp(i\pi/n)$  is in this case expressible using only iterated square root operation by starting from rationals. The known Fermat primes correspond to  $s = 0, 1, 2, 3, 4$  so that the hypothesis is very strong and predicts that p-adic length scales have satellite length scales given as multiples of  $n_F$  of fundamental p-adic length scale.  $n_F = 2^{11}$  corresponds in TGD framework to a fundamental constant expressible as a combination of Kähler coupling strength,  $CP_2$  radius and Planck length appearing in the expression for the tension of cosmic strings, and the powers of  $2^{11}$  was proposed to define favored as values of  $n_a$  in living matter [K8].

The hypothesis that Mersenne primes  $M_k = 2^k - 1$ ,  $k \in \{89, 107, 127\}$ , and Gaussian Mersennes  $M_{G,k} = (1+i)k - 1$ ,  $k \in \{113, 151, 157, 163, 167, 239, 241.. \}$  (the number theoretical miracle is that

all the four scaled up electron Compton lengths  $L_e(k) = \sqrt{5}L(k)$  with  $k \in \{151, 157, 163, 167\}$  are in the biologically highly interesting range 10 nm-2.5  $\mu\text{m}$ ) define scaled up copies of electroweak and QCD type physics with ordinary value of  $\hbar$  and that these physics are induced by dark variants of corresponding lower level physics leads to a prediction for the preferred values of  $r = 2^{k_d}$ ,  $k_d = k_i - k_j$ , and the resulting picture finds support from the ensuing models for biological evolution and for EEG [K8]. This hypothesis - to be referred to as Mersenne hypothesis - replaces the rather ad hoc proposal  $r = \hbar/\hbar_0 = 2^{11k}$  for the preferred values of Planck constant.

### 3.3.6 How Planck constants are visible in Kähler action?

$\hbar(M^4)$  and  $\hbar(CP_2)$  appear in the commutation and anti-commutation relations of various superconformal algebras. Only the ratio of  $M^4$  and  $CP_2$  Planck constants appears in Kähler action and is due to the fact that the  $M^4$  and  $CP_2$  metrics of the imbedding space sector with given values of Planck constants are proportional to the corresponding Planck. This implies that Kähler function codes for radiative corrections to the classical action, which makes possible to consider the possibility that higher order radiative corrections to functional integral vanish as one might expect at quantum criticality. For a given p-adic length scale space-time sheets with all allowed values of Planck constants are possible. Hence the spectrum of quantum critical fluctuations could in the ideal case correspond to the spectrum of  $\hbar$  coding for the scaled up values of Compton lengths and other quantal lengths and times. If so, large  $\hbar$  phases could be crucial for understanding of quantum critical superconductors, in particular high  $T_c$  superconductors.

## 4 Updated View About The Hierarchy Of Planck Constants

During last years the work with TGD proper has transformed from the discovery of brave visions to the work of clock smith. The challenge is to fill in the details, to define various notions more precisely, and to eliminate the numerous inconsistencies.

Few years has passed from the latest formulation for the hierarchy of Planck constant. The original hypothesis was that the hierarchy is real. In this formulation the imbedding space was replaced with its covering space assumed to decompose to a Cartesian product of singular finite-sheeted coverings of  $M^4$  and  $CP_2$ .

Few years ago came the realization that it could be only effective but have same practical implications. The basic observation was that the effective hierarchy need not be postulated separately but follows as a prediction from the vacuum degeneracy of Kähler action. In this formulation Planck constant at fundamental level has its standard value and its effective values come as its integer multiples so that one should write  $\hbar_{eff} = n\hbar$  rather than  $\hbar = n\hbar_0$  as I have done. For most practical purposes the states in question would behave as if Planck constant were an integer multiple of the ordinary one. It was no more necessary to assume that the covering reduces to a Cartesian product of singular coverings of  $M^4$  and  $CP_2$  but for some reason I kept this assumption.

It seems that the time is ripe for checking whether some polishing of this formulation might be needed. In particular, the work with TGD inspired quantum biology suggests a close connection between the hierarchy of Planck constants and negentropic entanglement. Also the connection with anyons and charge fractionalization has remained somewhat fuzzy [K17]. In particular, it seems that the formulation based on multi-furcations of space-time surfaces to  $N$  branches is not general enough: the  $N$  branches are very much analogous to single particle states and second quantization allowing all  $0 < n \leq N$ -particle states for given  $N$  rather than only  $N$ -particle states looks very natural: as a matter fact, this interpretation was the original one and led to the very speculative and fuzzy notion of  $N$ -atom, which I later more or less gave up. Quantum multi-furcation could be the root concept implying the effective hierarchy of Planck constants, anyons and fractional charges, and related notions- even the notions of  $N$ -nuclei,  $N$ -atoms, and  $N$ -molecules.

### 4.1 Basic Physical Ideas

The basic phenomenological rules are simple and there is no need to modify them.

1. The phases with non-standard values of effective Planck constant are identified as dark matter. The motivation comes from the natural assumption that only the particles with the same

value of effective Planck can appear in the same vertex. One can illustrate the situation in terms of the book metaphor. Imbedding spaces with different values of Planck constant form a book like structure and matter can be transferred between different pages only through the back of the book where the pages are glued together. One important implication is that light exotic charged particles lighter than weak bosons are possible if they have non-standard value of Planck constant. The standard argument excluding them is based on decay widths of weak bosons and has led to a neglect of large number of particle physics anomalies [K21].

2. Large effective or real value of Planck constant scales up Compton length - or at least de Broglie wave length - and its geometric correlate at space-time level identified as size scale of the space-time sheet assignable to the particle. This could correspond to the Kähler magnetic flux tube for the particle forming consisting of two flux tubes at parallel space-time sheets and short flux tubes at ends with length of order  $CP_2$  size.

This rule has far reaching implications in quantum biology and neuroscience since macroscopic quantum phases become possible as the basic criterion stating that macroscopic quantum phase becomes possible if the density of particles is so high that particles as Compton length sized objects overlap. Dark matter therefore forms macroscopic quantum phases. One implication is the explanation of mysterious looking quantal effects of ELF radiation in EEG frequency range on vertebrate brain:  $E = hf$  implies that the energies for the ordinary value of Planck constant are much below the thermal threshold but large value of Planck constant changes the situation. Also the phase transitions modifying the value of Planck constant and changing the lengths of flux tubes (by quantum classical correspondence) are crucial as also reconnections of the flux tubes.

The hierarchy of Planck constants suggests also a new interpretation for FQHE (fractional quantum Hall effect) [K17] in terms of anyonic phases with non-standard value of effective Planck constant realized in terms of the effective multi-sheeted covering of imbedding space: multi-sheeted space-time is to be distinguished from many-sheeted space-time.

3. In astrophysics and cosmology the implications are even more dramatic if one believes that also  $\hbar_{gr}$  corresponds to effective Planck constant interpreted as number of sheets of multifurcation. It was Nottale [E6] who first introduced the notion of gravitational Planck constant as  $\hbar_{gr} = GMm/v_0$ ,  $v_0 < 1$  has interpretation as velocity light parameter in units  $c = 1$ . This would be true for  $GMm/v_0 \geq 1$ . The interpretation of  $\hbar_{gr}$  in TGD framework is as an effective Planck constant associated with space-time sheets mediating gravitational interaction between masses  $M$  and  $m$ . The huge value of  $\hbar_{gr}$  means that the integer  $\hbar_{gr}/\hbar_0$  interpreted as the number of sheets of covering is gigantic and that Universe possesses gravitational quantum coherence in super-astronomical scales for masses which are large. This would suggest that gravitational radiation is emitted as dark gravitons which decay to pulses of ordinary gravitons replacing continuous flow of gravitational radiation.

It must be however emphasized that the interpretation of  $\hbar_{gr}$  could be different, and it will be found that one can develop an argument demonstrating how  $\hbar_{gr}$  with a correct order of magnitude emerges from the effective space-time metric defined by the anti-commutators appearing in the Kähler-Dirac equation.

4. Why Nature would like to have large effective value of Planck constant? A possible answer relies on the observation that in perturbation theory the expansion takes in powers of gauge couplings strengths  $\alpha = g^2/4\pi\hbar$ . If the effective value of  $\hbar$  replaces its real value as one might expect to happen for multi-sheeted particles behaving like single particle,  $\alpha$  is scaled down and perturbative expansion converges for the new particles. One could say that Mother Nature loves theoreticians and comes in rescue in their attempts to calculate. In quantum gravitation the problem is especially acute since the dimensionless parameter  $GMm/\hbar$  has gigantic value. Replacing  $\hbar$  with  $\hbar_{gr} = GMm/v_0$  the coupling strength becomes  $v_0 < 1$ .

## 4.2 Space-Time Correlates For The Hierarchy Of Planck Constants

The hierarchy of Planck constants was introduced to TGD originally as an additional postulate and formulated as the existence of a hierarchy of imbedding spaces defined as Cartesian products

of singular coverings of  $M^4$  and  $CP_2$  with numbers of sheets given by integers  $n_a$  and  $n_b$  and  $\hbar = n\hbar_0$ .  $n = n_a n_b$ .

With the advent of zero energy ontology, it became clear that the notion of singular covering space of the imbedding space could be only a convenient auxiliary notion. Singular means that the sheets fuse together at the boundary of multi-sheeted region. The effective covering space emerges naturally from the vacuum degeneracy of Kähler action meaning that all deformations of canonically imbedded  $M^4$  in  $M^4 \times CP_2$  have vanishing action up to fourth order in small perturbation. This is clear from the fact that the induced Kähler form is quadratic in the gradients of  $CP_2$  coordinates and Kähler action is essentially Maxwell action for the induced Kähler form. The vacuum degeneracy implies that the correspondence between canonical momentum currents  $\partial L_K / \partial(\partial_\alpha h^k)$  defining the Kähler-Dirac gamma matrices [K25] and gradients  $\partial_\alpha h^k$  is not one-to-one. Same canonical momentum current corresponds to several values of gradients of imbedding space coordinates. At the partonic 2-surfaces at the light-like boundaries of CD carrying the elementary particle quantum numbers this implies that the two normal derivatives of  $h^k$  are many-valued functions of canonical momentum currents in normal directions.

Multi-furcation is in question and multi-furcations are indeed generic in highly non-linear systems and Kähler action is an extreme example about non-linear system (see **Fig. <http://tgdtheory.fi/appfigures/planckhierarchy.jpg>** or **Fig. ??** in the appendix of this book). What multi-furcation means in quantum theory? The branches of multi-furcation are obviously analogous to single particle states. In quantum theory second quantization means that one constructs not only single particle states but also the many particle states formed from them. At space-time level single particle states would correspond to  $N$  branches  $b_i$  of multi-furcation carrying fermion number. Two-particle states would correspond to 2-fold covering consisting of 2 branches  $b_i$  and  $b_j$  of multi-furcation.  $N$ -particle state would correspond to  $N$ -sheeted covering with all branches present and carrying elementary particle quantum numbers. The branches coincide at the partonic 2-surface but since their normal space data are different they correspond to different tensor product factors of state space. Also now the factorization  $N = n_a n_b$  occurs but now  $n_a$  and  $n_b$  would relate to branching in the direction of space-like 3-surface and light-like 3-surface rather than  $M^4$  and  $CP_2$  as in the original hypothesis.

In light of this the working hypothesis adopted during last years has been too limited: for some reason I ended up to propose that only  $N$ -sheeted covering corresponding to a situation in which all  $N$  branches are present is possible. Before that I quite correctly considered more general option based on intuition that one has many-particle states in the multi-sheeted space. The erratic form of the working hypothesis has not been used in applications.

Multi-furcations relate closely to the quantum criticality of Kähler action. Feigenbaum bifurcations represent a toy example of a system which via successive bifurcations approaches chaos. Now more general multi-furcations in which each branch of given multi-furcation can multi-furcate further, are possible unless one poses any additional conditions. This allows to identify additional aspect of the geometric arrow of time. Either the positive or negative energy part of the zero energy state is “prepared” meaning that single  $n$ -sub-furcations of  $N$ -furcation is selected. The most general state of this kind involves superposition of various  $n$ -sub-furcations.

### 4.3 The Relationship To The Original View About The Hierarchy Of Planck Constants

Originally the hierarchy of Planck constant was assumed to correspond to a book like structure having as pages the  $n$ -fold coverings of the imbedding space for various values of  $n$  serving therefore as a page number. The pages are glued together along a 4-D “back” at which the branches of  $n$ -furcations are degenerate. This leads to a very elegant picture about how the particles belonging to the different pages of the book interact. All vertices are local and involve only particles with the same value of Planck constant: this is enough for darkness in the sense of particle physics. The interactions between particles belonging to different pages involve exchange of a particle travelling from page to another through the back of the book and thus experiencing a phase transition changing the value of Planck constant.

Is this picture consistent with the picture based on  $n$ -furcations? This seems to be the case. The conservation of energy in  $n$ -furcation in which several sheets are realized simultaneously is consistent with the conservation of classical conserved quantities only if the space-time sheet before

$n$ -furcation involves  $n$  identical copies of the original space-time sheet or if the Planck constant is  $h_{eff} = nh$ . This kind of degenerate many-sheetedness is encountered also in the case of branes. The first option means an  $n$ -fold covering of imbedding space and  $h_{eff}$  is indeed effective Planck constant. Second option means a genuine quantization of Planck constant due to the fact the value of Kähler coupling strength  $\alpha_K = g_K^2/4\pi\hbar_{eff}$  is scaled down by  $1/n$  factor. The scaling of Planck constant consistent with classical field equations since they involve  $\alpha_K$  as an overall multiplicative factor only.

#### 4.4 Basic Phenomenological Rules Of Thumb In The New Framework

It is important to check whether or not the refreshed view about dark matter is consistent with existent rules of thumb.

1. The interpretation of quantized multi-furcations as WCW anyons explains also why the effective hierarchy of Planck constants defines a hierarchy of phases which are dark relative to each other. This is trivially true since the phases with different number of branches in multi-furcation correspond to disjoint regions of WCW so that the particles with different effective value of Planck constant cannot appear in the same vertex.
2. The phase transitions changing the value of Planck constant are just the multi-furcations and can be induced by changing the values of the external parameters controlling the properties of preferred extremals. Situation is very much the same as in any non-linear system.
3. In the case of massless particles the scaling of wavelength in the effective scaling of  $\hbar$  can be understood if dark  $n$ -photons consist of  $n$  photons with energy  $E/n$  and wavelength  $n\lambda$ .
4. For massive particle it has been assumed that masses for particles and they dark counterparts are same and Compton wavelength is scaled up. In the new picture this need not be true. Rather, it would seem that wave length are same as for ordinary electron.

On the other hand, p-adic thermodynamics predicts that massive elementary particles are massless most of the time. ZEO predicts that even virtual wormhole throats are massless. Could this mean that the picture applying on massless particle should apply to them at least at relativistic limit at which mass is negligible. This might be the case for bosons but for fermions also fermion number should be fractionalized and this is not possible in the recent picture. If one assumes that the  $n$ -electron has same mass as electron, the mass for dark single electron state would be scaled down by  $1/n$ . This does not look sensible unless the p-adic length defined by prime is scaled down by this fact in good approximation.

This suggests that for fermions the basic scaling rule does not hold true for Compton length  $\lambda_c = \hbar/m$ . Could it however hold for de-Broglie lengths  $\lambda = \hbar/p$  defined in terms of 3-momentum? The basic overlap rule for the formation of macroscopic quantum states is indeed formulated for de Broglie wave length. One could argue that an  $1/N$ -fold reduction of density that takes place in the de-localization of the single particle states to the  $N$  branches of the cover, implies that the volume per particle increases by a factor  $N$  and single particle wave function is de-localized in a larger region of 3-space. If the particles reside at effectively one-dimensional 3-surfaces - say magnetic flux tubes - this would increase their de Broglie wave length in the direction of the flux tube and also the length of the flux tube. This seems to be enough for various applications.

One important notion in TGD inspired quantum biology is dark cyclotron state.

1. The scaling  $\hbar \rightarrow k\hbar$  in the formula  $E_n = (n + 1/2)\hbar eB/m$  implies that cyclotron energies are scaled up for dark cyclotron states. What this means microscopically has not been obvious but the recent picture gives a rather clearcut answer. One would have  $k$ -particle state formed from cyclotron states in  $N$ -fold branched cover of space-time surface. Each branch would carry magnetic field  $B$  and ion or electron. This would give a total cyclotron energy equal to  $kE_n$ . These cyclotron states would be excited by  $k$ -photons with total energy  $E = khf$  and for large enough value of  $k$  the energies involved would be above thermal threshold. In the case of  $Ca^{++}$  one has  $f = 15$  Hz in the field  $B_{end} = .2$  Gauss. This means that the value

of  $\hbar$  is at least the ratio of thermal energy at room temperature to  $E = hf$ . The thermal frequency is of order  $10^{12}$  Hz so that one would have  $k \simeq 10^{11}$ . The number branches would be therefore rather high.

2. It seems that this kinds of states which I have called cyclotron Bose-Einstein condensates could make sense also for fermions. The dark photons involved would be Bose-Einstein condensates of  $k$  photons and wall of them would be simultaneously absorbed. The biological meaning of this would be that a simultaneous excitation of large number of atoms or molecules can take place if they are localized at the branches of  $N$ -furcation. This would make possible coherent macroscopic changes. Note that also Cooper pairs of electrons could be  $n = 2$ -particle states associated with  $N$ -furcation.

There are experimental findings suggesting that photosynthesis involves de-localized excitations of electrons and it is interesting so see whether this could be understood in this framework.

1. The TGD based model relies on the assumption that cyclotron states are involved and that dark photons with the energy of visible photons but with much longer wavelength are involved. Single electron excitations (or single particle excitations of Cooper pairs) would generate negentropic entanglement (see **Fig.** <http://tgdtheory.fi/appfigures/cat.jpg> or **Fig.** ?? in the appendix of this book) automatically.
2. If cyclotron excitations are the primary ones, it would seem that they could be induced by dark  $n$ -photons exciting all  $n$  electrons simultaneously.  $n$ -photon should have energy of a visible photon. The number of cyclotron excited electrons should be rather large if the total excitation energy is to be above thermal threshold. In this case one could not speak about cyclotron excitation however. This would require that solar photons are transformed to  $n$ -photons in  $N$ -furcation in biosphere.
3. Second - more realistic looking - possibility is that the incoming photons have energy of visible photon and are therefore  $n = 1$  dark photons de-localized to the branches of the  $N$ -furcation. They would induce de-localized single electron excitation in WCW rather than 3-space.

## 4.5 Charge Fractionalization And Anyons

It is easy to see how the effective value of Planck constant as an integer multiple of its standard value emerges for multi-sheeted states in second quantization. At the level of Kähler action one can assume that in the first approximation the value of Kähler action for each branch is same so that the total Kähler action is multiplied by  $n$ . This corresponds effectively to the scaling  $\alpha_K \rightarrow \alpha_K/n$  induced by the scaling  $\hbar_0 \rightarrow n\hbar_0$ .

Also effective charge fractionalization and anyons emerge naturally in this framework.

1. In the ordinary charge fractionalization the wave function decomposes into sharply localized pieces around different points of 3-space carrying fractional charges summing up to integer charge. Now the same happens at the level of WCW ("world of classical worlds") rather than 3-space meaning that wave functions in  $E^3$  are replaced with wave functions in the space-time of 3-surfaces (4-surfaces by holography implied by General Coordinate Invariance) replacing point-like particles. Single particle wave function in WCW is a sum of  $N$  sharply localized contributions: localization takes place around one particular branch of the multi-sheeted space time surface. Each branch carries a fractional charge  $q/N$  for teh analogs of plane waves.

Therefore all quantum numbers are additive and fractionalization is only effective and observable in a localization of wave function to single branch occurring with probability  $p = 1/N$  from which one can deduce that charge is  $q/N$ .

2. The is consistent with the proposed interpretation of dark photons/gravitons since they could carry large spin and this kind of situation could decay to bunches of ordinary photons/gravitons. It is also consistent with electromagnetic charge fractionization and fractionization of spin.

3. The original - and it seems wrong - argument suggested what might be interpreted as a genuine fractionization for orbital angular momentum and also of color quantum numbers, which are analogous to orbital angular momentum in TGD framework. The observation was that a rotation through  $2\pi$  at space-time level moving the point along space-time surface leads to a new branch of multi-furcation and  $N + 1$ : th branch corresponds to the original one. This suggests that angular momentum fractionization should take place for  $M^4$  angle coordinate  $\phi$  because for it  $2\pi$  rotation could lead to a different sheet of the effective covering.

The orbital angular momentum eigenstates would correspond to waves  $\exp(i\phi m/N)$ ,  $m = 0, 2, \dots, N - 1$  and the maximum orbital angular momentum would correspond the sum  $\sum_{m=0}^{N-1} m/N = (N - 1)/2$ . The sum of spin and orbital angular momentum be therefore fractional.

The different prediction is due to the fact that rotations are now interpreted as flows rotating the points of 3-surface along 3-surface rather than rotations of the entire partonic surface in imbedding space. In the latter interpretation the rotation by  $2\pi$  does nothing for the 3-surface. Hence fractionization for the total charge of the single particle states does not take place unless one adopts the flow interpretation. This view about fractionization however leads to problems with fractionization of electromagnetic charge and spin for which there is evidence from fractional quantum Hall effect.

## 4.6 Negentropic Entanglement Between Branches Of Multi-Furcations

The application of negentropic entanglement and effective hierarchy of Planck constants to photosynthesis and metabolism [K11] suggests that these two notions might be closely related. Negentropic entanglement is possible for rational (and even algebraic) entanglement probabilities. If one allows number theoretic variant of Shannon entropy based on the p-adic norm for the probability appearing as argument of logarithm [K12], it is quite possible to have negative entanglement entropy and the interpretation is as genuine information carried by entanglement. The superposition of state pairs  $a_i \otimes b_i$  in entangled state would represent instances of a rule. In the case of Schrödinger cat the rule states that it is better to not open the bottle: understanding the rule consciously however requires that cat is somewhat dead! Entanglement provides information about the relationship between two systems. Shannon entropy represents lack of information about single particle state.

Negentropic entanglement would replace metabolic energy as the basic quantity making life possible. Metabolic energy could generate negentropic entanglement by exciting biomolecules to negentropically entangled states. ATP providing the energy for generating the metabolic entanglement could also itself carry negentropic entanglement, and transfer it to the target by the emission of large  $\hbar$  photons.

How the large  $\hbar$  photons could carry negentropic entanglement? There are several options to consider and at this stage it is not possible to pinpoint anyone of them as the only possible one. Several of them could also be realized.

1. In zero energy ontology large  $\hbar$  photons could carry the negentropic entanglement as entanglement between positive and negative energy parts of the photon state.
2. The negentropic entanglement of large  $\hbar$  photon could be also associated with its positive or energy part or both. Large  $\hbar_{eff} = n\hbar$  photon with  $n$ -fold energy  $E = n \times hf$  is  $n$ -sheeted structure consisting of  $n$ -photons with energy  $E = hf$  de-localized in the discrete space formed by the  $N$  space-time sheets. The  $n$  single photon states can entangle and since the branches effectively form a discrete space, rational and algebraic entanglement is very natural. There are many options for how this could happen. For instance, for  $N$ -fold branching the superposition of all  $N!/(N - n)!n!$  states obtained by selecting  $n$  branches are possible and the resulting state is entangled state. If this interpretation is correct, the vacuum degeneracy and multi-furcations implied by it would be the quintessence of life.
3. A further very attractive possibility discovered quite recently is that large  $\hbar_{eff} = n\hbar$  is closely related to the negentropic entanglement between the states of *two*  $n$ -furcated - that is dark - space-time sheets. In the most recent formulation negentropic entanglement corresponds



to a state characterized by  $n \times n$  identity matrix resulting from the measurement of density matrix. The number theoretic negentropy is positive for primes dividing  $p$  and there is unique prime for which it is maximal.

The identification of negentropic entanglement as entanglement between branches of a multifurcation is not the only possible option.

1. One proposal is that non-localized single particle excitations of cyclotron condensate at magnetic flux tubes give rise to negentropic entanglement relevant to living matter. Dark photons could transfer the negentropic entanglement possibly assignable to electron pairs of ATP molecule.

The negentropic entanglement associated with cyclotron condensate could be associated with the branches of the large  $\hbar$  variant of the condensate. In this case single particle excitation would not be sum of single particle excitations at various positions of 3-space but at various sheet of covering representing points of WCW. If each of the  $n$  branches carries  $1/n$ : th part of electron one would have an anyonic state in WCW.

2. One can also make a really crazy question. Could it be that ATP and various bio-molecules form  $n$ -particle states at the  $n$ -sheet of  $n$ -furcation and that the bio-chemistry involves simultaneous reactions of large numbers of biomolecules at these sheets? If so, the chemical reactions would take place as large number of copies.

Note that in this picture the breaking of time reversal symmetry [K2] in the presence of metabolic energy feed would be accompanied by evolution involving repeated multi-furcations leading to increased complexity. TGD based view about the arrow of time implies that for a given CD this evolution has definite direction of time. At the level of ensemble it implies second law but at the level of individual system means increasing complexity.

## 4.7 Dark Variants Of Nuclear And Atomic Physics

During years I have in rather speculative spirit considered the possibility of dark variants of nuclear and atomic - and perhaps even molecular physics. Also the notion of dark cyclotron state is central in the quantum model of living matter. One such notion is the idea that dark nucleons could realize vertebrate genetic code [K23].

Before the real understanding what charge fractionization means it was possible to imagine several variants of say dark atoms depending on whether both nuclei and electrons are dark or whether only electrons are dark and genuinely fractionally charged. The recent picture however fixes these notions completely. Basic building bricks are just ordinary nuclei and atoms and they form  $n$ -particle states associated with  $n$ -branches of  $N$ -furcation with  $n = 1, \dots, N$ . The fractionization for a single particle state de-localized completely to the discrete space of  $N$  branches as the analog of plane wave means that single branch carriers charge  $1/N$ .

The new element is the possibility of  $n$ -particle states populating  $n$  branches of the  $N$ -furcation: note that there is superposition over the states corresponding to different selections of these  $n$  branches.  $N - k$  and  $k$ -nuclei/atoms are in sense conjugates of each other and they can fuse to form  $N$ -nuclei/ $N$ -atoms which in fermionic case are analogous to Fermi sea with all states filled.

Bio-molecules seem to obey symbolic dynamics which does not depend much on the chemical properties: this has motivated various linguistic metaphors applied in bio-chemistry to describe the interactions between DNA and related molecules. This motivated the wild speculation was that  $N$ -atoms and even  $N$ -molecules could make possible the emergence of symbolic representations with  $n \leq N$  serving as a name of atom/molecule and that  $k$ - and  $N - k$  atom/molecule would be analogous to opposite sexes in that there would be strong tendency for them to fuse together to form  $N$ -atom/-molecule. For instance, in bio-catalysis  $k$ - and  $N - k$ -atoms/molecules would be paired. The recent picture about  $n$  and  $N - k$  atoms seems to be consistent with these speculations which I had already given up as too crazy. It is difficult to avoid even the speculation that bio-chemistry could replace chemical reactions with their  $n$ -multiples. Synchronized quantum jumps would allow to avoid the disastrous effects of state function reductions on quantum coherence. The second manner to say the same thing is that the effective value of Planck constant is large.

## 4.8 What About The Relationship Of Gravitational Planck Constant To Ordinary Planck Constant?

Gravitational Planck constant is given by the expression  $\hbar_{gr} = GMm/v_0$ , where  $v_0 < 1$  has interpretation as velocity parameter in the units  $c = 1$ . Can one interpret also  $\hbar_{gr}$  as effective value of Planck constant so that its values would correspond to multi-furcation with a gigantic number of sheets. This does not look reasonable.

Could one imagine any other interpretation for  $\hbar_{gr}$ ? Could the two Planck constants correspond to inertial and gravitational dichotomy for four-momenta making sense also for angular momentum identified as a four-vector? Could gravitational angular momentum and the momentum associated with the flux tubes mediating gravitational interaction be quantized in units of  $\hbar_{gr}$  naturally?

1. Gravitational four-momentum can be defined as a projection of the  $M^4$ -four-momentum to space-time surface. It's length can be naturally defined by the effective metric  $g_{eff}^{\alpha\beta}$  defined by the anti-commutators of the modified gamma matrices. Gravitational four-momentum appears as a measurement interaction term in the Kähler-Dirac action and can be restricted to the space-like boundaries of the space-time surface at the ends of CD and to the light-like orbits of the wormhole throats and which induced 4- metric is effectively 3-dimensional.
2. At the string world sheets and partonic 2-surfaces the effective metric degenerates to 2-D one. At the ends of braid strands representing their intersection, the metric is effectively 4-D. Just for definiteness assume that the effective metric is proportional to the  $M^4$  metric or rather - to its  $M^2$  projection:  $g_{eff}^{kl} = K^2 m^{kl}$ .

One can express the length squared for momentum at the flux tubes mediating the gravitational interaction between massive objects with masses  $M$  and  $m$  as

$$g_{eff}^{\alpha\beta} p_\alpha p_\beta = g_{eff}^{\alpha\beta} \partial_\alpha h^k \partial_\beta h^l p_k p_l \equiv g_{eff}^{kl} p_k p_l = n^2 \frac{\hbar^2}{L^2} . \quad (4.1)$$

Here  $L$  would correspond to the length of the flux tube mediating gravitational interaction and  $p_k$  would be the momentum flowing in that flux tube.  $g_{eff}^{kl} = K^2 m^{kl}$  would give

$$p^2 = \frac{n^2 \hbar^2}{K^2 L^2} .$$

$\hbar_{gr}$  could be identified in this simplified situation as  $\hbar_{gr} = \hbar/K$ .

3. Nottale's proposal requires  $K = GMm/v_0$  for the space-time sheets mediating gravitational interacting between massive objects with masses  $M$  and  $m$ . This gives the estimate

$$p_{gr} = \frac{GMm}{v_0} \frac{1}{L} . \quad (4.2)$$

For  $v_0 = 1$  this is of the same order of magnitude as the exchanged momentum if gravitational potential gives estimate for its magnitude.  $v_0$  is of same order of magnitude as the rotation velocity of planet around Sun so that the reduction of  $v_0$  to  $v_0 \simeq 2^{-11}$  in the case of inner planets does not mean that the propagation velocity of gravitons is reduced.

4. Nottale's formula requires that the order of magnitude for the components of the energy momentum tensor at the ends of braid strands at partonic 2-surface should have value  $GMm/v_0$ . Einstein's equations  $T = \kappa G + \Lambda g$  give a further constraint. For the vacuum solutions of Einstein's equations with a vanishing cosmological constant the value of  $\hbar_{gr}$  approaches infinity. At the flux tubes mediating gravitational interaction one expects  $T$  to be proportional to the factor  $GMm$  simply because they mediate the gravitational interaction.

5. One can consider similar equation for gravitational angular momentum:

$$g_{eff}^{\alpha\beta} L_\alpha L_\beta = g_{eff}^{kl} L_k L_l = l(l+1)\hbar^2 . \tag{4.3}$$

This would give under the same simplifying assumptions

$$L^2 = l(l+1) \frac{\hbar^2}{K^2} . \tag{4.4}$$

This would justify the Bohr quantization rule for the angular momentum used in the Bohr quantization of planetary orbits.

One might counter argue that if gravitational 4- momentum square is proportional to inertial 4-momentum squared, then Equivalence Principle implies that  $h_{gr}$  can have only single value. In ZEO however all wormhole throats - also virtual - are massless and the argument fails. The varying  $h_{gr}$  can be assigned only with longitudinal and transversal momentum squared separately but not to the ratio of gravitational and inertial 4-momenta squared which both vanish.

Maybe the proposed connection might make sense in some more refined formulation. In particular the proportionality between  $m_{eff}^{kl} = Km^{kl}$  could make sense as a quantum average. Also the fact, that the constant  $v_0$  varies, could be understood from the dynamical character of  $m_{eff}^{kl}$ .

#### 4.9 Hierarchy Of Planck Constants And Non-Determinism Of Kähler Action

Originally the hierarchy of Planck constant was inspired by empirical inputs from neuroscience, biology, and from Nottale's observations. That it is possible to understand the hierarchy in terms of non-determinism of Kähler action - the fundamental difference between TGD and quantum field theories and string models - is a victory for TGD approach (see **Fig.** <http://tgdtheory.fi/appfigures/planckhierarchy.jpg>, or **Fig. ??** in the appendix of this book).

Recall that non-determinism means that all space-time surfaces with  $CP_2$  projection, which is Lagrangian sub-manifold (at most 2-D) of  $CP_2$ , carries a vanishing induced Kähler form and is vacuum extremal. The first guess would be that there is a finite number  $n$  of space-time sheets connecting given pair of 3-surfaces at the ends of space-time surface at the light-like boundaries of causal diamond (CD). Planck constant would be given as  $h_{eff} = n \times h$  in accordance with the earlier interpretation. The degenerate extremals would have same Kähler action and conserved quantities as assumed also in the earlier approach. That the degenerate extremals co-incide at the ends of space-time surface was motivated by mathematical aesthetics in the earlier approach but finds an interpretation in terms of non-uniqueness of the preferred extremals.

It is essential that these  $n$  degrees of freedom are regarded as genuine physical degrees of freedom, which are however discrete. Negentropic entanglement and dark matter would be associated with them naturally. The effective description would be in terms of  $n$ -fold singular covering of imbedding space becoming singular at the ends of the space-time surface.

I have also assigned hierarchy of Planck constants with the quantum criticality. Quantum criticality means the existence of an entire continuous family of deformations of space-time sheet with same Kähler action and conserved quantities. The deformations would by definition vanish at the ends of space-time surface. The critical deformations would act as gauge transformations identifiable as conformal symmetries indeed expected to be presents since WCW isometries form a conformal algebra and there is also Kac-Moody type algebra present. The proposal has been that the sub-algebras of conformal algebra for which conformal weights are integer multiples of integer  $n = 1, 2, ..$  defined a hierarchy of gauge algebras so that the dynamical algebra reduces to  $n$ -dimensional one.

These two identifications seem to be mutually inconsistent. The resolution of the conflict comes from the gauge invariance. For a given Kähler action and conserved quantities there would be  $n$  conformal equivalence classes of these 4-surfaces rather than  $n$  surfaces, and one would have  $n$ -fold degeneracy but for conformal equivalence classes of 4-surfaces rather than 4-surfaces. In Minkowskian regions the degenerate preferred extremals are sheets (graphs of a map from  $M^4$  to  $CP_2$ ).

## 5 Vision About Dark Matter As Phases With Non-Standard Value Of Planck Constant

### 5.1 Dark Rules

It is useful to summarize the basic phenomenological view about dark matter.

#### 5.1.1 The notion of relative darkness

The essential difference between TGD and more conventional models of dark matter is that darkness is only relative concept.

1. Generalized imbedding space forms a book like structure and particles at different pages of the book are dark relative to each other since they cannot appear in the same vertex identified as the partonic 2-surface along which light-like 3-surfaces representing the lines of generalized Feynman diagram meet.
2. Particles at different space-time sheets act via classical gauge field and gravitational field and can also exchange gauge bosons and gravitons (as also fermions) provided these particles can leak from page to another. This means that dark matter can be even photographed [I4]. This interpretation is crucial for the model of living matter based on the assumption that dark matter at magnetic body controls matter visible to us. Dark matter can also suffer a phase transition to visible matter by leaking between the pages of the Big Book.
3. The notion of standard value  $\hbar_0$  of  $\hbar$  is not a relative concept in the sense that it corresponds to rational  $r = 1$ . In particular, the situation in which both CD and  $CP_2$  correspond to trivial coverings and factor spaces would naturally correspond to standard physics.

#### 5.1.2 Is dark matter anyonic?

In [K17] a detailed model for the Kähler structure of the generalized imbedding space is constructed. What makes this model non-trivial is the possibility that  $CP_2$  Kähler form can have gauge parts which would be excluded in full imbedding space but are allowed because of singular covering/factor-space property. The model leads to the conclusion that dark matter is anyonic if the partonic 2-surface, which can have macroscopic or even astrophysical size, encloses the tip of CD within it. Therefore the partonic 2-surface is homologically non-trivial when the tip is regarded as a puncture. Fractional charges for anyonic elementary particles imply confinement to the partonic 2-surface and the particles can escape the two surface only via reactions transforming them to ordinary particles. This would mean that the leakage between different pages of the big book is a rare phenomenon. This could partially explain why dark matter is so difficult to observe.

#### 5.1.3 Field body as carrier of dark matter

The notion of “field body” implied by topological field quantization is essential. There would be em,  $Z^0$ ,  $W$ , gluonic, and gravitonic field bodies, each characterized by its one prime. The motivation for considering the possibility of separate field bodies seriously is that the notion of induced gauge field means that all induced gauge fields are expressible in terms of four  $CP_2$  coordinates so that only single component of a gauge potential allows a representation as and independent field quantity. Perhaps also separate magnetic and electric field bodies for each interaction and identifiable as flux quanta must be considered. This kind of separation requires that the fermionic content of the flux quantum (say fermion and anti-fermion at the ends of color flux tube) is such that it conforms with the quantum numbers of the corresponding boson.

What is interesting that the conceptual separation of interactions to various types would have a direct correlate at the level of space-time topology. From a different perspective inspired by the general vision that many-sheeted space-time provides symbolic representations of quantum physics, the very fact that we make this conceptual separation of fundamental interactions could reflect the topological separation at space-time level.

p-Adic mass calculations for quarks encourage to think that the p-adic length scale characterizing the mass of particle is associated with its electromagnetic body and in the case of neutrinos

with its  $Z^0$  body.  $Z^0$  body can contribute also to the mass of charged particles but the contribution would be small. It is also possible that these field bodies are purely magnetic for color and weak interactions. Color flux tubes would have exotic fermion and anti-fermion at their ends and define colored variants of pions. This would apply not only in the case of nuclear strings but also to molecules and larger structures so that scaled variants of elementary particles and standard model would appear in all length scales as indeed implied by the fact that classical electro-weak and color fields are unavoidable in TGD framework.

One can also go further and distinguish between magnetic field body of free particle for which flux quanta start and return to the particle and “relative field” bodies associated with pairs of particles. Very complex structures emerge and should be essential for the understanding the space-time correlates of various interactions. In a well-defined sense they would define space-time correlate for the conceptual analysis of the interactions into separate parts. In order to minimize confusion it should be emphasized that the notion of field body used in this chapter relates to those space-time correlates of interactions, which are more or less *static* and related to the formation of *bound states*.

## 5.2 Phase Transitions Changing Planck Constant

The general picture is that p-adic length scale hierarchy corresponds to p-adic coupling constant evolution and hierarchy of Planck constants to the coupling constant evolution related to phase resolution. Both evolutions imply a book like structure of the generalized imbedding space.

### 5.2.1 Transition to large $\hbar$ phase and failure of perturbation theory

One of the first ideas was that the transition to large  $\hbar$  phase occurs when perturbation theory based on the expansion in terms of gauge coupling constant ceases to converge: Mother Nature would take care of the problems of theoretician. The transition to large  $\hbar$  phase obviously reduces the value of gauge coupling strength  $\alpha \propto 1/\hbar$  so that higher orders in perturbation theory are reduced whereas the lowest order “classical” predictions remain unchanged. A possible quantitative formulation of the criterion is that maximal 2-particle gauge interaction strength parameterized as  $Q_1 Q_2 \alpha$  satisfies the condition  $Q_1 Q_2 \alpha \simeq 1$ .

A justification for this picture would be that in non-perturbative phase large quantum fluctuations are present (as functional integral formalism suggests). At space-time level this could mean that space-time sheet is near to a non-deterministic vacuum extremal -at least if homologically trivial geodesic sphere defines the number theoretic braids. At certain critical value of coupling constant strength one expects that the transition amplitude for phase transition becomes very large. The resulting phase would be of course different from the original since typically charge fractionization would occur.

One should understand why the failure of the perturbation theory (expected to occur for  $\alpha Q_1 Q_2 > 1$ ) induces the reduction of Clifford algebra, scaling down of  $CP_2$  metric, and whether the  $G$ -symmetry is exact or only approximate. A partial understanding already exists. The discrete  $G$  symmetry and the reduction of the dimension of Clifford algebra would have interpretation in terms of a loss of degrees of freedom as a strongly bound state is formed. The multiple covering of  $M_{\pm}^4$  accompanying strong binding can be understood as an automatic consequence of  $G$ -invariance. A concrete realization for the binding could be charge fractionization which would not allow the particles bound on large light-like 3-surface to escape without transformation to ordinary particles.

Two examples perhaps provide more concrete view about this idea.

1. The proposed scenario can reproduce the huge value of the gravitational Planck constant. One should however develop a convincing argument why non-perturbative phase for the gravitating dark matter leads to a formation of  $G_a \times$  covering of  $CD \setminus M^2 \times CP_2 \setminus S_I^2$  with the huge value of  $\hbar_{eff} = n_a/n_b \simeq GM_1 M_2/v_0$ . The basic argument is that the dimensionless parameter  $\alpha_{gr} = GM_1 M_2/4\pi\hbar$  should be so small that perturbation theory works. This gives  $\hbar_{gr} \geq GM_1 M_2/4\pi$  so that order of magnitude is predicted correctly.
2. Color confinement represents the simplest example of a transition to a non-perturbative phase. In this case  $A_2$  and  $n = 3$  would be the natural option. The value of Planck constant would be 3 times higher than its value in perturbative QCD. Hadronic space-time sheets

would be 3-fold coverings of  $M_{\pm}^4$  and baryonic quarks of different color would reside on 3 separate sheets of the covering. This would resolve the color statistics paradox suggested by the fact that induced spinor fields do not possess color as spin like quantum number and by the facts that for orbifolds different quarks cannot move in independent  $CP_2$  partial waves assignable to  $CP_2$  cm degrees of freedom as in perturbative phase.

### 5.2.2 The mechanism of phase transition and selection rules

The mechanism of phase transition is at classical level similar to that for ordinary phase transitions. The partonic 2-surface decomposes to regions corresponding to difference values of  $\hbar$  at quantum criticality in such a manner that regions in which induced Kähler form is non-vanishing are contained within single page of imbedding space. It might be necessary to assume that only a region corresponding to single value of  $\hbar$  is possible for partonic 2-surfaces and  $\delta CD \times CP_2$  so that quantum criticality would be associated with the intermediate state described by the light-like 3-surface. One could also see the phase transition as a leakage of  $X^2$  from given page to another: this is like going through a closed door through a narrow slit between door and floor. By quantum criticality the points of number theoretic braid are already in the slit.

As in the case of ordinary phase transitions the allowed phase transitions must be consistent with the symmetries involved. This means that if the state is invariant under the maximal cyclic subgroups  $G_a$  and  $G_b$  then also the final state must satisfy this condition. This gives constraints to the orders of maximal cyclic subgroups  $Z_a$  and  $Z_b$  for initial and final state:  $n(Z_{a_i})$  resp.  $n(Z_{b_i})$  must divide  $n(Z_{a_f})$  resp.  $n(Z_{b_f})$  or vice versa in the case that factors of  $Z_i$  do not leave invariant the states. If this is the case similar condition must hold true for appropriate subgroups. In particular, powers of prime  $Z_{p^n}$ ,  $n = 1, 2, \dots$  define hierarchies of allowed phase transitions.

## 5.3 Coupling Constant Evolution And Hierarchy Of Planck Constants

If the overall vision is correct, quantum TGD would be characterized by two kinds of couplings constant evolutions. p-Adic coupling constant evolution would correspond to length scale resolution and the evolution with respect to Planck constant to phase resolution. Both evolution would have number theoretic interpretation.

### 5.3.1 Evolution with respect to phase resolution

The coupling constant evolution in phase resolution in p-adic degrees of freedom corresponds to emergence of algebraic extensions allowing increasing variety of phases  $\exp(i2\pi/n)$  expressible p-adically. This evolution can be assigned to the emergence of increasingly complex quantum phases and the increase of Planck constant.

One expects that quantum phases  $q = \exp(i\pi/n)$  which are expressible using only iterated square root operation are number theoretically very special since they correspond to algebraic extensions of p-adic numbers obtained by an iterated square root operation, which should emerge first. Therefore systems involving these values of  $q$  should be especially abundant in Nature. That arbitrarily high square roots are involved as becomes clear by studying the case  $n = 2^k$ :  $\cos(\pi/2^k) = \sqrt{[1 + \cos(\pi/2^{k-1})]/2}$ .

These polygons are obtained by ruler and compass construction and Gauss showed that these polygons, which could be called Fermat polygons, have  $n_F = 2^k \prod_s F_{n_s}$  sides/vertices: all Fermat primes  $F_{n_s}$  in this expression must be different. The analog of the p-adic length scale hypothesis emerges since larger Fermat primes are near a power of 2. The known Fermat primes  $F_n = 2^{2^n} + 1$  correspond to  $n = 0, 1, 2, 3, 4$  with  $F_0 = 3$ ,  $F_1 = 5$ ,  $F_2 = 17$ ,  $F_3 = 257$ ,  $F_4 = 65537$ . It is not known whether there are higher Fermat primes.  $n = 3, 5, 15$ -multiples of p-adic length scales clearly distinguishable from them are also predicted and this prediction is testable in living matter. I have already earlier considered the possibility that Fermat polygons could be of special importance for cognition and for biological information processing [K16].

This condition could be interpreted as a kind of resonance condition guaranteeing that scaled up sizes for space-time sheets have sizes given by p-adic length scales. The numbers  $n_F$  could take the same role in the evolution of Planck constant assignable with the phase resolution as Mersenne primes have in the evolution assignable to the p-adic length scale resolution.

The Dynkin diagrams of exceptional Lie groups  $E_6$  and  $E_8$  are exceptional as subgroups of rotation group in the sense that they cannot be reduced to symmetry transformations of plane. They correspond to the symmetry group  $S_4 \times Z_2$  of tetrahedron and  $A_5 \times Z_2$  of dodecahedron or its dual polytope icosahedron ( $A_5$  is 60-element subgroup of  $S_5$  consisting of even permutations). Maximal cyclic subgroups are  $Z_4$  and  $Z_5$  and thus their orders correspond to Fermat polygons. Interestingly,  $n = 5$  corresponds to minimum value of  $n$  making possible topological quantum computation using braids and also to Golden Mean.

### 5.3.2 Is there a correlation between the values of p-adic prime and Planck constant?

The obvious question is whether there is a correlation between p-adic length scale and the value of Planck constant. One-to-one correspondence is certainly excluded but loose correlation seems to exist.

1. In [K1] the information about the number theoretic anatomy of Kähler coupling strength is combined with input from p-adic mass calculations predicting  $\alpha_K$  to be the value of fine structure constant at the p-adic length scale associated with electron. One can also develop an explicit expression for gravitational constant assuming its renormalization group invariance on basis of dimensional considerations and this model leads to a model for the fraction of volume of the wormhole contact (piece of  $CP_2$  type extremal) from the volume of  $CP_2$  characterizing gauge boson and for similar volume fraction for the piece of the  $CP_2$  type vacuum extremal associated with fermion.
2. The requirement that gravitational constant is renormalization group invariant implies that the volume fraction depends logarithmically on p-adic length scale and Planck constant (characterizing quantum scale). The requirement that this fraction in the range  $(0, 1)$  poses a correlation between the rational characterizing Planck constant and p-adic length scale. In particular, for space-time sheets mediating gravitational interaction Planck constant must be larger than  $\hbar_0$  above length scale which is about .1 Angstrom. Also an upper bound for  $\hbar$  for given p-adic length scale results but is very large. This means that quantum gravitational effects should become important above atomic length scale [K1].

## 6 Some Applications

Below some applications of the hierarchy of Planck constants as a model of dark matter are briefly discussed. The range of applications varying from elementary particle physics to cosmology and I hope that this will convince the reader that the idea has strong physical motivations.

### 6.1 A Simple Model Of Fractional Quantum Hall Effect

The generalization of the imbedding space suggests that it could possible to understand fractional quantum Hall effect [D2] at the level of basic quantum TGD. This section represents the first rough model of QHE constructed for a couple of years ago is discussed. Needless to emphasize, the model represents only the basic idea and involves ad hoc assumption about charge fractionization.

Recall that the formula for the quantized Hall conductance is given by

$$\begin{aligned}\sigma &= \nu \times \frac{e^2}{h} , \\ \nu &= \frac{n}{m} .\end{aligned}\tag{6.1}$$

Series of fractions in  $\nu = 1/3, 2/5, 3/7, 4/9, 5/11, 6/13, 7/15, \dots, 2/3, 3/5, 4/7, 5/9, 6/11, 7/13, \dots, 5/3, 8/5, 11/7, 14/9, \dots, 4/3, 7/5, 10/7, 13/9, \dots, 1/5, 2/9, 3/13, \dots, 2/7, 3/11, \dots, 1/7, \dots$  with odd denominator have been observed as are also  $\nu = 1/2$  and  $\nu = 5/2$  states with even denominator [D2].

The model of Laughlin [D14] cannot explain all aspects of FQHE. The best existing model proposed originally by Jain is based on composite fermions resulting as bound states of electron and even number of magnetic flux quanta [D9]. Electrons remain integer charged but due to the

effective magnetic field electrons appear to have fractional charges. Composite fermion picture predicts all the observed fractions and also their relative intensities and the order in which they appear as the quality of sample improves.

The generalization of the notion of imbedding space suggests the possibility to interpret these states in terms of fractionized charge, spin, and electron number. There are  $2 \times 2 = 4$  combinations of covering and factors spaces of  $CP_2$  and three of them can lead to the increase of Planck constant. Besides this one can consider two options for the formula of Planck constant so that which the very meager theoretical background one can make only guesses. In the following a model based on option II for which the number of states is conserved in the phase transition changing  $\hbar$ .

1. The easiest manner to understand the observed fractions is by assuming that both CD and  $CP_2$  correspond to covering spaces so that both spin and electric charge and fermion number are fractionized. This means that  $e$  in electronic charge density is replaced with fractional charge. Quantized magnetic flux is proportional to  $e$  and the question is whether also here fractional charge appears. Assume that this does not occur.
2. With this assumption the expression for the Planck constant becomes for Option II as  $r = \hbar/\hbar_0 = n_a/n_b$  and charge and spin units are equal to  $1/n_b$  and  $1/n_a$  respectively. This gives  $\nu = nn_a/n_b$ . The values  $m = 2, 3, 5, 7, ..$  are observed. Planck constant can have arbitrarily large values. There are general arguments stating that also spin is fractionized in FQHE.
3. Both  $\nu = 1/2$  and  $\nu = 5/2$  state has been observed [D2, D5]. The fractionized charge is  $e/4$  in the latter case [D5, D3]. Since  $n_i > 3$  holds true if coverings and factor spaces are correlates for Jones inclusions, this requires  $n_a = 4$  and  $n_b = 8$  for  $\nu = 1/2$  and  $n_b = 4$  and  $n_a = 10$  for  $\nu = 5/2$ . Correct fractionization of charge is predicted. For  $n_b = 2$  also  $Z_2$  would appear as the fundamental group of the covering space. Filling fraction  $1/2$  corresponds in the composite fermion model and also experimentally to the limit of zero magnetic field [D9].  $n_b = 2$  is inconsistent with the observed fractionization of electric charge for  $\nu = 5/2$  and with the vision inspired by Jones inclusions.
4. A possible problematic aspect of the TGD based model is the experimental absence of even values of  $n_b$  except  $n_b = 2$  (Laughlin's model predicts only odd values of  $n$ ). A possible explanation is that by some symmetry condition possibly related to fermionic statistics (as in Laughlin model)  $n_a/n_b$  must reduce to a rational with an odd denominator for  $n_b > 2$ . In other words, one has  $n_a \propto 2^r$ , where  $2^r$  the largest power of 2 divisor of  $n_b$ .
5. Large values of  $n_a$  emerge as  $B$  increases. This can be understood from flux quantization. One has  $e \int BdS = n\hbar(M^4) = nn_a\hbar_0$ . By using actual fractional charge  $e_F = e/n_b$  in the flux factor would give  $e_F \int BdS = n(n_a/n_b)\hbar_0 = n\hbar$ . The interpretation is that each of the  $n_a$  sheets contributes one unit to the flux for  $e$ . Note that the value of magnetic field in given sheet is not affected so that the build-up of multiple covering seems to keep magnetic field strength below critical value.
6. The understanding of the thermal stability is not trivial. The original FQHE was observed in 80 mK temperature corresponding roughly to a thermal energy of  $T \sim 10^{-5}$  eV. For graphene the effect is observed at room temperature. Cyclotron energy for electron is (from  $f_e = 6 \times 10^5$  Hz at  $B = .2$  Gauss) of order thermal energy at room temperature in a magnetic field varying in the range 1-10 Tesla. This raises the question why the original FQHE requires so low temperature. The magnetic energy of a flux tube of length  $L$  is by flux quantization roughly  $e^2 B^2 S \sim E_c(e)m_e L$  ( $\hbar_0 = c = 1$ ) and exceeds cyclotron roughly by a factor  $L/L_e$ ,  $L_e$  electron Compton length so that thermal stability of magnetic flux quanta is not the explanation. A possible explanation is that since FQHE involves several values of Planck constant, it is quantum critical phenomenon and is characterized by a critical temperature. The differences of the energies associated with the phase with ordinary Planck constant and phases with different Planck constant would characterize the transition temperature.

As already noticed, it is possible to imagine several other options and the assumption about charge fractionization -although consistent with fractionization for  $\nu = 5/2$ , is rather ad hoc. Therefore the model can be taken as a warm-up exercise only. In [K17], where the delicacies of



Kähler structure of generalized imbedding space are discussed, also a more detailed of QHE is discussed.

## 6.2 Gravitational Bohr Orbitology

The basic question concerns justification for gravitational Bohr orbitology [K18]. The basic vision is that visible matter identified as matter with  $\hbar = \hbar_0$  ( $n_a = n_b = 1$ ) concentrates around dark matter at Bohr orbits for dark matter particles. The question is what these Bohr orbits really mean. Should one in improved approximation relate Bohr orbits to 3-D wave functions for dark matter as ordinary Bohr rules would suggest or do the Bohr orbits have some deeper meaning different from that in wave mechanics. Anyonic variants of partonic 2-surfaces with astrophysical size are a natural guess for the generalization of Bohr orbits.

### 6.2.1 Dark matter as large $\hbar$ phase

D. Da Rocha and Laurent Nottale have proposed that Schrödinger equation with Planck constant  $\hbar$  replaced with what might be called gravitational Planck constant  $\hbar_{gr} = \frac{GmM}{v_0}$  ( $\hbar = c = 1$ ).  $v_0$  is a velocity parameter having the value  $v_0 = 144.7 \pm .7$  km/s giving  $v_0/c = 4.6 \times 10^{-4}$ . This is rather near to the peak orbital velocity of stars in galactic halos. Also subharmonics and harmonics of  $v_0$  seem to appear. The support for the hypothesis coming from empirical data is impressive [K18].

Nottale and Da Rocha believe that their Schrödinger equation results from a fractal hydrodynamics. Many-sheeted space-time however suggests astrophysical systems are not only quantum systems at larger space-time sheets but correspond to a gigantic value of gravitational Planck constant. The gravitational (ordinary) Schrödinger equation -or at least Bohr rules with appropriate interpretation - would provide a solution of the black hole collapse (IR catastrophe) problem encountered at the classical level. The resolution of the problem inspired by TGD inspired theory of living matter is that it is the dark matter at larger space-time sheets which is quantum coherent in the required time scale.

### 6.2.2 Prediction for the parameter $v_0$

One of the key questions relate to the value of the parameter  $v_0$ . Before the introduction of the hierarchy of Planck constants I proposed that the value of the parameter  $v_0$  assuming that cosmic strings and their decay remnants are responsible for the dark matter. The harmonics of  $v_0$  can be understood as corresponding to perturbations replacing cosmic strings with their  $n$ -branched coverings so that tension becomes  $n$ -fold much like the replacement of a closed orbit with an orbit closing only after  $n$  turns.  $1/n$ -sub-harmonic would result when a magnetic flux tube split into  $n$  disjoint magnetic flux tubes. The planetary mass ratios can be produced with an accuracy better than 10 per cent assuming ruler and compass phases.

### 6.2.3 Further predictions

The study of inclinations (tilt angles with respect to the Earth's orbital plane) leads to a concrete model for the quantum evolution of the planetary system. Only a stepwise breaking of the rotational symmetry and angular momentum Bohr rules plus Newton's equation (or geodesic equation) are needed, and gravitational Schrödinger equation holds true only inside flux quanta for the dark matter.

1. During pre-planetary period dark matter formed a quantum coherent state on the ( $Z^0$ ) magnetic flux quanta (spherical cells or flux tubes). This made the flux quantum effectively a single rigid body with rotational degrees of freedom corresponding to a sphere or circle (full  $SO(3)$  or  $SO(2)$  symmetry).
2. In the case of spherical shells associated with inner planets the  $SO(3) \rightarrow SO(2)$  symmetry breaking led to the generation of a flux tube with the inclination determined by  $m$  and  $j$  and a further symmetry breaking, kind of an astral traffic jam inside the flux tube, generated a planet moving inside flux tube. The semiclassical interpretation of the angular momentum algebra predicts the inclinations of the inner planets. The predicted (real) inclinations are 6

(7) resp. 2.6 (3.4) degrees for Mercury resp. Venus). The predicted (real) inclination of the Earth's spin axis is 24 (23.5) degrees.

3. The  $v_0 \rightarrow v_0/5$  transition allowing to understand the radii of the outer planets in the model of Da Rocha and Nottale can be understood as resulting from the splitting of ( $Z^0$ ) magnetic flux tube to five flux tubes representing Earth and outer planets except Pluto, whose orbital parameters indeed differ dramatically from those of other planets. The flux tube has a shape of a disk with a hole glued to the Earth's spherical flux shell.

It is important to notice that effectively a multiplication  $n \rightarrow 5n$  of the principal quantum number is in question. This allows to consider also alternative explanations. Perhaps external gravitational perturbations have kicked dark matter from the orbit or Earth to  $n = 5k$ ,  $k = 2, 3, \dots, 7$  orbits: the fact that the tilt angles for Earth and all outer planets except Pluto are nearly the same, supports this explanation. Or perhaps there exist at least small amounts of dark matter at all orbits but visible matter is concentrated only around orbits containing some critical amount of dark matter and these orbits satisfy  $n \bmod 5 = 0$  for some reason.

4. A remnant of the dark matter is still in a macroscopic quantum state at the flux quanta. It couples to photons as a quantum coherent state but the coupling is extremely small due to the gigantic value of  $\hbar_{gr}$  scaling alpha by  $\hbar/\hbar_{gr}$ : hence the darkness.

The rather amazing coincidences between basic bio-rhythms and the periods associated with the states of orbits in solar system suggest that the frequencies defined by the energy levels of the gravitational Schrödinger equation might entrain with various biological frequencies such as the cyclotron frequencies associated with the magnetic flux tubes. For instance, the period associated with  $n = 1$  orbit in the case of Sun is 24 hours within experimental accuracy for  $v_0$ .

#### 6.2.4 Comparison with Bohr quantization of planetary orbits

The predictions of the generalization of the p-adic length scale hypothesis are consistent with the TGD based model for the Bohr quantization of planetary orbits and some new non-trivial predictions follow.

1. The model can explain the enormous values of gravitational Planck constant  $\hbar_{gr}/\hbar_0 = \simeq GMm/v_0 = n_a/n_b$ . The favored values of this parameter should correspond to  $n_{F_a}/n_{F_b}$  so that the mass ratios  $m_1/m_2 = n_{F_{a,1}}n_{F_{b,2}}/n_{F_{b,1}}n_{F_{a,2}}$  for planetary masses should be preferred. The general prediction  $GMm/v_0 = n_a/n_b$  is of course not testable.
2. Nottale [E6] has suggested that also the harmonics and sub-harmonics of  $\hbar_{gr}$  are possible and in fact required by the model for planetary Bohr orbits (in TGD framework this is not absolutely necessary [K18]). The prediction is that favored values of  $n$  should be of form  $n_F = 2^k \prod F_i$  such that  $F_i$  appears at most once. In Nottale's model for planetary orbits as Bohr orbits in solar system [K18]  $n = 5$  harmonics appear and are consistent with either  $n_{F,a} \rightarrow F_1 n_{F_a}$  or with  $n_{F,b} \rightarrow n_{F_b}/F_1$  if possible.

The prediction for the ratios of planetary masses can be tested. In the table below are the experimental mass ratios  $r_{exp} = m(pl)/m(E)$ , the best choice of  $r_R = [n_{F,a}/n_{F,b}] * X$ ,  $X$  common factor for all planets, and the ratios  $r_{pred}/r_{exp} = n_{F,a}(planet)n_{F,b}(Earth)/n_{F,a}(Earth)n_{F,b}(planet)$ . The deviations are at most 2 per cent.

A stronger prediction comes from the requirement that  $GMm/v_0$  equals to  $n = n_{F_a}/n_{F_b}$ ,  $n_F = 2^k \prod_k F_{n_k}$ , where  $F_i = 2^{2^i} + 1$ ,  $i = 0, 1, 2, 3, 4$  is Fibonacci prime. The fit using solar mass and Earth mass gives  $n_F = 2^{254} \times 5 \times 17$  for  $1/v_0 = 2044$ , which within the experimental accuracy equals to the value  $2^{11} = 2048$  whose powers appear as scaling factors of Planck constant in the model for living matter [K8]. For  $v_0 = 4.6 \times 10^{-4}$  reported by Nottale the prediction is by a factor 16/17.01 too small (6 per cent discrepancy).

A possible solution of the discrepancy is that the empirical estimate for the factor  $GMm/v_0$  is too large since  $m$  contains also the the visible mass not actually contributing to the gravitational

**Table 1:** Table compares the ratios  $x = m(pl)/(m(E))$  of planetary mass to the mass of Earth to prediction for these ratios in terms of integers  $n_F$  associated with Fermat polygons.  $y$  gives the best fit for the allowed factors of the known part  $y$  of the rational  $n_{F,a}/n_{F,b} = yX$  characterizing planet, and the ratios  $y/x$ . Errors are at most 2 per cent.

<i>planet</i>	<i>Me</i>	<i>V</i>	<i>E</i>	<i>M</i>	<i>J</i>
<i>y</i>	$\frac{2^{13} \times 5}{17}$	$2^{11} \times 17$	$2^9 \times 5 \times 17$	$2^8 \times 17$	$\frac{2^{23} \times 5}{7}$
<i>y/x</i>	1.01	.98	1.00	.98	1.01
<i>planet</i>	<i>S</i>	<i>U</i>	<i>N</i>	<i>P</i>	
<i>y</i>	$2^{14} \times 3 \times 5 \times 17$	$\frac{2^{21} \times 5}{17}$	$\frac{2^{17} \times 17}{3}$	$\frac{2^4 \times 17}{3}$	
<i>y/x</i>	1.01	.98	.99	.99	

force between dark matter objects whereas  $M$  is known correctly. The assumption that the dark mass is a fraction  $1/(1 + \epsilon)$  of the total mass for Earth gives

$$1 + \epsilon = \frac{17}{16} \quad (6.2)$$

in an excellent approximation. This gives for the fraction of the visible matter the estimate  $\epsilon = 1/16 \simeq 6$  per cent. The estimate for the fraction of visible matter in cosmos is about 4 per cent so that estimate is reasonable and would mean that most of planetary and solar mass would be also dark (as a matter dark energy would be in question).

That  $v_0(eff) = v_0/(1 - \epsilon) \simeq 4.6 \times 10^{-4}$  equals with  $v_0(eff) = 1/(2^7 \times F_2) = 4.5956 \times 10^{-4}$  within the experimental accuracy suggests a number theoretical explanation for the visible-to-dark fraction.

The original unconsciously performed identification of the gravitational and inertial Planck constants leads to some confusing conclusions but it seems that the new view about the quantization of Planck constants resolves these problems and allows to see  $\hbar_{gr}$  as a special case of  $\hbar_I$ .

1.  $\hbar_{gr}$  is proportional to the product of masses of interacting systems and not a universal constant like  $\hbar$ . One can however express the gravitational Bohr conditions as a quantization of circulation  $\oint v \cdot dl = n(GM/v_0)\hbar_0$  so that the dependence on the planet mass disappears as required by Equivalence Principle. This would suggest that gravitational Bohr rules relate to velocity rather than inertial momentum as is indeed natural. The quantization of circulation is consistent with the basic prediction that space-time surfaces are analogous to Bohr orbits.
2.  $\hbar_{gr}$  seems to characterize a relationship between planet and central mass and quite generally between two systems with the property that smaller system is topologically condensed at the space-time sheet of the larger system. Thus it would seem that  $\hbar_{gr}$  is not a universal constant and cannot correspond to a special value of ordinary Planck constant. Certainly this would be the case if  $\hbar_I$  is quantized as  $\lambda^k$ -multiplet of ordinary Planck constant with  $\lambda \simeq 2^{11}$ .

The recent view about the quantization of Planck constant in terms of coverings of CD seems to resolve these problems.

1. The integer quantization of Planck constants is consistent with the huge values of gravitational Planck constant within experimental resolution and the killer test for  $\hbar = \hbar_{gr}$  emerges if one takes seriously the stronger prediction  $\hbar_{gr} = n_{F,a}/n_{F,b}$ .
2. One can also regard  $\hbar_{gr}$  as ordinary Planck constant  $\hbar_{eff}$  associated with the space-time sheet along which the masses interact provided each pair  $(M, m_i)$  of masses is characterized by its own sheets. These sheets could correspond to flux tube like structures carrying the gravitational flux of dark matter. If these sheets corresponds to  $n_{F_a}$ -fold covering of CD, one can understand  $\hbar_{gr}$  as a particular instance of the  $\hbar_{eff}$ .

### 6.2.5 Quantum Hall effect and dark anyonic systems in astrophysical scales

Bohr orbitology could be understood if dark matter concentrates on 2-dimensional partonic surfaces usually assigned with elementary particles and having size of order  $CP_2$  radius. The interpretation is in terms of wormhole throats assignable to topologically condensed  $CP_2$  type extremals (fermions) and pairs of them assignable to wormhole contacts (gauge bosons). Wormhole throat defines the light-like 3-surface at which the signature of metric of space-time surface changes from Minkowskian to Euclidian.

Large value of Planck constant would allow partons with astrophysical size. Since anyonic systems are 2-dimensional, the natural idea is that dark matter corresponds to systems carrying large fermion number residing at partonic 2-surfaces of astrophysical size and that visible matter condenses around these. Not only black holes but also ordinary stars, planetary systems, and planets could correspond at the level of dark matter to atom like structures consisting of anyonic 2-surfaces which can have complex topology (flux tubes associated with planetary orbits connected by radial flux tubes to the central spherical anyonic surface). Charge and spin fractionization are key features of anyonic systems and Jones inclusions inspiring the generalization of imbedding space indeed involve quantum groups central in the modelling of anyonic systems. Hence one has could hopes that a coherent theoretical picture could emerge along these lines.

This seems to be the case. Anyons and charge and spin fractionization are discussed in detail [K17] and leads to a precise identification of the delicacies involved with the Kähler gauge potential of  $CP_2$  Kähler form in the sectors of the generalized imbedding space corresponding to various pages of boook like structures assignable to CD and  $CP_2$ . The basic outcome is that anyons correspond geometrically to partonic 2-surfaces at the light-like boundaries of CD containing the tip of CD inside them. This is what gives rise to charge fractionization and also to confinement like effects since elementary particles in anyonic states cannot as such leak to the other pages of the generalized imbedding space.  $G_a$  and  $G_b$  invariance of the states imply that fractionization occurs only at single particle level and total charge is integer valued.

This picture is much more flexible that that based on  $G_a$  symmetries of CD orbifold since partonic 2-surfaces do not possess any orbifold symmetries in CD sector anymore. In this framework various astrophysical structures such as spokes and circles would be parts of anyonic 2-surfaces with complex topology representing quantum geometrically quantum coherence in the scale of say solar system. Planets would have formed by the condensation of ordinary matter in the vicinity of the anyonic matter. This would predict stars, planetary system, and even planets to have onion-like structure consisting of shells at the level of dark matter. Similar conclusion is suggested also by purely classical model for the final state of star predicting that matter is strongly concentrated at the surface of the star [K22].

### 6.2.6 Anyonic view about blackholes

A new element to the model of black hole comes from the vision that black hole horizon as a light-like 3-surface corresponds to a light-like orbit of light-like partonic 2-surface. This allows two kinds of black holes. Fermion like black hole would correspond to a deformed  $CP_2$  type extremal which Euclidian signature of metric and topologically condensed at a space-time sheet with a Minkowskian signature. Boson like black hole would correspond to a wormhole contact connecting two space-time sheets with Minkowskian signature. Wormhole contact would be a piece deformed  $CP_2$  type extremal possessing two light-like throats defining two black hole horizons very near to each other. It does not seem absolutely necessary to assume that the interior metric of the black-hole is realized in another space-time sheet with Minkowskian signature.

Second new element relates to the value of Planck constant. For  $\hbar_{gr} = 4GM^2$  the Planck length  $L_P(\hbar) = \sqrt{\hbar G}$  equals to Schwarzschild radius and Planck mass equals to  $M_P(\hbar) = \sqrt{\hbar/G} = 2M$ . If the mass of the system is below the ordinary Planck mass:  $M \leq m_P(\hbar_0)/2 = \sqrt{\hbar_0/4G}$ , gravitational Planck constant is smaller than the ordinary Planck constant.

Black hole surface contains ultra dense matter so that perturbation theory is not expected to converge for the standard value of Planck constant but do so for gravitational Planck constant. If the phase transition increasing Planck constant is a friendly gesture of Nature making perturbation theory convergent, one expects that only the black holes for which Planck constant is such that  $GM^2/4\pi\hbar < 1$  holds true are formed. Black hole entropy -being proportional to  $1/\hbar-$  is of order

unity so that TGD black holes are not very entropic.

If the partonic 2-surface surrounds the tip of causal diamond CD, the matter at its surface is in anyonic state with fractional charges. Anyonic black hole can be seen as single gigantic elementary particle stabilized by fractional quantum numbers of the constituents preventing them from escaping from the system and transforming to ordinary visible matter. A huge number of different black holes are possible for given value of  $\hbar$  since there is infinite variety of pairs  $(n_a, n_b)$  of integers giving rise to same value of  $\hbar$ .

One can imagine that the partonic surface is not exact sphere except for ideal black holes but contains large number of magnetic flux tubes giving rise to handles. Also a pair of spheres with different radii can be considered with surfaces of spheres connected by braided flux tubes. The braiding of these handles can represent information and one can even consider the possibility that black hole can act as a topological quantum computer. There would be no sharp difference between the dark parts of black holes and those of ordinary stars. Only the volume containing the complex flux tube structures associated with the orbits of planets and various objects around star would become very small for black hole so that the black hole might code for the topological information of the matter collapsed into it.

### **6.3 Accelerating Periods Of Cosmic Expansion As PhaseTransitions Increasing The Value Of Planck Constant**

There are several pieces of evidence for accelerated expansion, which need not mean cosmological constant, although this is the interpretation adopted in [E4, E2]. Quantum cosmology predicts that astrophysical objects do not follow cosmic expansion except in jerk-wise quantum leaps increasing the value of the gravitational Planck constant. This assumption provides explanation for the apparent cosmological constant. Also planets are predicted to expand in this manner. This provides a new version of Expanding Earth theory originally postulated to explain the intriguing findings suggesting that continents have once formed a connected continent covering the entire surface of Earth but with radius which was one half of the recent one.

#### **6.3.1 The four pieces of evidence for accelerated expansion**

##### *1. Supernovas of type Ia*

Supernovas of type *Ia* define standard candles since their luminosity varies in an oscillatory manner and the period is proportional to the luminosity. The period gives luminosity and from this the distance can be deduced by using Hubble's law:  $d = cz/H_0$ ,  $H_0$  Hubble's constant. The observation was that the farther the supernova was the more dimmer it was as it should have been. In other words, Hubble's constant increased with distance and the cosmic expansion was accelerating rather than decelerating as predicted by the standard matter dominated and radiation dominated cosmologies.

##### *2. Mass density is critical and 3-space is flat*

It is known that the contribution of ordinary and dark matter explaining the constant velocity of distance stars rotating around galaxy is about 25 per cent from the critical density. Could it be that total mass density is critical?

From the anisotropy of cosmic microwave background one can deduce that this is the case. What criticality means geometrically is that 3-space defined as surface with constant value of cosmic time is flat. This reflects in the spectrum of microwave radiation. The spots representing small anisotropies in the microwave background temperature is 1 degree and this correspond to flat 3-space. If one had dark matter instead of dark energy the size of spot would be 5 degrees!

Thus in a cosmology based on general relativity cosmological constant remains the only viable option. The situation is different in TGD based quantum cosmology based on sub-manifold gravity and hierarchy of gravitational Planck constants.

##### *3. The energy density of vacuum is constant in the size scale of big voids*

It was observed that the density of dark energy would be constant in the scale of  $10^8$  light years. This length scale corresponds to the size of big voids containing galaxies at their boundaries.

#### 4. Integrated Sachs-Wolf effect

Also so called integrated Sachs-Wolf effect supports accelerated expansion. Very slow variations of mass density are considered. These correspond to gravitational potentials. Cosmic expansion tends to flatten them but mass accretion to form structures compensates this effect so that gravitational potentials are unaffected and there is no effect of CMB. Situation changes if dark matter is replaced with dark energy the accelerated expansion flattening the gravitational potentials wins the tendency of mass accretion to make them deeper. Hence if photon passes by an over-dense region, it receives a little energy. Similarly, photon loses energy when passign by an under-dense region. This effect has been observed.

### 6.3.2 Accelerated expansion in classical TGD

The minimum TGD based explanation for accelerated expansion involves only the fact that the imbeddings of critical cosmologies correspond to accelerated expansion. A more detailed model allows to understand why the critical cosmology appears during some periods.

The first observation is that critical cosmologies (flat 3-space) imbeddable to 8-D imbedding space  $H$  correspond to negative pressure cosmologies and thus to accelerating expansion. The negativity of the counterpart of pressure in Einstein tensor is due to the fact that space-time sheet is forced to be a 4-D surface in 8-D imbedding space. This condition is analogous to a force forcing a particle at the surface of 2-sphere and gives rise to what could be called constraint force. Gravitation in TGD is sub-manifold gravitation whereas in GRT it is manifold gravitation. This would be minimum interpretation involving no assumptions about what mechanism gives rise to the critical periods.

### 6.3.3 Accelerated expansion and hierarchy of Planck constants

One can go one step further and introduce the hierarchy of Planck constants. The basic difference between TGD and GRT based cosmologies is that TGD cosmology is quantum cosmology. Smooth cosmic expansion is replaced by an expansion occurring in discrete jerks corresponding to the increase of gravitational Planck constant. At space-time level this means the replacement of 8-D imbedding space  $H$  with a book like structure containing almost-copies of  $H$  with various values of Planck constant as pages glued together along critical manifold through which space-time sheet can leak between sectors with different values of  $\hbar$ . This process is the geometric correlate for the the phase transition changing the value of Planck constant.

During these phase transition periods critical cosmology applies and predicts automatically accelerated expansion. Neither genuine negative pressure due to “quintessence” nor cosmological constant is needed. Note that quantum criticality replaces inflationary cosmology and predicts a unique cosmology apart from single parameter. Criticality also explains the fluctuations in microwave temperature as long range fluctuations characterizing criticality.

### 6.3.4 Accelerated expansion and flatness of 3-cosmology

Observations 1) and 2) about super-novae and critical cosmology (flat 3-space) are consistent with this cosmology. In TGD dark energy must be replaced with dark matter because the mass density is critical during the phase transition. This does not lead to wrong sized spots since it is the increase of Planck constant which induces the accelerated expansion understandable also as a constraint force due to imbedding to  $H$ .

### 6.3.5 The size of large voids is the characteristic scale

The TGD based model in its simplest form model assigns the critical periods of expansion to large voids of size  $10^8$  ly. Also larger and smaller regions can express similar periods and dark space-time sheets are expected to obey same universal “cosmology” apart from a parameter characterizing the duration of the phase transition. Observation 3) that just this length scale defines the scale below which dark energy density is constant is consistent with TGD based model.

The basic prediction is jerk-wise cosmic expansion with jerks analogous to quantum transitions between states of atom increasing the size of atom. The discovery of large voids with size of order

$10^8$  ly but age much longer than the age of galactic large voids conforms with this prediction. One the other hand, it is known that the size of galactic clusters has not remained constant in very long time scale so that jerk-wise expansion indeed seems to occur.

### 6.3.6 Do cosmic strings with negative gravitational mass cause the phase transition inducing accelerated expansion

Quantum classical correspondence is the basic principle of quantum TGD and suggest that the effective antigravity manifested by accelerated expansion might have some kind of concrete space-time correlate. A possible correlate is super heavy cosmic string like objects at the center of large voids which have negative gravitational mass under very general assumptions. The repulsive gravitational force created by these objects would drive galaxies to the boundaries of large voids. At some state the pressure of galaxies would become too strong and induce a quantum phase transition forcing the increase of gravitational Planck constant and expansion of the void taking place much faster than the outward drift of the galaxies. This process would repeat itself. In the average sense the cosmic expansion would not be accelerating.

## 6.4 Phase Transition Changing Planck Constant And Expanding Earth Theory

TGD predicts that cosmic expansion at the level of individual astrophysical systems does not take place continuously as in classical gravitation but through discrete quantum phase transitions increasing gravitational Planck constant and thus various quantum length and time scales. The reason would be that stationary quantum states for dark matter in astrophysical length scales cannot expand. One would have the analog of atomic physics in cosmic scales. Increases of  $\hbar$  by a power of two are favored in these transitions but also other scalings are possible.

This has quite far reaching implications.

1. These periods have a highly unique description in terms of a critical cosmology for the expanding space-time sheet. The expansion is accelerating. The accelerating cosmic expansion can be assigned to this kind of phase transition in some length scale (TGD Universe is fractal). There is no need to introduce cosmological constant and dark energy would be actually dark matter.
2. The recently observed void which has same size of about  $10^8$  light years as large voids having galaxies near their boundaries but having an age which is much higher than that of the large voids, would represent one example of jerk-wise expansion.
3. This picture applies also to solar system and planets might be perhaps seen as having once been parts of a more or less connected system, the primordial Sun. The Bohr orbits for inner and outer planets correspond to gravitational Planck constant which is 5 times larger for outer planets. This suggests that the space-time sheet of outer planets has suffered a phase transition increasing the size scale by a factor of 5. Earth can be regarded either as  $n=1$  orbit for Planck constant associated with outer planets or  $n=5$  orbit for inner planetary system. This might have something to do with the very special position of Earth in planetary system. One could even consider the possibility that both orbits are present as dark matter structures. The phase transition would also explain why  $n=1$  and  $n=2$  Bohr orbits are absent and one only  $n=3, 4,$  and  $5$  are present.
4. Also planets should have experienced this kind of phase transitions increasing the radius: the increase by a factor two would be the simplest situation.

The obvious question - that I did not ask - is whether this kind of phase transition might have occurred for Earth and led from a completely granite covered Earth - Pangeia without seas - to the recent Earth. Neither it did not occur to me to check whether there is any support for a rapid expansion of Earth during some period of its history.

Situation changed when my son visited me last Saturday and told me about a Youtube video [F7] by Neal Adams, an American comic book and commercial artist who has also produced animations for geologists. We looked the amazing video a couple of times and I looked it again yesterday. The

video is very impressive artwork but in the lack of references skeptic probably cannot avoid the feeling that Neal Adams might use his highly developed animation skills to cheat you. I found also a polemic article [F1] of Adams but again the references were lacking. Perhaps the reason of polemic tone was that the concrete animation models make the expanding Earth hypothesis very convincing but geologists refuse to consider seriously arguments by a layman without a formal academic background.

#### 6.4.1 The claims of Adams

The basic claims of Adams were following.

1. The radius of Earth has increased during last 185 million years (dinosaurs [I1] appeared for about 230 million years ago) by about factor 2. If this is assumed all continents have formed at that time a single super-continent, Pangeia, filling the entire Earth surface rather than only 1/4 of it since the total area would have grown by a factor of 4. The basic argument was that it is very difficult to imagine Earth with 1/4 of surface containing granite and 3/4 covered by basalt. If the initial situation was covering by mere granite -as would look natural- it is very difficult for a believer in thermodynamics to imagine how the granite would have gathered to a single connected continent.
2. Adams claims that Earth has grown by keeping its density constant, rather than expanded, so that the mass of Earth has grown linearly with radius. Gravitational acceleration would have thus doubled and could provide a partial explanation for the disappearance of dinosaurs: it is difficult to cope in evolving environment when you get slower all the time.
3. Most of the sea floor is very young and the areas covered by the youngest basalt are the largest ones. This Adams interprets this by saying that the expansion of Earth is accelerating. The alternative interpretation is that the flow rate of the magma slows down as it recedes from the ridge where it erupts. The upper bound of 185 million years for the age of sea floor requires that the expansion period - if it is already over - lasted about 185 million years after which the flow increasing the area of the sea floor transformed to a convective flow with subduction so that the area is not increasing anymore.
4. The fact that the continents fit together - not only at the Atlantic side - but also at the Pacific side gives strong support for the idea that the entire planet was once covered by the super-continent. After the emergence of subduction theory this evidence as been dismissed.
5. I am not sure whether Adams mentions the following objections [F2]. Subduction only occurs on the other side of the subduction zone so that the other side should show evidence of being much older in the case that oceanic subduction zones are in question. This is definitely not the case. This is explained in plate tectonics as a change of the subduction direction. My explanation would be that by the symmetry of the situation both oceanic plates bend down so that this would represent new type of boundary not assumed in the tectonic plate theory.
6. As a master visualizer Adams notices that Africa and South-America do not actually fit together in absence of expansion unless one assumes that these continents have suffered a deformation. Continents are not easily deformable stuff. The assumption of expansion implies a perfect fit of *all* continents without deformation.

Knowing that the devil is in the details, I must admit that these arguments look rather convincing to me and what I learned from Wikipedia articles supports this picture.

#### 6.4.2 The critic of Adams of the subduction mechanism

The prevailing tectonic plate theory [F5] has been compared to the Copernican revolution in geology. The theory explains the young age of the seafloor in terms of the decomposition of the lithosphere to tectonic plates and the convective flow of magma to which oceanic tectonic plates participate. The magma emerges from the crests of the mid ocean ridges representing a boundary of two plates and leads to the expansion of sea floor. The variations of the polarity of Earth's



magnetic field coded in sea floor provide a strong support for the hypothesis that magma emerges from the crests.

The flow back to would take place at so called oceanic trenches [F3] near continents which represent the deepest parts of ocean. This process is known as subduction. In subduction oceanic tectonic plate bends and penetrates below the continental tectonic plate, the material in the oceanic plate gets denser and sinks into the magma. In this manner the oceanic tectonic plate suffers a metamorphosis returning back to the magma: everything which comes from Earth's interior returns back. Subduction mechanism explains elegantly formation of mountains [F4] (orogeny), earth quake zones, and associated zones of volcanic activity [F6] .

Adams is very polemic about the notion of subduction, in particular about the assumption that it generates steady convective cycle. The basic objections of Adams against subduction are following.

1. There are not enough subduction zones to allow a steady situation. According to Adams, the situation resembles that for a flow in a tube which becomes narrower. In a steady situation the flow should accelerate as it approaches subduction zones rather than slow down. Subduction zones should be surrounded by large areas of sea floor with constant age. Just the opposite is suggested by the fact that the youngest portion of sea-floor near the ridges is largest. The presence of zones at which both ocean plates bend down could improve the situation. Also jamming of the flow could occur so that the thickness of oceanic plate increases with the distance from the eruption ridge. Jamming could increase also the density of the oceanic plate and thus the effectiveness of subduction.
2. There is no clear evidence that subduction has occurred at other planets. The usual defense is that the presence of sea is essential for the subduction mechanism.
3. One can also wonder what is the mechanism that led to the formation of single super continent Pangeia covering 1/4 of Earth's surface. How probable the gathering of all separate continents to form single cluster is? The later events would suggest that just the opposite should have occurred from the beginning.

#### 6.4.3 Expanding Earth theories are not new

After I had decided to check the claims of Adams, the first thing that I learned is that Expanding Earth theory [F2], whose existence Adams actually mentions, is by no means new. There are actually many of them.

The general reason why these theories were rejected by the main stream community was the absence of a convincing physical mechanism of expansion or of growth in which the density of Earth remains constant.

1. 1888 Yarkovski postulated some sort of aether absorbed by Earth and transforming to chemical elements (TGD version of aether could be dark matter). 1909 Mantovani postulated thermal expansion but no growth of the Earth's mass.
2. Paul Dirac's idea about changing Planck constant led Pascual Jordan in 1964 to a modification of general relativity predicting slow expansion of planets. The recent measurement of the gravitational constant imply that the upper bound for the relative change of gravitational constant is 10 time too small to produce large enough rate of expansion. Also many other theories have been proposed but they are in general conflict with modern physics.
3. The most modern version of Expanding Earth theory is by Australian geologist Samuel W. Carey. He calculated that in Cambrian period (about 500 million years ago) all continents were stuck together and covered the entire Earth. Deep seas began to evolve then.

#### 6.4.4 Summary of TGD based theory of Expanding Earth

TGD based model differs from the tectonic plate model but allows subduction which cannot imply considerable back-flow of magma. Let us sum up the basic assumptions and implications.

1. The expansion is or was due to a quantum phase transition increasing the value of gravitational Planck constant and forced by the cosmic expansion in the average sense.
2. Tectonic plates do not participate to the expansion and therefore new plate must be formed and the flow of magma from the crests of mid ocean ridges is needed. The decomposition of a single plate covering the entire planet to plates to create the mid ocean ridges is necessary for the generation of new tectonic plate. The decomposition into tectonic plates is thus prediction rather than assumption.
3. The expansion forced the decomposition of Pangeia super-continent covering entire Earth for about 530 million years ago to split into tectonic plates which began to recede as new non-expanding tectonic plate was generated at the ridges creating expanding sea floor. The initiation of the phase transition generated formation of deep seas.
4. The eruption of plasma from the crests of ocean ridges generated oceanic tectonic plates which did not participate to the expansion by density reduction but by growing in size. This led to a reduction of density in the interior of the Earth roughly by a factor 1/8. From the upper bound for the age of the seafloor one can conclude that the period lasted for about 185 million years after which it transformed to convective flow in which the material returned back to the Earth interior. Subduction at continent-ocean floor boundaries and downwards double bending of tectonic plates at the boundaries between two ocean floors were the mechanisms. Thus tectonic plate theory would be more or less the correct description for the recent situation.
5. One can consider the possibility that the subducted tectonic plate does not transform to magma but is fused to the tectonic layer below continent so that it grows to an iceberg like structure. This need not lead to a loss of the successful predictions of plate tectonics explaining the generation of mountains, earthquake zones, zones of volcanic activity, etc...
6. From the video of Adams it becomes clear that the tectonic flow is East-West asymmetric in the sense that the western side is more irregular at large distances from the ocean ridge at the western side. If the magma rotates with slightly lower velocity than the surface of Earth (like liquid in a rotating vessel), the erupting magma would rotate slightly slower than the tectonic plate and asymmetry would be generated.
7. If the planet has not experienced a phase transition increasing the value of Planck constant, there is no need for the decomposition to tectonic plates and one can understand why there is no clear evidence for tectonic plates and subduction in other planets. The conductive flow of magma could occur below this plate and remain invisible.

The biological implications might provide a possibility to test the hypothesis.

1. Great steps of progress in biological evolution are associated with catastrophic geological events generating new evolutionary pressures forcing new solutions to cope in the new situation. Cambrian explosion indeed occurred about 530 years ago (the book "Wonderful Life" of Stephen Gould [17] explains this revolution in detail) and led to the emergence of multicellular creatures, and generated huge number of new life forms living in seas. Later most of them suffered extinction: large number of phylae and groups emerged which are not present nowadays.

Thus Cambrian explosion is completely exceptional as compared to all other dramatic events in the evolution in the sense that it created something totally new rather than only making more complex something which already existed. Gould also emphasizes the failure to identify any great change in the environment as a fundamental puzzle of Cambrian explosion. Cambrian explosion is also regarded in many quantum theories of consciousness (including TGD) as a revolution in the evolution of consciousness: for instance, micro-tubuli emerged at this time. The periods of expansion might be necessary for the emergence of multicellular life forms on planets and the fact that they unavoidably occur sooner or later suggests that also life develops unavoidably.

2. TGD predicts a decrease of the surface gravity by a factor 1/4 during this period. The reduction of the surface gravity would have naturally led to the emergence of dinosaurs 230 million years ago as a response coming 45 million years after the accelerated expansion ceased. Other reasons led then to the decline and eventual catastrophic disappearance of the dinosaurs. The reduction of gravity might have had some gradually increasing effects on the shape of organisms also at microscopic level and manifest itself in the evolution of genome during expansion period.
3. A possibly testable prediction following from angular momentum conservation ( $\omega R^2 = constant$ ) is that the duration of day has increased gradually and was four times shorter during the Cambrian era. For instance, genetically coded bio-clocks of simple organisms during the expansion period could have followed the increase of the length of day with certain lag or failed to follow it completely. The simplest known circadian clock is that of the prokaryotic cyanobacteria. Recent research has demonstrated that the circadian clock of *Synechococcus elongatus* can be reconstituted in vitro with just the three proteins of their central oscillator. This clock has been shown to sustain a 22 hour rhythm over several days upon the addition of ATP: the rhythm is indeed faster than the circadian rhythm. For humans the average innate circadian rhythm is however 24 hours 11 minutes and thus conforms with the fact that human genome has evolved much later than the expansion ceased.
4. Scientists have found a fossil of a sea scorpion with size of 2.5 meters [I2], which has lived for about 10 million years for 400 million years ago in Germany. The gigantic size would conform nicely with the much smaller value of surface gravity at that time. The finding would conform nicely with the much smaller value of surface gravity at that time. Also the emergence of trees could be understood in terms of a gradual growth of the maximum plant size as the surface gravity was reduced. The fact that the oldest known tree fossil is 385 million years old [I5] conforms with this picture.

#### 6.4.5 Did intra-terrestrial life burst to the surface of Earth during Cambrian expansion?

The possibility of intra-terrestrial life [K10] is one of the craziest TGD inspired ideas about the evolution of life and it is quite possible that in its strongest form the hypothesis is unrealistic. One can however try to find what one obtains from the combination of the IT hypothesis with the idea of pre-Cambrian granite Earth. Could the harsh pre-Cambrian conditions have allowed only intra-terrestrial multicellular life? Could the Cambrian explosion correspond to the moment of birth for this life in the very concrete sense that the magma flow brought it into the day-light?

1. Gould emphasizes the mysterious fact that very many life forms of Cambrian explosion looked like final products of a long evolutionary process. Could the eruption of magma from the Earth interior have induced a burst of intra-terrestrial life forms to the Earth's surface? This might make sense: the life forms living at the bottom of sea do not need direct solar light so that they could have had intra-terrestrial origin. It is quite possible that Earth's mantle contained low temperature water pockets, where the complex life forms might have evolved in an environment shielded from meteoric bombardment and UV radiation.
2. Sea water is salty. It is often claimed that the average salt concentration inside cell is that of the primordial sea: I do not know whether this claim can be really justified. If the claim is true, the cellular salt concentration should reflect the salt concentration of the water inside the pockets. The water inside water pockets could have been salty due to the diffusion of the salt from ground but need not have been same as that for the ocean water (higher than for cell interior and for obvious reasons). Indeed, the water in the underground reservoirs in arid regions such as Sahara is salty, which is the reason for why agriculture is absent in these regions. Note also that the cells of marine invertebrates are osmoconformers able to cope with the changing salinity of the environment so that the Cambrian revolutionaries could have survived the change in the salt concentration of environment.
3. What applies to Earth should apply also to other similar planets and Mars [E3] is very similar to Earth. The radius is .533 times that for Earth so that after quantum leap doubling the

radius and thus Schumann frequency scale (7.8 Hz would be the lowest Schumann frequency) would be essentially same as for Earth now. Mass is .131 times that for Earth so that surface gravity would be .532 of that for Earth now and would be reduced to .131 meaning quite big dinosaurs! have learned that Mars probably contains large water reservoirs in it's interior and that there is an un-identified source of methane gas usually assigned with the presence of life. Could it be that Mother Mars is pregnant and just waiting for the great quantum leap when it starts to expand and gives rise to a birth of multicellular life forms. Or expressing freely how Bible describes the moment of birth: in the beginning there was only darkness and water and then God said Let the light come!

To sum up, TGD would provide only the long sought mechanism of expansion and a possible connection with the biological evolution. It would be indeed fascinating if Planck constant changing quantum phase transitions in planetary scale would have profoundly affected the biosphere.

## 6.5 Allais Effect As Evidence For Large Values Of Gravitational Planck Constant?

Allais effect [E1, E8] is a fascinating gravitational anomaly associated with solar eclipses. It was discovered originally by M. Allais, a Nobelist in the field of economy, and has been reproduced in several experiments but not as a rule. The experimental arrangement uses so called paraconical pendulum, which differs from the Foucault pendulum in that the oscillation plane of the pendulum can rotate in certain limits so that the motion occurs effectively at the surface of sphere.

### 6.5.1 Experimental findings

Consider first a brief summary of the findings of Allais and others [E8].

a) In the ideal situation (that is in the absence of any other forces than gravitation of Earth) paraconical pendulum should behave like a Foucault pendulum. The oscillation plane of the paraconical pendulum however begins to rotate.

b) Allais concludes from his experimental studies that the orbital plane approach always asymptotically to a limiting plane and the effect is only particularly spectacular during the eclipse. During solar eclipse the limiting plane contains the line connecting Earth, Moon, and Sun. Allais explains this in terms of what he calls the anisotropy of space.

c) Some experiments carried out during eclipse have reproduced the findings of Allais, some experiments not. In the experiment carried out by Jeverdan and collaborators in Romania it was found that the period of oscillation of the pendulum decreases by  $\Delta f/f \simeq 5 \times 10^{-4}$  [E1, E7] which happens to correspond to the constant  $v_0 = 2^{-11}$  appearing in the formula of the gravitational Planck constant. It must be however emphasized that the overall magnitude of  $\Delta f/f$  varies by five orders of magnitude. Even the sign of  $\Delta f/f$  varies from experiment to experiment.

d) There is also quite recent finding by Popescu and Olenici, which they interpret as a quantization of the plane of oscillation of paraconical oscillator during solar eclipse [E9].

### 6.5.2 TGD based models for Allais effect

I have already earlier proposed an explanation of the effect in terms of classical  $Z^0$  force [K4]. If the  $Z^0$  charge to mass ratio of pendulum varies and if Earth and Moon are  $Z^0$  conductors, the resulting model is quite flexible and one might hope it could explain the high variation of the experimental results.

The rapid variation of the effect during the eclipse is however a problem for this approach and suggests that gravitational screening or some more general interference effect might be present. Gravitational screening alone cannot however explain Allais effect.

A model based on the idea that gravitational interaction is mediated by topological light rays (MEs) and that gravitons correspond to a gigantic value of the gravitational Planck constant however explains the Allais effect as an interference effect made possible by macroscopic quantum coherence in astrophysical length scales. Equivalence Principle fixes the model to a high degree and one ends up with an explicit formula for the anomalous gravitational acceleration and the general order of magnitude and the large variation of the frequency change as being due to the variation of

the distance ratio  $r_{S,P}/r_{M,P}$  ( $S$ ,  $M$ , and  $P$  refer to Sun, Moon, and pendulum respectively). One can say that the pendulum acts as an interferometer.

## 6.6 Applications To Elementary Particle Physics, Nuclear Physics, And Condensed Matter Physics

The hierarchy of Planck constants could have profound implications for even elementary particle physics since the strong constraints on the existence of new light particles coming from the decay widths of intermediate gauge bosons can be circumvented because direct decays to dark matter are not possible. On the other hand, if light scaled versions of elementary particles exist they must be dark since otherwise their existence would be visible in these decay widths. The constraints on the existence of dark nuclei and dark condensed matter are much milder. Cold fusion and some other anomalies of nuclear and condensed matter physics - in particular the anomalies of water-might have elegant explanation in terms of dark nuclei.

### 6.6.1 Leptohadron hypothesis

TGD suggests strongly the existence of lepto-hadron [K21]. Lepto-hadrons are bound states of color excited leptons and the anomalous production of  $e^+e^-$  pairs in heavy ion collisions finds a nice explanation as resulting from the decays of lepto-hadrons with basic condensate level  $k = 127$  and having typical mass scale of one  $MeV$ . The recent indications on the existence of a new fermion with quantum numbers of muon neutrino and the anomaly observed in the decay of orto-positronium give further support for the lepto-hadron hypothesis. There is also evidence for anomalous production of low energy photons and  $e^+e^-$  pairs in hadronic collisions.

The identification of lepto-hadrons as a particular instance in the predicted hierarchy of dark matters interacting directly only via graviton exchange allows to circumvent the lethal counter arguments against the lepto-hadron hypothesis ( $Z^0$  decay width and production of colored lepton jets in  $e^+e^-$  annihilation) even without assumption about the loss of asymptotic freedom.

PCAC hypothesis and its sigma model realization lead to a model containing only the coupling of the lepto-pion to the axial vector current as a free parameter. The prediction for  $e^+e^-$  production cross section is of correct order of magnitude only provided one assumes that lepto-pions (or electropions) decay to lepto-nucleon pair  $e_{ex}^+e_{ex}^-$  first and that lepto-nucleons, having quantum numbers of electron and having mass only slightly larger than electron mass, decay to lepton and photon. The peculiar production characteristics are correctly predicted. There is some evidence that the resonances decay to a final state containing  $n > 2$  particle and the experimental demonstration that lepto-nucleon pairs are indeed in question, would be a breakthrough for TGD.

During 18 years after the first published version of the model also evidence for colored  $\mu$  has emerged [C4]. Towards the end of 2008 CDF anomaly [C2] gave a strong support for the colored excitation of  $\tau$ . The lifetime of the light long lived state identified as a charged  $\tau$ -pion comes out correctly and the identification of the reported 3 new particles as p-adically scaled up variants of neutral  $\tau$ -pion predicts their masses correctly. The observed muon jets can be understood in terms of the special reaction kinematics for the decays of neutral  $\tau$ -pion to 3  $\tau$ -pions with mass scale smaller by a factor 1/2 and therefore almost at rest. A spectrum of new particles is predicted. The discussion of CDF anomaly [K21] led to a modification and generalization of the original model for lepto-pion production and the predicted production cross section is consistent with the experimental estimate.

### 6.6.2 Cold fusion, plasma electrolysis, and burning salt water

The article of Kanarev and Mizuno [D10] reports findings supporting the occurrence of cold fusion in NaOH and KOH hydrolysis. The situation is different from standard cold fusion where heavy water  $D_2O$  is used instead of  $H_2O$ .

In nuclear string model nucleon are connected by color bonds representing the color magnetic body of nucleus and having length considerably longer than nuclear size. One can consider also dark nuclei for which the scale of nucleus is of atomic size [L1], [L1]. In this framework can understand the cold fusion reactions reported by Mizuno as nuclear reactions in which part of what I call dark proton string having negatively charged color bonds (essentially a zoomed up variant of ordinary

nucleus with large Planck constant) suffers a phase transition to ordinary matter and experiences ordinary strong interactions with the nuclei at the cathode. In the simplest model the final state would contain only ordinary nuclear matter. The generation of plasma in plasma electrolysis can be seen as a process analogous to the positive feedback loop in ordinary nuclear reactions.

Rather encouragingly, the model allows to understand also deuterium cold fusion and leads to a solution of several other anomalies.

1. The so called lithium problem of cosmology (the observed abundance of lithium is by a factor 2.5 lower than predicted by standard cosmology [E5] ) can be resolved if lithium nuclei transform partially to dark lithium nuclei.
2. The so called  $H_{1.5}O$  anomaly of water [D11, D8, D13, D6] can be understood if 1/4 of protons of water forms dark lithium nuclei or heavier dark nuclei formed as sequences of these just as ordinary nuclei are constructed as sequences of  ${}^4He$  and lighter nuclei in nuclear string model. The results force to consider the possibility that nuclear isotopes unstable as ordinary matter can be stable dark matter.
3. The mysterious behavior burning salt water [D1] can be also understood in the same framework.
4. The model explains the nuclear transmutations observed in Kanarev's plasma electrolysis. This kind of transmutations have been reported also in living matter long time ago [C1, C6]. Intriguingly, several biologically important ions belong to the reaction products in the case of NaOH electrolysis. This raises the question whether cold nuclear reactions occur in living matter and are responsible for generation of biologically most important ions.

## 6.7 Applications To Biology And Neuroscience

The notion of field or magnetic body regarded as carrier of dark matter with large Planck constant and quantum controller of ordinary matter is the basic idea in the TGD inspired model of living matter.

### 6.7.1 Do molecular symmetries in living matter relate to non-standard values of Planck constant?

Water is exceptional element and the possibility that  $G_a$  as symmetry of singular factor space of CD in water and living matter is intriguing.

1. There is evidence for an icosahedral clustering in [D15] [D12]. Synaptic contacts contain clathrin molecules which are truncated icosahedrons and form lattice structures and are speculated to be involved with quantum computation like activities possibly performed by microtubules. Many viruses have the shape of icosahedron. One can ask whether these structures could be formed around templates formed by dark matter corresponding to 120-fold covering of  $CP_2$  points by CD points and having  $\hbar(CP_2) = 5\hbar_0$  perhaps corresponding color confined light dark quarks. Of course, a similar covering of CD points by  $CP_2$  could be involved.
2. It should be noticed that single nucleotide in DNA double strands corresponds to a twist of  $2\pi/10$  per single DNA triplet so that 10 DNA strands corresponding to length  $L(151) = 10$  nm (cell membrane thickness) correspond to  $3 \times 2\pi$  twist. This could be perhaps interpreted as evidence for group  $C_{10}$  perhaps making possible quantum computation at the level of DNA.
3. What makes realization of  $G_a$  as a symmetry of singular factor space of CD is that the biomolecules most relevant for the functioning of brain (DNA nucleotides, amino-acids acting as neurotransmitters, molecules having hallucinogenic effects) contain aromatic 5- and 6-cycles.

These observations led to an identification of the formula for Planck constant (two alternatives were allowed by the condition that Planck constant is algebraic homomorphism) which was not consistent with the model for dark gravitons. If one accepts the proposed formula of Planck constant, the dark space-time sheets with large Planck constant correspond to factor spaces of both  $\hat{CD}\backslash M^2$  and of  $CP_2\backslash S_T^2$ . This identification is of course possible and it remains to be seen whether it leads to any problems. For gravitational space-time sheets only coverings of both CD and  $CP_2$  make sense and the covering group  $G_a$  has very large order and does not correspond to geometric symmetries analogous to those of molecules.

### 6.7.2 High $T_c$ super-conductivity in living matter

The model for high  $T_c$  super-conductivity realized as quantum critical phenomenon predicts the basic scales of cell membrane [K5] from energy minimization and p-adic length scale hypothesis. This leads to the vision that cell membrane and possibly also its scaled up dark fractal variants define Josephson junctions generating Josephson radiation communicating information about the nearby environment to the magnetic body.

Any model of high  $T_c$  superconductivity should explain various strange features of high  $T_c$  superconductors. One should understand the high value of  $T_c$ , the ambivalent character of high  $T_c$  superconductors suggesting both BCS type Cooper pairs and exotic Cooper pairs with non-vanishing spin, the existence of pseudogap temperature  $T_{c_1} > T_c$  and scaling law for resistance for  $T_c \leq T < T_{c_1}$ , the role of fluctuating charged stripes which are anti-ferromagnetic defects of a Mott insulator, the existence of a critical doping, etc... [D7, D4].

There are reasons to believe that high  $T_c$  super-conductors correspond to quantum criticality in which at least two (cusp catastrophe as in van der Waals model), or possibly three or even more phases, are competing. A possible analogy is provided by the triple critical point for water vapor, liquid phase and ice coexist. Instead of long range thermal fluctuations long range quantum fluctuations manifesting themselves as fluctuating stripes are present [D7].

The TGD based model for high  $T_c$  super-conductivity [K5] relies on the notions of quantum criticality, general ideas of catastrophe theory, dynamical Planck constant, and many-sheeted space-time. The 4-dimensional spin glass character of space-time dynamics deriving from the vacuum degeneracy of the Kähler action defining the basic variational principle would realize space-time correlates for quantum fluctuations.

1. Two kinds of super-conductivities and ordinary non-super-conducting phase would be competing at quantum criticality at  $T_c$  and above it only one super-conducting phase and ordinary conducting phase located at stripes representing ferromagnetic defects making possible formation of  $S = 1$  Cooper pairs.
2. The first super-conductivity would be based on exotic Cooper pairs of large  $\hbar$  dark electrons with  $\hbar = 2^{11}\hbar_0$  and able to have spin  $S = 1$ , angular momentum  $L = 2$ , and total angular momentum  $J = 2$ . Second type of super-conductivity would be based on BCS type Cooper pairs having vanishing spin and bound by phonon interaction. Also they have large  $\hbar$  so that gap energy and critical temperature are scaled up in the same proportion. The exotic Cooper pairs are possible below the pseudo gap temperature  $T_{c_1} > T_c$  but are unstable against decay to BCS type Cooper pairs which above  $T_c$  are unstable against a further decay to conduction electrons flowing along stripes. This would reduce the exotic super-conductivity to finite conductivity obeying the observed scaling law for resistance.
3. The mere assumption that electrons of exotic Cooper pairs feed their electric flux to larger space-time sheet via *two* elementary particle sized wormhole contacts rather than only *one* wormhole contacts implies that the throats of wormhole contacts defining analogs of Higgs field must carry quantum numbers of quark and anti-quark. This inspires the idea that cylindrical space-time sheets, the radius of which turns out to be about about 5 nm, representing zoomed up dark electrons of Cooper pair with Planck constant  $\hbar = 2^{11}\hbar_0$  are colored and bound by a scaled up variant of color force to form a color confined state. Formation of Cooper pairs would have nothing to do with direct interactions between electrons. Thus high  $T_c$  super-conductivity could be seen as a first indication for the presence of scaled up variant of QCD in mesoscopic length scales.

This picture leads to a concrete model for high  $T_c$  superconductors as quantum critical superconductors [K5]. p-Adic length scale hypothesis stating that preferred p-adic primes  $p \simeq 2^k$ ,  $k$  integer, with primes (in particular Mersenne primes) preferred, makes the model quantitative.

1. An unexpected prediction is that coherence length  $\xi$  is actually  $\hbar_{eff}/\hbar_0 = 2^{11}$  times longer than the coherence length 5-10 Angstroms deduced theoretically from gap energy using conventional theory and varies in the range 1 – 5  $\mu\text{m}$ , the cell nucleus length scale. Hence type I super-conductor would be in question with stripes as defects of anti-ferromagnetic Mott insulator serving as duals for the magnetic defects of type I super-conductor in nearly critical magnetic field.
2. At quantitative level the model reproduces correctly the four poorly understood photon absorption lines and allows to understand the critical doping ratio from basic principles.
3. The current carrying structures have structure locally similar to that of axon including the double layered structure of cell membrane and also the size scales are predicted to be same. One of the characteristic absorption lines has energy of 0.05 eV which corresponds to the Josephson energy for neuronal membrane for activation potential  $V = 50$  mV. Hence the idea that axons are high  $T_c$  superconductors is highly suggestive. Dark matter hierarchy coming in powers  $\hbar/\hbar_0 = 2^{k^{11}}$  suggests hierarchy of Josephson junctions needed in TGD based model of EEG [K8].

### 6.7.3 Magnetic body as a sensory perceiver and intentional agent

The hypothesis that dark magnetic body serves as an intentional agent using biological body as a motor instrument and sensory receptor is consistent with Libet's findings about strange time delays of consciousness. Magnetic body would carry cyclotron Bose-Einstein condensates of various ions. Magnetic body must be able to perform motor control and receive sensory input from biological body.

Cell membrane would be a natural sensor providing information about cell interior and exterior to the magnetic body and dark photons at appropriate frequency range would naturally communicate this information. The strange quantitative co-incidences with the physics of cell membrane and high  $T_c$  super-conductivity support the idea that Josephson radiation generated by Josephson currents of dark electrons through cell membrane is responsible for this communication [K8].

Also fractally scaled up versions of cell membrane at higher levels of dark matter hierarchy (in particular those corresponding to powers  $n = 2^{k^{11}}$ ) are possible and the model for EEG indeed relies on this hypothesis. The thickness for the fractal counterpart of cell membrane thickness would be  $2^{44}$  fold and of order of depth of ionosphere! Although this looks weird it is completely consistent with the notion of magnetic body as an intentional agent.

Motor control would be most naturally performed via genome: this is achieved if flux sheets traverse through DNA strands. Flux quantization for large values of Planck constant requires rather large widths for the flux sheets. If flux sheet contains sequences of genomes like the page of book contains lines of text, a coherent gene expression becomes possible at level of organs and even populations and one can speak about super- and hyper-genomes. Introns might relate to the collective gene expression possibly realized electromagnetically rather than only chemically [K5, K6].

Dark cyclotron radiation with photon energy above thermal energy could be used for coordination purposes at least. The predicted hierarchy of copies of standard model physics leads to ask whether also dark copies of electro-weak gauge bosons and gluons could be important in living matter. As already mentioned, dark  $W$  bosons could make possible charge entanglement and non-local quantum bio-control by inducing voltage differences and thus ionic currents in living matter.

The identification of plasmoids as rotating magnetic flux structures carrying dark ions and electrons as primitive life forms is natural in this framework. There exists experimental support for this identification [I6] but the main objection is the high temperature involved: this objection could be circumvented if large  $\hbar$  phase is involved. A model for the pre-biotic evolution relying also on this idea is discussed in [K10].



At the level of biology there are now several concrete applications leading to a rich spectrum of predictions. Magnetic flux quanta would carry charged particles with large Planck constant.

1. The shortening of the flux tubes connecting biomolecules in a phase transition reducing Planck constant could be a basic mechanism of bio-catalysis and explain the mysterious ability of biomolecules to find each other. Similar process in time direction could explain basic aspects of symbolic memories as scaled down representations of actual events.
2. The strange behavior of cell membrane suggests that a dominating portion of important biological ions are actually dark ions at magnetic flux tubes so that ionic pumps and channels are needed only for visible ions. This leads to a model of nerve pulse explaining its unexpected thermodynamical properties with basic properties of Josephson currents making it unnecessary to use pumps to bring ions back after the pulse. The model predicts automatically EEG as Josephson radiation and explains the synchrony of both kHz radiation and of EEG.
3. The DC currents of Becker could be accompanied by Josephson currents running along flux tubes making possible dissipation free energy transfer and quantum control over long distances and meridians of chinese medicine could correspond to these flux tubes.
4. The model of DNA as topological quantum computer assumes that nucleotides and lipids are connected by ordinary or “wormhole” magnetic flux tubes acting as strands of braid and carrying dark matter with large Planck constant. The model leads to a new vision about TGD in which the assignment of nucleotides to quarks allows to understand basic regularities of DNA not understood from biochemistry.
5. Each physical system corresponds to an onion-like hierarchy of field bodies characterized by p-adic primes and value of Planck constant. The highest value of Planck constant in this hierarchy provides kind of intelligence quotient characterizing the evolutionary level of the system since the time scale of planned action and memory correspond to the temporal distance between tips of corresponding causal diamond (CD). Also the spatial size of the system correlates with the Planck constant. This suggests that great evolutionary leaps correspond to the increase of Planck constant for the highest level of hierarchy of personal magnetic bodies. For instance, neurons would have much more evolved magnetic bodies than ordinary cells.
6. At the level of DNA this vision leads to an idea about hierarchy of genomes. Magnetic flux sheets traversing DNA strands provide a natural mechanism for magnetic body to control the behavior of biological body by controlling gene expression. The quantization of magnetic flux states that magnetic flux is proportional to  $\hbar$  and thus means that the larger the value of  $\hbar$  is the larger the width of the flux sheet is. For larger values of  $\hbar$  single genome is not enough to satisfy this condition. This leads to the idea that the genomes of organs, organism, and even population, can organize like lines of text at the magnetic flux sheets and form in this manner a hierarchy of genomes responsible for a coherent gene expression at level of cell, organ, organism and population and perhaps even entire biosphere. This would also provide a mechanism by which collective consciousness would use its biological body - biosphere.

#### 6.7.4 DNA as topological quantum computer

I ended up with the recent model of TQC in bottom-up manner and this representation is followed also in the text. The model which looks the most plausible one relies on two specific ideas.

1. Sharing of labor means conjugate DNA would do TQC and DNA would “print” the outcome of TQC in terms of mRNA yielding amino-acids in the case of exons. RNA could result also in the case of introns but not always. The experience about computers and the general vision provided by TGD suggests that introns could express the outcome of TQC also electromagnetically in terms of standardized field patterns as Gariaev’s findings suggest [I3]. Also speech would be a form of gene expression. The quantum states braid (in zero energy ontology) would entangle with characteristic gene expressions. This argument turned out

to be based on a slightly wrong belief about DNA: later I learned that both strand and its conjugate are transcribed but in different directions. The symmetry breaking in the case of transcription is only local which is also visible in DNA replication as symmetry breaking between leading and lagging strand. Thus the idea about *entire* leading strand devoted to printing and second strand to TQC must be weakened appropriately.

2. The manipulation of braid strands transversal to DNA must take place at 2-D surface. Here dancing metaphor for topological quantum computation [C3] generalizes. The ends of the space-like braid are like dancers whose feet are connected by thin threads to a wall so that the dancing pattern entangles the threads. Dancing pattern defines both the time-like braid, the running of classical TQC program and its representation as a dynamical pattern. The space-like braid defined by the entangled threads represents memory storage so that TQC program is automatically written to memory as the braiding of the threads during the TQC. The inner membrane of the nuclear envelope and cell membrane with entire endoplasmic reticulum included are good candidates for dancing halls. The 2-surfaces containing the ends of the hydrophobic ends of lipids could be the parquets and lipids the dancers. This picture seems to make sense.

One ends up to the model also in top-down manner.

1. Darwinian selection for which standard theory of self-organization [B1] provides a model, should apply also to TQC programs. TQC programs should correspond to asymptotic self-organization patterns selected by dissipation in the presence of metabolic energy feed. The spatial and temporal pattern of the metabolic energy feed characterizes the TQC program - or equivalently - sub-program call.
2. Since braiding characterizes the TQC program, the self-organization pattern should correspond to a hydrodynamical flow or a pattern of magnetic field inducing the braiding. Braid strands must correspond to magnetic flux tubes of the magnetic body of DNA. If each nucleotide is transversal magnetic dipole it gives rise to transversal flux tubes, which can also connect to the genome of another cell.
3. The output of TQC sub-program is probability distribution for the outcomes of state function reduction so that the sub-program must be repeated very many times. It is represented as four-dimensional patterns for various rates (chemical rates, nerve pulse patterns, EEG power distributions, ...) having also identification as temporal densities of zero energy states in various scales. By the fractality of TGD Universe there is a hierarchy of TQC's corresponding to p-adic and dark matter hierarchies. Programs (space-time sheets defining coherence regions) call programs in shorter scale. If the self-organizing system has a periodic behavior each TQC module defines a large number of almost copies of itself asymptotically. Generalized EEG could naturally define this periodic pattern and each period of EEG would correspond to an initiation and halting of TQC. This brings in mind the periodically occurring sol-gel phase transition inside cell near the cell membrane.
4. Fluid flow must induce the braiding which requires that the ends of braid strands must be anchored to the fluid flow. Recalling that lipid mono-layers of the cell membrane are liquid crystals and lipids of interior mono-layer have hydrophilic ends pointing towards cell interior, it is easy to guess that DNA nucleotides are connected to lipids by magnetic flux tubes and hydrophilic lipid ends are stuck to the flow.
5. The topology of the braid traversing cell membrane cannot be affected by the hydrodynamical flow. Hence braid strands must be split during TQC. This also induces the desired magnetic isolation from the environment. Halting of TQC reconnects them and makes possible the communication of the outcome of TQC.
6. There are several problems related to the details of the realization. How nucleotides A, T, C, G are coded to strand color and what this color corresponds to? The prediction that wormhole contacts carrying quark and anti-quark at their ends appear in all length scales in TGD Universe resolves the problem. How to split the braid strands in a controlled manner?

High  $T_c$  super conductivity provides a partial understanding of the situation: braid strand can be split only if the supra current flowing through it vanishes. From the proportionality of Josephson current to the quantity  $\sin(\int 2eV dt)$  it follows that a suitable voltage pulse  $V$  induces DC supra-current and its negative cancels it. The conformation of the lipid controls whether it can follow the flow or not. How magnetic flux tubes can be cut without breaking the conservation of the magnetic flux? The notion of wormhole magnetic field saves the situation now: after the splitting the flux returns back along the second space-time sheet of wormhole magnetic field.

To sum up, it seems that essentially all new physics involved with TGD based view about quantum biology enter to the model in crucial manner.

### 6.7.5 Quantum model of nerve pulse and EEG

In this article a unified model of nerve pulse and EEG is discussed.

1. In TGD Universe the function of EEG and its variants is to make possible communications from the cell membrane to the magnetic body and the control of the biological body by the magnetic body via magnetic flux sheets traversing DNA by inducing gene expression. This leads to the notions of super- and hyper-genome predicting coherent gene expression at level of organs and population.
2. The assignment the predicted ranged classical weak and color gauge fields to dark matter hierarchy was a crucial step in the evolution of the model, and led among other things to a model of high  $T_c$  superconductivity predicting the basic scales of cell, and also to a generalization of EXG to a hierarchy of ZXGs, WXGs, and GXGs corresponding to  $Z^0$ ,  $W$  bosons and gluons.
3. Dark matter hierarchy and the associated hierarchy of Planck constants plays a key role in the model. For instance, in the case of EEG Planck constant must be so large that the energies of dark EEG photons are above thermal energy at physiological temperatures. The assumption that a considerable fraction of the ionic currents through the cell membrane are dark currents flowing along the magnetic flux tubes explains the strange findings about ionic currents through cell membrane. Concerning the model of nerve pulse generation, the newest input comes from the model of DNA as a topological quantum computer and experimental findings challenging Hodgkin-Huxley model as even approximate description of the situation.
4. The identification of the cell interior as gel phase containing most of water as structured water around cytoskeleton - rather than water containing bio-molecules as solutes as assumed in Hodgkin-Huxley model - allows to understand many of the anomalous behaviors associated with the cell membrane and also the different densities of ions in the interior and exterior of cell at qualitative level. The proposal of Pollack that basic biological functions involve phase transitions of gel phase generalizes in TGD framework to a proposal that these phase transitions are induced by quantum phase transitions changing the value of Planck constant. In particular, gel-sol phase transition for the peripheral cytoskeleton induced by the primary wave would accompany nerve pulse propagation. This view about nerve pulse is not consistent with Hodgkin-Huxley model.

The model leads to the following picture about nerve pulse and EEG.

1. The system would consist of two superconductors- microtubule space-time sheet and the space-time sheet in cell exterior- connected by Josephson junctions represented by magnetic flux tubes defining also braiding in the model of TQC. The phase difference between two super-conductors would obey Sine-Gordon equation allowing both standing and propagating solitonic solutions. A sequence of rotating gravitational penduli coupled to each other would be the mechanical analog for the system. Soliton sequences having as a mechanical analog penduli rotating with constant velocity but with a constant phase difference between them would generate moving kHz synchronous oscillation. Periodic boundary conditions at the ends of the axon rather than chemistry determine the propagation velocities of kHz waves

and kHz synchrony is an automatic consequence since the times taken by the pulses to travel along the axon are multiples of same time unit. Also moving oscillations in EEG range can be considered and would require larger value of Planck constant in accordance with vision about evolution as gradual increase of Planck constant.

2. During nerve pulse one pendulum would be kicked so that it would start to oscillate instead of rotating and this oscillation pattern would move with the velocity of kHz soliton sequence. The velocity of kHz wave and nerve pulse is fixed by periodic boundary conditions at the ends of the axon implying that the time spent by the nerve pulse in traveling along axon is always a multiple of the same unit: this implies kHz synchrony. The model predicts the value of Planck constant for the magnetic flux tubes associated with Josephson junctions and the predicted force caused by the ionic Josephson currents is of correct order of magnitude for reasonable values of the densities of ions. The model predicts kHz em radiation as Josephson radiation generated by moving soliton sequences. EEG would also correspond to Josephson radiation: it could be generated either by moving or standing soliton sequences (latter are naturally assignable to neuronal cell bodies for which  $\hbar$  should be correspondingly larger): synchrony is predicted also now.

## 7 Appendix

### 7.1 About Inclusions Of Hyper-Finite Factors Of Type $II_1$

Many names have been assigned to inclusions: Jones, Wenzl, Ocneacnu, Pimsner-Popa, Wasserman [A3]. It would seem to me that the notion Jones inclusion includes them all so that various names would correspond to different concrete realizations of the inclusions conjugate under outer automorphisms.

1. According to [A3] for inclusions with  $\mathcal{M} : \mathcal{N} \leq 4$  (with  $A_1^{(1)}$  excluded) there exists a countable infinity of sub-factors which are pairwise non inner conjugate but conjugate to  $\mathcal{N}$ .
2. Also for any finite group  $G$  and its outer action there exists uncountably many sub-factors which are pairwise non inner conjugate but conjugate to the fixed point algebra of  $G$  [A3]. For any amenable group  $G$  the inclusion is also unique apart from outer automorphism [A1].

Thus it seems that not only Jones inclusions but also more general inclusions are unique apart from outer automorphism.

Any \*-endomorphism  $\sigma$ , which is unit preserving, faithful, and weakly continuous, defines a sub-factor of type  $II_1$  factor [A3]. The construction of Jones leads to a standard inclusion sequence  $\mathcal{N} \subset \mathcal{M} \subset \mathcal{M}^1 \subset \dots$ . This sequence means addition of projectors  $e_i$ ,  $i < 0$ , having visualization as an addition of braid strand in braid picture. This hierarchy exists for all factors of type  $II$ . At the limit  $\mathcal{M}^\infty = \cup_i \mathcal{M}^i$  the braid sequence extends from  $-\infty$  to  $\infty$ . Inclusion hierarchy can be understood as a hierarchy of Connes tensor powers  $\mathcal{M} \otimes_{\mathcal{N}} \mathcal{M} \dots \otimes_{\mathcal{N}} \mathcal{M}$ . Also the ordinary tensor powers of hyper-finite factors of type  $II_1$  (HFF) as well as their tensor products with finite-dimensional matrix algebras are isomorphic to the original HFF so that these objects share the magic of fractals.

Under certain assumptions the hierarchy can be continued also in opposite direction. For a finite index an infinite inclusion hierarchy of factors results with the same value of index.  $\sigma$  is said to be basic if it can be extended to \*-endomorphisms from  $\mathcal{M}^1$  to  $\mathcal{M}$ . This means that the hierarchy of inclusions can be continued in the opposite direction: this means elimination of strands in the braid picture. For finite factors (as opposed to hyper-finite ones) there are no basic \*-endomorphisms of  $\mathcal{M}$  having fixed point algebra of non-abelian  $G$  as a sub-factor [A3].

#### 1. Jones inclusions

For hyper-finite factors of type  $II_1$  Jones inclusions allow basic \*-endomorphism. They exist for all values of  $\mathcal{M} : \mathcal{N} = r$  with  $r \in \{4\cos^2(\pi/n) | n \geq 3\} \cap [4, \infty)$  [A3]. They are defined for an algebra defined by projectors  $e_i$ ,  $i \geq 1$ . All but nearest neighbor projectors commute.  $\lambda = 1/r$  appears in the relations for the generators of the algebra given by  $e_i e_j e_i = \lambda e_i$ ,  $|i-j|=1$ .  $\mathcal{N} \subset \mathcal{M}$  is identified as the double commutator of algebra generated by  $e_i$ ,  $i \geq 2$ .

This means that principal graph and its dual are equivalent and the braid defined by projectors can be continued not only to  $-\infty$  but that also the dropping of arbitrary number of strands is possible [A3]. It would seem that ADE property of the principal graph meaning single root length codes for the duality in the case of  $r \leq 4$  inclusions.

Irreducibility holds true for  $r < 4$  in the sense that the intersection of  $Q' \cap P = P' \cap P = C$ . For  $r \geq 4$  one has  $\dim(Q' \cap P) = 2$ . The operators commuting with  $Q$  contain besides identify operator of  $Q$  also the identify operator of  $P$ .  $Q$  would contain a single finite-dimensional matrix factor less than  $P$  in this case. Basic \*-endomorphisms with  $\sigma(P) = Q$  is  $\sigma(e_i) = e_{i+1}$ . The difference between genuine symmetries of quantum TGD and symmetries which can be mimicked by TGD could relate to the irreducibility for  $r < 4$  and raise these inclusions in a unique position. This difference could partially justify the hypothesis that only the groups  $G_a \times G_b \subset SU(2) \times SU(2) \subset SL(2, C) \times SU(3)$  define orbifold coverings of  $H_{\pm} = CD \times CP_2 \rightarrow H_{\pm}/G_a \times G_b$ .

### 2. Wasserman's inclusion

Wasserman's construction of  $r = 4$  factors clarifies the role of the subgroup of  $G \subset SU(2)$  for these inclusions. Also now  $r = 4$  inclusion is characterized by a discrete subgroup  $G \subset SU(2)$  and is given by  $(1 \otimes \mathcal{M})^G \subset (M_2(C) \times \mathcal{M})^G$ . According to [A3] Jones inclusions are irreducible also for  $r = 4$ . The definition of Wasserman inclusion for  $r = 4$  seems however to imply that the identity matrices of both  $\mathcal{M}^G$  and  $(M(2, C) \otimes \mathcal{M})^G$  commute with  $\mathcal{M}^G$  so that the inclusion should be reducible for  $r = 4$ .

Note that  $G$  leaves both the elements of  $\mathcal{N}$  and  $\mathcal{M}$  invariant whereas  $SU(2)$  leaves the elements of  $\mathcal{N}$  invariant.  $M(2, C)$  is effectively replaced with the orbifold  $M(2, C)/G$ , with  $G$  acting as automorphisms. The space of these orbits has complex dimension  $d = 4$  for finite  $G$ .

For  $r < 4$  inclusion is defined as  $M^G \subset M$ . The representation of  $G$  as outer automorphism must change step by step in the inclusion sequence  $\dots \subset \mathcal{N} \subset \mathcal{M} \subset \dots$  since otherwise  $G$  would act trivially as one proceeds in the inclusion sequence. This is true since each step brings in additional finite-dimensional tensor factor in which  $G$  acts as automorphisms so that although  $\mathcal{M}$  can be invariant under  $G_{\mathcal{M}}$  it is not invariant under  $G_{\mathcal{N}}$ .

These two inclusions might accompany each other in TGD based physics. One could consider  $r < 4$  inclusion  $\mathcal{N} = \mathcal{M}^G \subset \mathcal{M}$  with  $G$  acting non-trivially in  $\mathcal{M}/\mathcal{N}$  quantum Clifford algebra.  $\mathcal{N}$  would decompose by  $r = 4$  inclusion to  $\mathcal{N}_1 \subset \mathcal{N}$  with  $SU(2)$  taking the role of  $G$ .  $\mathcal{N}/\mathcal{N}_1$  quantum Clifford algebra would transform non-trivially under  $SU(2)$  but would be  $G$  singlet.

In TGD framework the  $G$ -invariance for  $SU(2)$  representations means a reduction of  $S^2$  to the orbifold  $S^2/G$ . The coverings  $H_{\pm} \rightarrow H_{\pm}/G_a \times G_b$  should relate to these double inclusions and  $SU(2)$  inclusion could mean Kac-Moody type gauge symmetry for  $\mathcal{N}$ . Note that the presence of the factor containing only unit matrix should relate directly to the generator  $d$  in the generator set of affine algebra in the McKay construction. The physical interpretation of the fact that almost all ADE type extended diagrams ( $D_n^{(1)}$  must have  $n \geq 4$ ) are allowed for  $r = 4$  inclusions whereas  $D_{2n+1}$  and  $E_6$  are not allowed for  $r < 4$ , remains open.

## 7.2 Generalization From $Su(2)$ To Arbitrary Compact Group

The inclusions with index  $\mathcal{M} : \mathcal{N} < 4$  have one-dimensional relative commutant  $\mathcal{N}' \cup \mathcal{M}$ . The most obvious conjecture that  $\mathcal{M} : \mathcal{N} \geq 4$  corresponds to a non-trivial relative commutant is wrong. The index for Jones inclusion is identifiable as the square of quantum dimension of the fundamental representation of  $SU(2)$ . This identification generalizes to an arbitrary representation of arbitrary compact Lie group.

In his thesis Wenzl [A2] studied the representations of Hecke algebras  $H_n(q)$  of type  $A_n$  obtained from the defining relations of symmetric group by the replacement  $e_i^2 = (q - 1)e_i + q$ .  $H_n$  is isomorphic to complex group algebra of  $S_n$  if  $q$  is not a root of unity and for  $q = 1$  the irreducible representations of  $H_n(q)$  reduce trivially to Young's representations of symmetric groups. For primitive roots of unity  $q = \exp(i2\pi/l)$ ,  $l = 4, 5, \dots$ , the representations of  $H_n(\infty)$  give rise to inclusions for which index corresponds to a quantum dimension of any irreducible representation of  $SU(k)$ ,  $k \geq 2$ . For  $SU(2)$  also the value  $l = 3$  is allowed for spin 1/2 representation.

The inclusions are obtained by dropping the first  $m$  generators  $e_k$  from  $H_{\infty}(q)$  and taking double commutant of both  $H_{\infty}$  and the resulting algebra. The relative commutant corresponds to  $H_m(q)$ . By reducing by the minimal projection to relative commutant one obtains an inclusion

with a trivial relative commutant. These inclusions are analogous to a discrete states superposed in continuum. Thus the results of Jones generalize from the fundamental representation of  $SU(2)$  to all representations of all groups  $SU(k)$ , and in fact to those of general compact groups as it turns out.

The generalization of the formula for index to square of quantum dimension of an irreducible representation of  $SU(k)$  reads as

$$\mathcal{M} : \mathcal{N} = \prod_{1 \leq r < s \leq k} \frac{\sin^2((\lambda_r - \lambda_s + s - r)\pi/l)}{\sin^2((s - r)n/l)}. \quad (7.1)$$

Here  $\lambda_r$  is the number of boxes in the  $r^{th}$  row of the Yang diagram with  $n$  boxes characterizing the representations and the condition  $1 \leq k \leq l - 1$  holds true. Only Young diagrams satisfying the condition  $l - k = \lambda_1 - \lambda_{r_{max}}$  are allowed.

The result would allow to restrict the generalization of the imbedding space in such a manner that only cyclic group  $Z_n$  appears in the covering of  $M^4 \rightarrow M^4/G_a$  or  $CP_2 \rightarrow CP_2/G_b$  factor. Be as it may, it seems that quantum representations of any compact Lie group can be realized using the generalization of the imbedding space. In the case of  $SU(2)$  the interpretation of higher-dimensional quantum representations in terms of Connes tensor products of 2-dimensional fundamental representations is highly suggestive.

The groups  $SO(3, 1) \times SU(3)$  and  $SL(2, C) \times U(2)_{ew}$  have a distinguished position both in physics and quantum TGD and the vision about physics as a generalized number theory implies them. Also the general pattern for inclusions selects these groups, and one can say that the condition that all possible statistics are realized is guaranteed by the choice  $M^4 \times CP_2$ .

1.  $n > 2$  for the quantum counterparts of the fundamental representation of  $SU(2)$  means that braid statistics for Jones inclusions cannot give the usual fermionic statistics. That Fermi statistics cannot “emerge” conforms with the role of infinite- $D$  Clifford algebra as a canonical representation of HFF of type  $II_1$ .  $SO(3, 1)$  as isometries of  $H$  gives  $Z_2$  statistics via the action on spinors of  $M^4$  and  $U(2)$  holonomies for  $CP_2$  realize  $Z_2$  statistics in  $CP_2$  degrees of freedom.
2.  $n > 3$  for more general inclusions in turn excludes  $Z_3$  statistics as braid statistics in the general case.  $SU(3)$  as isometries induces a non-trivial  $Z_3$  action on quark spinors but trivial action at the imbedding space level so that  $Z_3$  statistics would be in question.

## REFERENCES

### Mathematics

- [A1] Kawahigashi Y Evans D. *Quantum symmetries on operator algebras*. Oxford University Press, New York, 1998.
- [A2] Wenzl H. Hecke algebra of type  $A_n$  and subfactors. *Invent Math*, 92, 1988.
- [A3] Choda M. Conjugate but non inner conjugate subfactors. *Proc AMS*. Available at: <http://tinyurl.com/n1e4p7b>, 124(1), January 1996.

### Theoretical Physics

- [B1] Self organization. Available at: [http://en.wikipedia.org/wiki/Self\\_organization](http://en.wikipedia.org/wiki/Self_organization).

## Particle and Nuclear Physics

- [C1] Kervran CL. *Biological transmutations, and their applications in chemistry, physics, biology, ecology, medicine, nutrition, agriculture, geology*. Swan House Publishing Co., 1972.
- [C2] CDF Collaboration. Study of multi-muon events produced in p-pbar collisions at  $\sqrt{s}=1.96$  TeV. Available at: [http://arxiv.org/PS\\_cache/arxiv/pdf/0810/0810.0714v1.pdf](http://arxiv.org/PS_cache/arxiv/pdf/0810/0810.0714v1.pdf), 2008.
- [C3] Samuel E. Ghost in the Atom. *New Scientist*, (2366):30, October 2002.
- [C4] Valencia G He XG, Tandean J. Has HyperCP Observed a Light Higgs Boson? *Phys Rev D*. Available at: <http://arxiv.org/abs/hep-ph/0610274>, 74, 2007.
- [C5] McLerran L Ludham T. What Have We Learned From the Relativistic Heavy Ion Collider? *Phys Today*. Available at: <http://www.physicstoday.org/vol-56/iss-10/p48.html>, October 2003.
- [C6] Bird C Tompkins P. *The secret life of plants*. Harper & Row, New York, 1973.

## Condensed Matter Physics

- [D1] Burning salt water. Available at: <http://www.youtube.com/watch?v=aGgOATfoBgo>.
- [D2] Fractional quantum Hall Effect. Available at: [http://en.wikipedia.org/wiki/Fractional\\_quantum\\_Hall\\_effect](http://en.wikipedia.org/wiki/Fractional_quantum_Hall_effect).
- [D3] Monroe D. Know Your Anyons. *New Scientist*, (2676), 2008.
- [D4] Tranquada JM Emery VJ, Kivelson SA. Stripe phases in high-temperature superconductors. *Perspective*. Available at: <http://www.pnas.org/cgi/reprint/96/16/8814.pdf>, 96(16), August 1999.
- [D5] Miller JB et al. Fractional Quantum Hall effect in a quantum point contact at filling fraction  $5/2$ . Available at: <http://arxiv.org/abs/cond-mat/0703161v2>, 2007.
- [D6] Moreh R et al. Search for anomalous scattering of keV neutrons from H<sub>2</sub>O-D<sub>2</sub>O mixtures. *Phys Rev*, 94, 2005.
- [D7] Zaanen J. Superconductivity: Quantum Stripe Search. *Nature*. Available at: <http://www.lorentz.leidenuniv.nl/~jan/nature03/qustripes06.pdf>, April 2006.
- [D8] Borchardt JK. The chemical formula H<sub>2</sub>O - a misnomer. *Alchemist*, August 2003.
- [D9] Jain JK. *Phys Rev*, 63, 1989.
- [D10] Mizuno T Kanarev P. Cold fusion by plasma electrolysis of water. Available at: <http://www.guns.connect.fi/innoplaza/energy/story/Kanarev/codlfusion/>, 2002.
- [D11] Chaplin M. Water Structure and Behavior. Available at: <http://www.lsbu.ac.uk/water/index.html>, 2005.
- [D12] Chaplin M. Water as a Network of Icosahedral Water Clusters. Available at: <http://www.lsbu.ac.uk/water/clusters.html>, 2006.
- [D13] Cowley RA. Neutron-scattering experiments and quantum entanglement. *Phys B*, 350:243–245, 2004.
- [D14] Laughlin RB. *Phys Rev*, 50, 1990.
- [D15] Glashow SL. Can Sci Save the World? Available at: [http://www.hypothesis.it/nobel/nobel199/eng/pro/pro\\_2.htm](http://www.hypothesis.it/nobel/nobel199/eng/pro/pro_2.htm), 1999.

## Cosmology and Astro-Physics

- [E1] Allais effect. Available at: [http://en.wikipedia.org/wiki/Allais\\_effect](http://en.wikipedia.org/wiki/Allais_effect).
- [E2]  $\Lambda$ -CDM model. Available at: [http://en.wikipedia.org/wiki/Lambda-CDM\\_model](http://en.wikipedia.org/wiki/Lambda-CDM_model).
- [E3] Mars. Available at: <http://en.wikipedia.org/wiki/Mars>.
- [E4] Dark energy. *Physics World*. Available at: <http://physicsworld.com/cws/article/print/19419>, May 2004.
- [E5] Primas F Charbonnel C. The lithium content of the Galactic Halo stars. Available at: <http://arxiv.org/abs/astro-ph/0505247>, 2005.
- [E6] Nottale L Da Rocha D. Gravitational Structure Formation in Scale Relativity. Available at: <http://arxiv.org/abs/astro-ph/0310036>, 2003.
- [E7] Antonescu V Jeverdan GT, Rusu GI. Experiments using the Foucault pendulum during the solar eclipse of 15 February, 1961. *Bibl Astr*, 1(55), 1981.
- [E8] Allais M. Should the Laws of Gravitation Be Reconsidered: Part I,II,III? Available at: <http://home.t01.itscom.net/allais/blackprior/allais/lawgravit/lawgrav-one.pdf>, 1959.
- [E9] Olenici D Popescu VA. A confirmation of the Allais and Jeverdan-Rusu-Antonescu effects during the solar eclipse from 22 September 2006, and the quantization behavior of pendulum. Available at: <http://www.hessdalen.org/sse/program/Articol.pdf>, 2007.

## Physics of Earth

- [F1] A challenge to all geologists of Earth. Available at: <http://www.nealadams.com/challenge.html>.
- [F2] Expanding Earth Theory. Available at: [http://en.wikipedia.org/wiki/Expanding\\_earth\\_theory](http://en.wikipedia.org/wiki/Expanding_earth_theory).
- [F3] Oceanic trench. Available at: [http://en.wikipedia.org/wiki/Oceanic\\_trench](http://en.wikipedia.org/wiki/Oceanic_trench).
- [F4] Orogenies. Available at: <http://en.wikipedia.org/wiki/Orogenies>.
- [F5] Plate tectonics. Available at: [http://en.wikipedia.org/wiki/Plate\\_tectonics](http://en.wikipedia.org/wiki/Plate_tectonics).
- [F6] Volcano. Available at: <http://en.wikipedia.org/wiki/Volcano>.
- [F7] Adams N. Conspiracy of Sci , Earth is in fact growing. Available at: <http://www.youtube.com/watch?v=VjgidAICoQI>, 2006.

## Biology

- [I1] Dinosaurs. Available at: <http://en.wikipedia.org/wiki/Dinosaur>.
- [I2] Tetlie OE Braddy SJ, Poschmann M. Giant claw reveals the largest ever arthropod. *Biol Lett*. Available at: <http://www.journals.royalsoc.ac.uk/content/t15r2588mn27n0w1>, November 2007.
- [I3] Gariaev PP et al. The spectroscopy of bio-photons in non-local genetic regulation. *J Non-Locality and Remote Mental Interactions*. Available at: <http://www.emergentmind.org/gariaevI3.htm>, (3), 2002.
- [I4] Tovmash AV Gariaev PP, Tertishni GG. Experimental investigation in vitro of holographic mapping and holographic transposition of DNA in conjunction with the information pool encircling DNA. *New Medical Technologies*, 9:42–53, 2007.



- [15] O'Donoghue J. How trees changed the world? *New Scientist*. Available at: <http://tinyurl.com/our2ldk>, 2631, November 2007.
- [16] Sanduloviciu M Lozneau E. Minimal-cell system created in laboratory by self-organization. *Chaos , Solitons & Fractals*, 18(2):335, September 2003.
- [17] Gould SJ. *Wonderful Life*. Penguin Books, 1991.

## Books related to TGD

- [K1] Pitkänen M. Is it Possible to Understand Coupling Constant Evolution at Space-Time Level? In *Towards M-Matrix*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdquantum/tgdquantum.html#rgflow](http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#rgflow), 2006.
- [K2] Pitkänen M. About Nature of Time. In *TGD Inspired Theory of Consciousness*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdconsc/tgdconsc.html#timenature](http://tgdtheory.fi/public_html/tgdconsc/tgdconsc.html#timenature), 2006.
- [K3] Pitkänen M. About the New Physics Behind Qualia. In *Quantum Hardware of Living Matter*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/bioware/bioware.html#newphys](http://tgdtheory.fi/public_html/bioware/bioware.html#newphys), 2006.
- [K4] Pitkänen M. Anomalies Related to the Classical  $Z^0$  Force and Gravitation. In *TGD and Fringe Physics*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/freenergy/freenergy.html#Zanom](http://tgdtheory.fi/public_html/freenergy/freenergy.html#Zanom), 2006.
- [K5] Pitkänen M. Bio-Systems as Super-Conductors: part I. In *Quantum Hardware of Living Matter*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/bioware/bioware.html#superc1](http://tgdtheory.fi/public_html/bioware/bioware.html#superc1), 2006.
- [K6] Pitkänen M. Bio-Systems as Super-Conductors: part II. In *Quantum Hardware of Living Matter*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/bioware/bioware.html#superc2](http://tgdtheory.fi/public_html/bioware/bioware.html#superc2), 2006.
- [K7] Pitkänen M. Dark Forces and Living Matter. In *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/neuplanck/neuplanck.html#darkforces](http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#darkforces), 2006.
- [K8] Pitkänen M. Dark Matter Hierarchy and Hierarchy of EEGs. In *TGD and EEG*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdeeg/tgdeeg.html#eegdark](http://tgdtheory.fi/public_html/tgdeeg/tgdeeg.html#eegdark), 2006.
- [K9] Pitkänen M. Dark Nuclear Physics and Condensed Matter. In *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/neuplanck/neuplanck.html#exonuclear](http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#exonuclear), 2006.
- [K10] Pitkänen M. Evolution in Many-Sheeted Space-Time. In *Genes and Memes*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/genememe/genememe.html#prebio](http://tgdtheory.fi/public_html/genememe/genememe.html#prebio), 2006.
- [K11] Pitkänen M. Macroscopic Quantum Coherence and Quantum Metabolism as Different Sides of the Same Coin: Part I. In *Bio-Systems as Conscious Holograms*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/hologram/hologram.html#metab](http://tgdtheory.fi/public_html/hologram/hologram.html#metab), 2006.
- [K12] Pitkänen M. Negentropy Maximization Principle. In *TGD Inspired Theory of Consciousness*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdconsc/tgdconsc.html#nmpc](http://tgdtheory.fi/public_html/tgdconsc/tgdconsc.html#nmpc), 2006.
- [K13] Pitkänen M. New Particle Physics Predicted by TGD: Part I. In *p-Adic Physics*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/padphys/padphys.html#mass4](http://tgdtheory.fi/public_html/padphys/padphys.html#mass4), 2006.

- [K14] Pitkänen M. Nuclear String Hypothesis. In *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/neuplanck/neuplanck.html#nuclstring](http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#nuclstring), 2006.
- [K15] Pitkänen M. p-Adic Particle Massivation: Hadron Masses. In *p-Adic Length Scale Hypothesis and Dark Matter Hierarchy*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/padphys/padphys.html#mass3](http://tgdtheory.fi/public_html/padphys/padphys.html#mass3), 2006.
- [K16] Pitkänen M. p-Adic Physics as Physics of Cognition and Intention. In *TGD Inspired Theory of Consciousness*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdconsc/tgdconsc.html#cognic](http://tgdtheory.fi/public_html/tgdconsc/tgdconsc.html#cognic), 2006.
- [K17] Pitkänen M. Quantum Hall effect and Hierarchy of Planck Constants. In *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/neuplanck/neuplanck.#anyontgd](http://tgdtheory.fi/public_html/neuplanck/neuplanck.#anyontgd), 2006.
- [K18] Pitkänen M. TGD and Astrophysics. In *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdclass/tgdclass.html#astro](http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#astro), 2006.
- [K19] Pitkänen M. TGD and Cosmology. In *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdclass/tgdclass.html#cosmo](http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#cosmo), 2006.
- [K20] Pitkänen M. The Notion of Wave-Genome and DNA as Topological Quantum Computer. In *Genes and Memes*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/genememe/genememe.html#gari](http://tgdtheory.fi/public_html/genememe/genememe.html#gari), 2006.
- [K21] Pitkänen M. The Recent Status of Lepto-hadron Hypothesis. In *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/neuplanck/neuplanck.html#leptc](http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#leptc), 2006.
- [K22] Pitkänen M. The Relationship Between TGD and GRT. In *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdclass/tgdclass.html#tgdgrt](http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#tgdgrt), 2006.
- [K23] Pitkänen M. Three new physics realizations of the genetic code and the role of dark matter in bio-systems. In *Genes and Memes*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/genememe/genememe.html#dnatqccodes](http://tgdtheory.fi/public_html/genememe/genememe.html#dnatqccodes), 2006.
- [K24] Pitkänen M. Was von Neumann Right After All? In *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/neuplanck/neuplanck.html#vNeumann](http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#vNeumann), 2006.
- [K25] Pitkänen M. WCW Spinor Structure. In *Quantum Physics as Infinite-Dimensional Geometry*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdgeom/tgdgeom.html#cspin](http://tgdtheory.fi/public_html/tgdgeom/tgdgeom.html#cspin), 2006.

## Articles about TGD

- [L1] Pitkänen M. Further Progress in Nuclear String Hypothesis. Available at: <http://tgdtheory.fi/articles/nuclstring.pdf>, 2007.
- [L2] Pitkänen M. CMAP representations about TGD. Available at: <http://www.tgdtheory.fi/cmaphtml.html>, 2014.
- [L3] Pitkänen M. CMAP representations about TGD, and TGD inspired theory of consciousness and quantum biology. Available at: <http://www.tgdtheory.fi/tgdglossary.pdf>, 2014.
- [L4] Pitkänen M. Geometrization of fields. Available at: <http://www.tgdtheory.fi/webCMAPs/Geometrizationoffields.html>. 2014.
- [L5] Pitkänen M. Hierarchy of Planck constants. Available at: <http://www.tgdtheory.fi/webCMAPs/HierarchyofPlanckconstants.html>. 2014.

- [L6] Pitkänen M. Magnetic body. Available at: <http://www.tgdtheory.fi/webCMAPs/Magneticbody.html>. 2014.