

Recent Status of Lepto-Hadron Hypothesis

M. Pitkänen,

February 14, 2018

Email: matpitka6@gmail.com.

http://tgdtheory.com/public_html/.

Recent postal address: Rinnekatu 2-4 A 8, 03620, Karkkila, Finland.

Contents

1	Introduction	4
2	Lepto-Hadron Hypothesis	7
2.1	Anomalous E^+E^- Pairs In Heavy Ion Collisions	7
2.2	Lepto-Pions And Generalized PCAC Hypothesis	9
2.2.1	PCAC for ordinary pions	9
2.2.2	PCAC in leptonic sector	10
2.3	Lepto-Pion Decays And PCAC Hypothesis	12
2.3.1	$\Gamma(\pi_L \rightarrow \gamma\gamma)$	12
2.3.2	Lepto-pion-lepton coupling	13
2.3.3	$\Gamma(\pi_L \rightarrow e + \bar{\nu}_e)$	13
2.3.4	$\Gamma(\pi_L/\sigma_L \rightarrow e_{ex}\bar{e}_{ex})$ and $\Gamma(\pi_L/\sigma_L \rightarrow e_{ex}\bar{e})$	13
2.3.5	$\Gamma(e_{ex} \rightarrow e + \gamma)$	14
2.3.6	Some implications	15
2.3.7	Karmen anomaly	15
2.4	Lepto-Pions And Weak Decays	15
2.4.1	Lepto-hadrons and lepton decays	16
2.4.2	Lepto-pions and beta decay	16
2.5	Ortopositronium Puzzle And Lepto-Pion In Photon Photon Scattering	17
2.6	Spontaneous Vacuum Expectation Of Lepto-Pion Field As Source Of Lepto-Pions	18
2.7	Sigma Model And Creation Of Lepto-Hadrons In Electromagnetic Fields	20
2.7.1	Why sigma model approach?	20
2.7.2	Simplest sigma model	21
2.8	Classical Model For Lepto-Pion Production	23
2.8.1	General considerations	23

2.8.2	Expression of the classical cross section	24
2.9	Quantum Model For Lepto-Pion Production	25
2.9.1	Two possible approaches	25
2.9.2	Production amplitude	27
2.9.3	Differential cross section in the quantum model	29
2.9.4	Calculation of the lepto-pion production amplitude in the quantum model	31
2.9.5	Formula for the production cross section	31
2.9.6	Dominating contribution to production cross section and diffractive effects	32
2.9.7	Cutoff length scales	33
2.9.8	Numerical estimate for the electro-pion production cross section	34
2.9.9	$e_{ex}^+e_{ex}^-$ pairs from lepto-pions or e^+e^- pairs from lepto-sigmas?	35
2.9.10	Summary	36
3	Further Developments	37
3.1	How To Observe Leptonic Color?	37
3.2	New Experimental Evidence	38
3.2.1	Could lepto-hadrons correspond to dark matter?	38
3.2.2	Lightnings and lepto-pions	39
3.2.3	Lightnings, dark matter, and lepto-pion hypothesis again	40
3.2.4	Experimental evidence for colored muons	42
3.3	Evidence For τ -Hadrons	42
3.3.1	The first gift	43
3.3.2	The second gift	43
3.3.3	Explanation of the CDF anomaly	43
3.3.4	PAMELA and ATIC anomalies	45
3.3.5	Comparison of TGD model with the model of CDF collaboration	46
3.3.6	Numerical estimate for the production cross section	48
3.3.7	Does the production of lepto-pions involve a phase transition increasing Planck constant?	50
3.3.8	Tau-pions again but now as dark matter candidate in galactic center	51
3.3.9	Could it have been otherwise?	52
3.4	Dark Matter Puzzle	53
3.4.1	How to detect dark matter and what's the problem?	53
3.4.2	TGD based explanation of the DAMA events and related anomalies	53
3.4.3	TGD based explanation for the dark matter puzzle	54
3.4.4	Has Fermi observed dark matter?	55
3.4.5	Two different lifetimes for neutron as evidence for dark protons	56
3.5	Has Pamela Observed Evidence For The Non-Dark Electro-PionOf M_{89} Lepto-Hadron Physics?	57
3.6	Positron anomaly nine years later	58
3.6.1	Dark matter explanations for the positron excess	59
3.6.2	Standard physics explanation for the positron excess	59
3.6.3	TGD based model for positron excess	59
3.6.4	Other evidence for dark pion like states	60
3.7	Could Lepto-Hadrons Be Replaced With Bound States Of Exotic Quarks?	60
3.8	About The Masses Of Lepto-Hadrons	61
3.9	Do X And Y Mesons Provide Evidence For Color Excited Quarks Or Squarks?	61
3.9.1	Could these be the good questions?	62
3.9.2	Both leptons and quarks have color excitations in TGD Universe	62
3.9.3	Also squarks could explain exotic charmonium states	63
3.9.4	Could exotic charmonium states consist of color excited c and \bar{c} or of their spartners?	63
3.9.5	Why the color excitations/spartners of light quarks would be effectively absent?	64
3.9.6	The option based on heavy color excitations/spartners of light quarks	66
3.9.7	How to test the dark squark option?	66
3.9.8	Objection against covariantly constant neutrinos as SUSY generators	67

4	Appendix	68
4.1	Evaluation Of Lepto-Pion Production Amplitude	68
4.1.1	General form of the integral	68
4.1.2	k_y -integration	69
4.1.3	k_x -integration	70
4.2	Production Amplitude In Quantum Model	77
4.2.1	General expression of the production amplitude	77
4.2.2	The limit $\Delta k = 0$	77
4.2.3	Improved approximation to the production cross section	80
4.3	Evaluation Of The Singular Parts Of The Amplitudes	80

Abstract

TGD suggests strongly the existence of lepto-hadron physics. Lepto-hadrons would be bound states of color excited leptons and the anomalous production of e^+e^- pairs in heavy ion collisions finds a nice explanation as resulting from the decays of lepto-hadrons with basic condensate level $k = 127$ and having typical mass scale of one MeV . The recent indications on the existence of a new fermion with quantum numbers of muon neutrino and the anomaly observed in the decay of orthopositronium give further support for the lepto-hadron hypothesis. There is also evidence for anomalous production of low energy photons and e^+e^- pairs in hadronic collisions.

The identification of lepto-hadrons as a particular instance in the predicted hierarchy of dark matters interacting directly only via graviton exchange allows to circumvent the lethal counter arguments against the lepto-hadron hypothesis (Z^0 decay width and production of colored lepton jets in e^+e^- annihilation) even without assumption about the loss of asymptotic freedom.

PCAC hypothesis and its sigma model realization lead to a model containing only the coupling of the lepto-pion to the axial vector current as a free parameter. The prediction for e^+e^- production cross section is of correct order of magnitude only provided one assumes that lepto-pions decay to lepto-nucleon pair $e_{e_x}^+e_{e_x}^-$ first and that lepto-nucleons, having quantum numbers of electron and having mass only slightly larger than electron mass, decay to lepton and photon. The peculiar production characteristics are correctly predicted. There is some evidence that the resonances decay to a final state containing $n > 2$ particle and the experimental demonstration that lepto-nucleon pairs are indeed in question, would be a breakthrough for TGD.

During 18 years after the first published version of the model also evidence for colored μ has emerged. Towards the end of 2008 CDF anomaly gave support for the colored excitation of τ . The lifetime of the light long lived state identified as a charged τ -pion comes out correctly and the identification of the reported 3 new particles as p-adically scaled up variants of neutral τ -pion predicts their masses correctly. The observed muon jets can be understood in terms of the special reaction kinematics for the decays of neutral τ -pion to 3 τ -pions with mass scale smaller by a factor 1/2 and therefore almost at rest. A spectrum of new particles is predicted. The discussion of CDF anomaly led to a modification and generalization of the original model for lepto-pion production and the predicted production cross section is consistent with the experimental estimate.

1 Introduction

TGD suggest strongly (“predicts” is perhaps too strong expression) the existence of color excited leptons. The mass calculations based on p-adic thermodynamics and p-adic conformal invariance lead to a rather detailed picture about color excited leptons.

1. The simplest color excited neutrinos and charged leptons belong to the color octets ν_8 and L_{10} and $L_{\bar{10}}$ decouplet representations respectively and lepto-hadrons are formed as the color singlet bound states of these and possible other representations. Electro-weak symmetry suggests strongly that the minimal representation content is octet and decouplets for both neutrinos and charged leptons.
2. The basic mass scale for lepto-hadron physics is completely fixed by p-adic length scale hypothesis. The first guess is that color excited leptons have the levels $k = 127, 113, 107, \dots$ ($p \simeq 2^k$, k prime or power of prime) associated with charged leptons as primary condensation levels. p-Adic length scale hypothesis allows however also the level $k = 11^2 = 121$ in case of electronic lepto-hadrons. Thus both $k = 127$ and $k = 121$ must be considered as a candidate for the level associated with the observed lepto-hadrons. If also lepto-hadrons correspond non-perturbatively to exotic Super Virasoro representations, lepto-pion mass relates to pion mass by the scaling factor $L(107)/L(k) = k^{(107-k)/2}$. For $k = 121$ one has $m_{\pi_L} \simeq 1.057$ MeV which compares favorably with the mass $m_{\pi_L} \simeq 1.062$ MeV of the lowest observed state: thus $k = 121$ is the best candidate contrary to the earlier beliefs. The mass spectrum of lepto-hadrons is expected to have same general characteristics as hadronic mass spectrum and a satisfactory description should be based on string tension concept. Regge slope is

predicted to be of order $\alpha' \simeq 1.02/MeV^2$ for $k = 121$. The masses of ground state lepto-hadrons are calculable once primary condensation levels for colored leptons and the CKM matrix describing the mixing of color excited lepton families is known.

The strongest counter arguments against color excited leptons are the following ones.

1. The decay widths of Z^0 and W boson allow only $N = 3$ light particles with neutrino quantum numbers. The introduction of new light elementary particles seems to make the decay widths of Z^0 and W intolerably large.
2. Lepto-hadrons should have been seen in e^+e^- scattering at energies above few MeV . In particular, lepto-hadronic counterparts of hadron jets should have been observed.

A possible resolution of these problems is provided by the loss of asymptotic freedom in lepto-hadron physics. Lepto-hadron physics would effectively exist in a rather limited energy range about one MeV .

The development of the ideas about dark matter hierarchy [K7, K14, K5, K4] led however to a much more elegant solution of the problem.

1. TGD predicts an infinite hierarchy of various kinds of dark matters which in particular means a hierarchy of color and electro-weak physics with weak mass scales labelled by appropriate p-adic primes different from M_{89} : the simplest option is that also ordinary photons and gluons are labelled by M_{89} .
2. There are number theoretical selection rules telling which particles can interact with each other. The assignment of a collection of primes to elementary particle as characterizer of p-adic primes characterizing the particles coupling directly to it, is inspired by the notion of infinite primes [K15]. and discussed in [K7]. Only particles characterized by integers having common prime factors can interact by the exchange of elementary bosons: the p-adic length scale of boson corresponds to a common primes.
3. Also the physics characterized by different values of \hbar are dark with respect to each other as far quantum coherent gauge interactions are considered. Laser beams might well correspond to photons characterized by p-adic prime different from M_{89} and de-coherence for the beam would mean decay to ordinary photons. De-coherence interaction involves scaling down of the Compton length characterizing the size of the space-time of particle implying that particles do not anymore overlap so that macroscopic quantum coherence is lost.
4. Those dark physics which are dark relative to each other can interact only via graviton exchange. If lepto-hadrons correspond to a physics for which weak bosons correspond to a p-adic prime different from M_{89} , intermediate gauge bosons cannot have direct decays to colored excitations of leptons irrespective of whether the QCD in question is asymptotically free or not. Neither are there direct interactions between the QED:s and QCD:s in question if M_{89} characterizes also ordinary photons and gluons. These ideas are discussed and applied in detail in [K7, K14, K5].

Skeptic reader might stop the reading after these counter arguments unless there were definite experimental evidence supporting the lepto-hadron hypothesis.

1. The production of anomalous e^+e^- pairs in heavy ion collisions (energies just above the Coulomb barrier) suggests the existence of pseudo-scalar particles decaying to e^+e^- pairs. A natural identification is as lepto-pions that is bound states of color octet excitations of e^+ and e^- .
2. The second puzzle, Karmen anomaly, is quite recent [C20]. It has been found that in charge pion decay the distribution for the number of neutrinos accompanying muon in decay $\pi \rightarrow \mu + \nu_\mu$ as a function of time seems to have a small shoulder at $t_0 \sim ms$. A possible explanation is the decay of charged pion to muon plus some new weakly interacting particle with mass of order $30 MeV$ [C7] : the production and decay of this particle would proceed via mixing with muon neutrino. TGD suggests the identification of this state as color singlet leptobaryon of, say type $L_B = f_{abc}L_8^aL_8^b\bar{L}_8^c$, having electro-weak quantum numbers of neutrino.

3. The third puzzle is the anomalously high decay rate of orto-positronium. [C35]. e^+e^- annihilation to virtual photon followed by the decay to real photon plus virtual lepto-pion followed by the decay of the virtual lepto-pion to real photon pair, $\pi_L\gamma\gamma$ coupling being determined by axial anomaly, provides a possible explanation of the puzzle.
4. There exists also evidence for anomalously large production of low energy e^+e^- pairs [C19, C28, C21, C47] in hadronic collisions, which might be basically due to the production of lepto-hadrons via the decay of virtual photons to colored leptons.

In this chapter a revised form of lepto-hadron hypothesis is described.

1. Sigma model realization of PCAC hypothesis allows to determine the decay widths of lepto-pion and lepto-sigma to photon pairs and e^+e^- pairs. Ortopositronium anomaly determines the value of $f(\pi_L)$ and therefore the value of lepto-pion-lepto-nucleon coupling and the decay rate of lepto-pion to two photons. Various decay widths are in accordance with the experimental data and corrections to electro-weak decay rates of neutron and muon are small.
2. One can consider several alternative interpretations for the resonances.

Option 1: For the minimal color representation content, three lepto-pions are predicted corresponding to 8, 10, $\bar{10}$ representations of the color group. If the lightest lepto-nucleons e_{ex} have masses only slightly larger than electron mass, the anomalous e^+e^- could be actually $e_{ex}^+ + e_{ex}^-$ pairs produced in the decays of lepto-pions. One could identify 1.062, 1.63 and 1.77 MeV states as the three lepto-pions corresponding to 8, 10, $\bar{10}$ representations and also understand why the latter two resonances have nearly degenerate masses. Since d and s quarks have same primary condensation level and same weak quantum numbers as colored e and μ , one might argue that also colored e and μ correspond to $k = 121$. From the mass ratio of the colored e and μ , as predicted by TGD, the mass of the muonic lepto-pion should be about 1.8 MeV in the absence of topological mixing. This suggests that 1.83 MeV state corresponds to the lightest $g = 1$ lepto-pion.

Option 2: If one believes sigma model (in ordinary hadron physics the existence of sigma meson is not established and its width is certainly very large if it exists), then lepto-pions are accompanied by sigma scalars. If lepto-sigmas decay dominantly to e^+e^- pairs (this might be forced by kinematics) then one could adopt the previous scenario and could identify 1.062 state as lepto-pion and 1.63, 1.77 and 1.83 MeV states as lepto-sigmas rather than lepto-pions. The fact that muonic lepto-pion should have mass about 1.8 MeV in the absence of topological mixing, suggests that the masses of lepto-sigma and lepto-pion should be rather close to each other.

Option 3: One could also interpret the resonances as string model “satellite states” having interpretation as radial excitations of the ground state lepto-pion and lepto-sigma. This identification is not however so plausible as the genuinely TGD based identification and will not be discussed in the sequel.

3. PCAC hypothesis and sigma model leads to a general model for lepto-hadron production in the electromagnetic fields of the colliding nuclei and production rates for lepto-pion and other lepto-hadrons are closely related to the Fourier transform of the instanton density $\vec{E} \cdot \vec{B}$ of the electromagnetic field created by nuclei. The first source of anomalous e^+e^- pairs is the production of $\sigma_L\pi_L$ pairs from vacuum followed by $\sigma_L \rightarrow e^+e^-$ decay. If $e_{ex}^+e_{ex}^-$ pairs rather than genuine e^+e^- pairs are in question, the production is production of lepto-pions from vacuum followed by lepto-pion decay to lepto-nucleon pair.

Option 1: For the production of lepto-nucleon pairs the cross section is only slightly below the experimental upper bound for the production of the anomalous e^+e^- pairs and the decay rate of lepto-pion to lepto-nucleon pair is of correct order of magnitude.

Option 2: The rough order of magnitude estimate for the production cross section of anomalous e^+e^- pairs via $\sigma_l\pi_l$ pair creation followed by $\sigma_L \rightarrow e^+e^-$ decay, is by a factor of order $1/\sum N_c^2$ (N_c is the total number of states for a given colour representation and sum over the representations contributing to the ortopositronium anomaly appears) smaller than the reported cross section in case of 1.8 MeV resonance. The discrepancy could be due to the

neglect of the large radiative corrections (the coupling $g(\pi_L\pi_L\sigma_L) = g(\sigma_L\sigma_L\sigma_L)$ is very large) and also due to the uncertainties in the value of the measured cross section.

Given the unclear status of sigma in hadron physics, one has a temptation to conclude that anomalous e^+e^- pairs actually correspond to lepto-nucleon pairs.

4. The vision about dark matter suggests that direct couplings between leptons and lepto-hadrons are absent in which case no new effects in the direct interactions of ordinary leptons are predicted. If colored leptons couple directly to ordinary leptons, several new physics effects such as resonances in photon-photon scattering at cm energy equal to lepto-pion masses and the production of $e_{ex}\bar{e}_{ex}$ (e_{ex} is leptobaryon with quantum numbers of electron) and $e_{ex}\bar{e}$ pairs in heavy ion collisions, are possible. Lepto-pion exchange would give dominating contribution to $\nu - e$ and $\bar{\nu} - e$ scattering at low energies. Lepto-hadron jets should be observed in e^+e^- annihilation at energies above few MeV:s unless the loss of asymptotic freedom restricts lepto-hadronic physics to a very narrow energy range and perhaps to entirely non-perturbative regime of lepto-hadronic QCD.

This chapter is a revised version of the earlier chapter [K1] and still a work in progress. I apologize for the reader for possible inconvenience. The motivation for the re-writing came from the evidence for the production of τ -pions in high energy proton-antiproton collisions [C10]. Since the kinematics of these collisions differs dramatically from that for heavy ion collisions, a critical re-examination of the earlier model - which had admittedly somewhat ad hoc character- became necessary. As a consequence the earlier model simplified dramatically. As far as basic calculations are considered, the modification makes itself visible only at the level of coefficients. Even more remarkably, it turned out possible to calculate exactly the lepto-pion production amplitude under a very natural approximation, which can be also generalized so that the calculation of production amplitude can be made analytically in high accuracy and only the integration over lepto-pion momentum must be carried out numerically. As a consequence, a rough analytic estimate for the production cross section follows and turns out to be of correct order of magnitude. It must be however stressed that the cross section is highly sensitive to the value of the cutoff parameter (at least in this naive estimate) and only a precise calculation can settle the situation.

The appendix of the book gives a summary about basic concepts of TGD with illustrations. Pdf representation of same files serving as a kind of glossary can be found at <http://tgdtheory.fi/tgdglossary.pdf> [L2].

2 Lepto-Hadron Hypothesis

2.1 Anomalous E^+E^- Pairs In Heavy Ion Collisions

Heavy ion collision experiments carried out at the Gesellschaft für Schwerionenforschung in Darmstadt, West Germany [C32, C23, C24, C33] have yielded a rather puzzling set of results. The expectation was that in heavy ion collisions in which the combined charge of the two colliding ions exceeds 173, a composite nucleus with $Z > Z_{cr}$ would form and the probability for spontaneous positron emission would become appreciable.

Indeed, narrow peaks of widths of roughly 50-70 keV and energies about 350 ± 50 keV were observed in the positron spectra but it turned out that the position of the peaks seems to be a constant function of Z rather than vary as Z^{20} as expected and that peaks are generated also for Z smaller than the critical Z. The collision energies at which peaks occur lie in the neighborhood of 5.7-6 MeV/nucleon. Also it was found that positrons are accompanied by e^- emission. Data are consistent with the assumption that some structure at rest in cm is formed and decays subsequently to e^+e^- pair.

Various theoretical explanations for these peaks have been suggested [C4, C41]. For example, lines might be created by pair conversion in the presence of heavy nuclei. In nuclear physics explanations the lines are due to some nuclear transition that occurs in the compound nucleus formed in the collision or in the fragments formed. The Z-independence of the peaks seems however to exclude both atomic and nuclear physics explanations [C4]. Elementary particle physics explanations [C4, C41] seem to be excluded already by the fact that several peaks have been observed in the range 1.6 – 1.8 MeV with widths of order $10^1 - 10^2$ keV. These states decay to e^+e^- pairs.

There is evidence for one narrow peak with width of order one keV at 1.062 Mev [C4]: this state decays to photon-photon pairs.

Thus it seems that the structures produced might be composite, perhaps resonances in e^+e^- system. The difficulty of this explanation is that conventional QED seems to offer no natural explanation for the strong force needed to explain the energy scale of the states. One idea is that the strong electromagnetic fields create a new phase of QED [C4] and that the resonances are analogous to pseudo-scalar mesons appearing as resonances in strongly interacting systems.

TGD based explanation relies on the following hypothesis motivated by Topological Geometro-dynamics.

1. Ordinary leptons are not point like particles and can have colored excitations, which form color singlet bound states. A natural identification for the primary condensate level is $k = 121$ so that the mass scale is of order one MeV for the states containing lowest generation colored leptons. The fact that d and s quarks, having the same weak quantum numbers as charged leptons, have same primary condensation level, suggests that both colored electron and muon condense to the same level. The expectation that lepto-hadron physics exists in a narrow energy interval only, suggests that also colored τ should condense on the same level.
2. The states in question are lepto-hadrons, that is color confined states formed from the colored excitations of e^+ and e^- . The decay rate to lepto-nucleon pairs $e_{ex}^+e_{ex}^-$ is large and turns out to give rise to correct order of magnitude for the decay width. Hence two options emerge.

Option 1: Lepto-nucleons e_{ex} have masses only slightly above the electron mass and since they behave like electrons, anomalous e^+e^- pairs could actually correspond to lepto-nucleon pairs created in the decays of lepto-pions. 1.062, 1.63 and 1.77 MeV states can be identified as lowest generation lepto-pions correspond to octet and two decouplets. 1.83 MeV state could be identified as the second generation lepto-pion corresponding to colored muon. The small branching fraction to gamma pairs explains why the decays of the higher mass lepto-pions to gamma pairs has not been observed. $g = 0$ lepto-pion decays to lepto-nucleon pairs can be visualized as occurring via dual diagrams obeying Zweig's rule (annihilation is not allowed inside incoming or outgoing particle states). The decay of $g = 1$ colored muon pair occurs via Zweig rule violating annihilation to two gluon intermediate state, which transforms back to virtual $g = 0$ colored electron pair decaying via dual diagram: the violation of Zweig's rule suggests that the decay rate for 1.8 MeV state is smaller than for the lighter states. Quantitative model shows that this scenario is the most plausible one.

Option 2: Lepto-sigmas, which are the scalar partners of lepto-pions predicted by sigma model, are the source of anomalous (and genuine) e^+e^- pairs. In this case 1.062 state must correspond to lepto-pion whereas higher states must be identified as lepto-sigmas. Also now new lepto-pion states decaying to gamma pairs are predicted and one could hence argue that this prediction is not consistent with what has been observed. A crucial assumption is that lepto-sigmas are light and cannot decay to other lepto-mesons. Ordinary hadronic physics suggests that this need not be the case: the hadronic decay width of the ordinary sigma, if it exists, is very large.

The program of the section is following:

1. PCAC hypothesis, successful in low energy pion physics, is generalized to the case of lepto-pion. Hypothesis allows to deduce the coupling of lepto-pion to leptons and lepto-baryons in terms of leptobaryon-lepton mixing angles. Ortopositronium anomaly allows to deduce precise value of $f(\pi_L)$ characterizing the decay rate of lepto-pion so that the crucial parameters of the model are completely fixed. The decay rates of lepto-pion to photon pair and of lepto-sigma to ordinary e^+e^- pairs are within experimental bounds and corrections to muon and beta decay rates are small. New calculable resonance contributions to photon-photon scattering at cm energy equal to lepto-pion masses are predicted.
2. If anomalous e^+e^- pairs are actually lepto-nucleon pairs, only a model for the creation of lepto-pions from vacuum is needed. In an external electromagnetic field lepto-pion develops a vacuum expectation value proportional to electromagnetic anomaly term [B1] so that the production amplitude for the lepto-pion is essentially the Fourier transform of the scalar

product of the electric field of the stationary target nucleus with the magnetic field of the colliding nucleus.

3. If anomalous e^+e^- pairs are produced in the decays of lepto-sigmas, the starting point is sigma model providing a realization of PCAC hypothesis. Sigma model makes it possible to relate the production amplitude for $\sigma_L\pi_L$ pairs to the lepto-pion production amplitude: the key element of the model is the large value of the $\sigma\pi_L\pi_L$ coupling constant.
4. Lepto-hadron production amplitudes are proportional to lepto-pion production amplitude and this motivates a detailed study of lepto-pion production. Two models for lepto-pion production are developed: in classical model colliding nucleus is treated classically whereas in quantum model the colliding nucleus is described quantum mechanically. It turns out that classical model explains the peculiar production characteristics of lepto-pion but that production cross section is too small by several orders of magnitude. Quantum mechanical model predicts also diffractive effects: production cross section varies rapidly as a function of the scattering angle and for a fixed value of scattering angle there is a rapid variation with the collision velocity. The estimate for the total lepto-pion production cross section increases by several orders of magnitude due to the coherent summation of the contributions to the amplitude from different values of the impact parameter at the peak.
5. The production rate for lepto-nucleon pairs is only slightly smaller than the experimental upper bound but the e^+e^- production rate predicted by sigma model approach is still by a factor of order $1/\sum N_c^2$ smaller than the reported maximum cross section. A possible explanation for this discrepancy is the huge value of the coupling $g(\pi_L, \pi_L, \sigma_L) = g(\sigma_L, \sigma_L, \sigma_L)$ implying that the diagram involving the exchange of virtual sigma can give the dominant contribution to the production cross section of $\sigma_L\pi_L$ pair.

2.2 Lepto-Pions And Generalized PCAC Hypothesis

One can say that the PCAC hypothesis predicts the existence of pions and a connection between the pion nucleon coupling strength and the pion decay rate to leptons. In the following we give the PCAC argument and its generalization and consider various consequences.

2.2.1 PCAC for ordinary pions

The PCAC argument for ordinary pions goes as follows [B3] :

1. Consider the contribution of the hadronic axial current to the matrix element describing lepton nucleon scattering (say $N + \nu \rightarrow P + e^-$) by weak interactions. The contribution in question reduces to the well-known current-current form

$$\begin{aligned}
 M &= \frac{G_F}{\sqrt{2}} g_A L_\alpha \langle P | A^\alpha | P \rangle , \\
 L_\alpha &= \bar{e} \gamma_\alpha (1 + \gamma_5) \nu , \\
 \langle P | A^\alpha | P \rangle &= \bar{P} \gamma^\alpha N ,
 \end{aligned} \tag{2.1}$$

where $G_F = \frac{\pi\alpha}{2m_W^2 \sin^2(\theta_W)} \simeq 10^{-5}/m_p^2$ denotes the dimensional weak interaction coupling strength and g_A is the nucleon axial form factor: $g_A \simeq 1.253$.

2. The matrix element of the hadronic axial current is not divergenceless, due to the non-vanishing nucleon mass,

$$a_\alpha \langle P | A^\alpha | P \rangle \simeq 2m_p \bar{P} \gamma_5 N . \tag{2.2}$$

Here q^α denotes the momentum transfer vector. In order to obtain divergenceless current, one can modify the expression for the matrix element of the axial current

$$\langle P|A^\alpha|N\rangle \rightarrow \langle P|A^\alpha|N\rangle - q^\alpha 2m_p \bar{P}\gamma_5 N \frac{1}{q^2} . \quad (2.3)$$

3. The modification introduces a new term to the lepton-hadron scattering amplitude identifiable as an exchange of a massless pseudo-scalar particle

$$\delta T = \frac{G_F g_A}{\sqrt{2}} L_\alpha \frac{2m_p q^\alpha}{q^2} \bar{P}\gamma_5 N . \quad (2.4)$$

The amplitude is identifiable as the amplitude describing the exchange of the pion, which gets its mass via the breaking of chiral invariance and one obtains by the straightforward replacement $q^2 \rightarrow q^2 - m_\pi^2$ the correct form of the amplitude.

4. The nontrivial point is that the interpretations as pion exchange is indeed possible since the amplitude obtained is to a good approximation identical to that obtained from the Feynman diagram describing pion exchange, where the pion nucleon coupling constant and pion decay amplitude appear

$$T_2 = \frac{G}{\sqrt{2}} f_\pi q^\alpha L_\alpha \frac{1}{q^2 - m_\pi^2} g \sqrt{2} \bar{P}\gamma_5 N . \quad (2.5)$$

The condition $\delta T \sim T_2$ gives from Goldberger-Treiman [B3]

$$g_A (\simeq 1.25) = \sqrt{2} \frac{f_\pi g}{2m_p} (\simeq 1.3) , \quad (2.6)$$

satisfied in a good accuracy experimentally.

2.2.2 PCAC in leptonic sector

A natural question is why not generalize the previous argument to the leptonic sector and look at what one obtains. The generalization is based on following general picture.

1. There are two levels to be considered: the level of ordinary leptons and the level of leptobaryons of, say type $f_{ABC} \nu_8^A \nu_8^B \bar{L}_{10}^C$, possessing same quantum numbers as leptons. The interaction transforming these states to each other causes in mass eigenstates mixing of leptobaryons with ordinary leptons described by mixing angles. The masses of lepton and corresponding leptobaryon could be quite near to each other and in case of electron this should be the case as it turns out.
2. A counterargument against the applications of PCAC hypothesis at level of ordinary leptons is that baryons and mesons are both bound states of quarks whereas ordinary leptons are not bound states of colored leptons. The divergence of the axial current is however completely independent of the possible internal structure of leptons and microscopic emission mechanism. Ordinary lepton cannot emit lepto-pion directly but must first transform to leptobaryon with same quantum numbers: phenomenologically this process can be described using mixing angle $\sin(\theta_B)$. The emission of lepto-pion proceeds as $L \rightarrow B_L : B_L \rightarrow B_L + \pi_L : B_L \rightarrow L$, where B_L denotes leptobaryon of type structure $f_{ABC} L_8^A L_8^B \bar{L}_8^C$. The transformation amplitude $L \rightarrow B_L$ is proportional to the mixing angle $\sin(\theta_L)$.

Three different PCAC type identities are assumed to hold true:

PCAC1) The vertex for the emission of lepto-pion by ordinary lepton is equivalent with the graph in which lepton L transforms to leptobaryon L^{ex} with same quantum numbers, emits lepto-pion

and transforms back to ordinary lepton. The assumption relates the couplings $g(L_1, L_2)$ and $g(L_1^{ex}, L_2^{ex})$ (analogous to strong coupling) and mixing angles to each other

$$g(L_1, L_2) = g(L_1^{ex}, L_2^{ex}) \sin(\theta_1) \sin(\theta_2) . \quad (2.7)$$

The condition implies that in electro-weak interactions ordinary leptons do not transform to their exotic counterparts.

PCAC2) The generalization of the ordinary Goldberger-Treiman argument holds true, when ordinary baryons are replaced with leptobaryons. This gives the condition expressing the coupling $f(\pi_L)$ of the lepto-pion state to axial current defined as

$$\langle vac | A_\alpha | \pi_L \rangle = i p_\alpha f(\pi_L) , \quad (2.8)$$

in terms of the masses of leptobaryons and strong coupling g .

$$f(\pi_L) = \sqrt{2} g_A \frac{(m_{ex}(1) + m_{ex}(2)) \sin(\theta_1) \sin(\theta_2)}{g(L_1, L_2)} , \quad (2.9)$$

where g_A is parameter characterizing the deviation of weak coupling strength associated with leptobaryon from ideal value: $g_A \sim 1$ holds true in good approximation.

PCAC3) The elimination of leptonic axial anomaly from leptonic current fixes the values of $g(L_i, L_j)$.

1. The standard contribution to the scattering of leptons by weak interactions given by the expression

$$\begin{aligned} T &= \frac{G_F}{\sqrt{2}} \langle L_1 | A^\alpha | L_2 \rangle \langle L_3 | A_\alpha | L_4 \rangle , \\ \langle L_i | A^\alpha | L_j \rangle &= \bar{L}_i \gamma^\alpha \gamma_5 L_j . \end{aligned} \quad (2.10)$$

2. The elimination of the leptonic axial anomaly

$$q_\alpha \langle L_i | A^\alpha | L_j \rangle = (m(L_i) + m(L_j)) \bar{L}_i \gamma_5 L_j , \quad (2.11)$$

by modifying the axial current by the anomaly term

$$\langle L_i | A^\alpha | L_j \rangle \rightarrow \langle L_i | A^\alpha | L_j \rangle - (m(L_i) + m(L_j)) \frac{q^\alpha}{q^2} \bar{L}_i \gamma_5 L_j , \quad (2.12)$$

induces a new interaction term in the scattering of ordinary leptons.

3. It is assumed that this term is equivalent with the exchange of lepto-pion. This fixes the value of the coupling constant $g(L_1, L_2)$ to

$$\begin{aligned} g(L_1, L_2) &= 2^{1/4} \sqrt{G_F} (m(L_1) + m(L_2)) \xi , \\ \xi(\text{charged}) &= 1 , \\ \xi(\text{neutral}) &= \cos(\theta_W) . \end{aligned} \quad (2.13)$$

Here the coefficient ξ is related to different values of masses for gauge bosons W and Z appearing in charged and neutral current interactions. An important factor 2 comes from the modification of the axial current in both matrix elements of the axial current.

Lepto-pion exchange interaction couples right and left handed leptons to each other and its strength is of the same order of magnitude as the strength of the ordinary weak interaction at energies not considerably large than the mass of the lepto-pion. At high energies this interaction is negligible and the existence of the lepto-pion predicts no corrections to the parameters of the standard model since these are determined from weak interactions at much higher energies. If lepto-pion mass is sufficiently small (as found, $m(\pi_L) < 2m_e$ is allowed by the experimental data), the interaction mediated by lepto-pion exchange can become quite strong due to the presence of the lepto-pion propagator. The value of the lepton-lepto-pion coupling is $g(e, e) \equiv g \sim 5.6 \cdot 10^{-6}$. It is perhaps worth noticing that the value of the coupling constant is of the same order as lepton-Higgs coupling constant and also proportional to the mass of the lepton.

PCAC identities fix the values of coupling constants apart from the values of mixing angles. If one assumes that the strong interaction mediated by lepto-pions is really strong and the coupling strength $g(L_{ex}, L_{ex})$ is of same order of magnitude as the ordinary pion nucleon coupling strength $g(\pi NN) \simeq 13.5$ one obtains an estimate for the value of the mixing angle $\sin(\theta_e)$

$\sin^2(\theta_e) \sim \frac{g(\pi NN)}{g(L,L)} \sim 2.4 \cdot 10^{-6}$. This implies the order of magnitude $f(\pi_L) \sim 10^{-6} m_W \sim 10^2 \text{ keV}$ for $f(\pi_L)$. The order of magnitude is correct as will be found. Ortopositronium decay rate anomaly $\Delta\Gamma/\Gamma \sim 10^{-3}$ and the assumption $m_{ex} \geq 1.3 \text{ MeV}$ (so that $e_{ex}\bar{e}$ decay is not possible) gives the upper bound $\sin(\theta_e) \leq x \cdot \sqrt{N_c} \cdot 10^{-4}$, where the value of $x \sim 1$ depends on the number of the lepto-pion type states and on the precise value of the Op anomaly.

2.3 Lepto-Pion Decays And PCAC Hypothesis

The PCAC argument makes it possible to predict the lepto-pion coupling and decay rates of the lepto-pion to various channels. Actually the orders of magnitude for the decay rates of the lepto-sigma and other lepto-mesons can be deduced also. The comparison with the experimental data is made difficult by the uncertainty of the identifications. The lightest candidate has mass 1.062 MeV and decay width of order 1 keV [C4]: only photon photon decay has been observed for this state. The next lepto-meson candidates are in the mass range $1.6 - 1.8 \text{ MeV}$. Perhaps the best status is possessed by ‘‘Darmstadtium’’ with mass 1.8 MeV . For these states decays to final states identified as e^+e^- pairs dominate: if indeed e^+e^- pairs, these states probably correspond to the decay products of lepto-sigma. Another possibility is that pairs are actually lepto-nucleon pairs with the mass of the lepto-nucleon only slightly larger than electron mass. Hadron physics experience suggests that the decay widths of the lepto-hadrons (lepto-pion forming a possible exception) should be about 1-10 per cent of particle mass as in hadron physics. The upper bounds for the widths are indeed in the range $50 - 70 \text{ keV}$ [C4].

2.3.1 $\Gamma(\pi_L \rightarrow \gamma\gamma)$

As in the case of the ordinary pion, anomaly considerations give the following approximate expression for the decay rate of the lepto-pion to two-photon final states [B1])

$$\Gamma(\pi_L \rightarrow \gamma\gamma) = \frac{N_c^2 \alpha^2 m^3(\pi_L)}{64 f(\pi_L)^2 \pi^3} . \quad (2.14)$$

$N_c = 8, 10$ is the number of the colored lepton states coming from the axial anomaly loop. For $m(\pi_L) = 1.062 \text{ MeV}$ and $f(\pi_L) = N_c \cdot 7.9 \text{ keV}$ implied by the ortopositronium decay rate anomaly $\Delta\Gamma/\Gamma = 10^{-3}$ one has $\Gamma(\gamma\gamma) = .52 \text{ keV}$, which is consistent with the experimental estimate of order 1 keV [C4].

In fact, several lepto-pion states could exist (4 at least corresponding to the resonances at 1.062, 1.63, 1.77 and 1.83 MeV). Since all these lepto-pion states contribute to Op decay rate, the actual value of $f(\pi_L)$ assumed to scale as $m(\pi_L)$, is actually larger in this case: it turns out that $f(\pi_L)$ for the lightest lepto-pion increases to $f(\pi_L)(\text{lightest}) = N_c \cdot 15 \text{ keV}$ and gives $\Gamma(\gamma\gamma) \simeq .13 \text{ keV}$ in case of the lightest lepto-pion if lepto-pions are assumed to correspond the resonances. Note that the order of magnitude for $f(\pi_L)$ is same as deduced from the assumption that lepto-hadronic counterpart of $g(\pi NN)$ equals to the ordinary $g(\pi NN)$. The increase of the ortopositronium anomaly by a factor of, say 4, implies corresponding decrease in $f(\pi_L)^2$. The value of $f(\pi_L)$ is also sensitive to the precise value of the mass of the lightest lepto-pion.

2.3.2 Lepto-pion-lepton coupling

The value of the lepto-pion-lepton coupling can be used to predict the decay rate of lepto-pion to leptons. One obtains for the decay rate $\pi_L^0 \rightarrow e^+e^-$ the estimate

$$\begin{aligned}\Gamma(\pi_L \rightarrow e^+e^-) &= 4 \frac{g(e,e)^2 \pi}{2(2\pi)^2} (1-4x^2) m(\pi_L) \\ &= 16 G m_e^2 \cos^2(\theta_W) \frac{\sqrt{2}}{4\pi} (1-4x^2) m(\pi_L) , \\ x &= \frac{m_e}{m(\pi_L)} .\end{aligned}\tag{2.15}$$

for the decay rate of the lepto-pion: for lepto-pion mass $m(\pi_L) \simeq 1.062 \text{ MeV}$ one obtains for the decay rate the estimate $\Gamma \sim 1/(1.3 \cdot 10^{-8} \text{ sec})$: the low decay rate is partly due to the phase space suppression and implies that e^+e^- decay products cannot be observed in the measurement volume. The low decay rate is in accordance with the identification of the lepto-pion as the $m = 1.062 \text{ MeV}$ lepto-pion candidate. In sigma model lepto-pion and lepto-sigma have identical lifetimes and for lepto-sigma mass of order 1.8 MeV one obtains $\Gamma(\sigma_L \rightarrow e^+e^-) \simeq 1/(8.2 \cdot 10^{-10} \text{ sec})$: the prediction is larger than the lower limit $\sim 1/(10^{-9} \text{ sec})$ for the decay rate implied by the requirement that σ_L decays inside the measurement volume. The estimates of the lifetime obtained from heavy ion collisions [C30] give the estimate $\tau \geq 10^{-10} \text{ sec}$. The large value of the lifetime is in accordance with the limits for the lifetime obtained from Bhabha scattering [C29], which indicate that the lifetime must be longer than 10^{-12} sec .

For lepto-meson candidates with mass above 1.6 MeV no experimental evidence for other decay modes than $X \rightarrow e^+e^-$ has been found and the empirical upper limit for $\gamma\gamma/e^+e^-$ branching ratio [C26] is $\Gamma(\gamma\gamma)/\Gamma(e^+e^-) \leq 10^{-3}$. If the identification of the decay products as e^+e^- pairs is correct then the only possible conclusion is that these states cannot correspond to lepto-pion since lepto-pion should decay dominantly into photon photon pairs. Situation changes if pairs of lepton-ucleons $e_{ex}\bar{e}_{ex}$ of type $e_{ex} = e_8\nu_8\bar{\nu}_8$ pair are in question.

I realized that this conclusion might be questioned for more than decade after writing the above text as I developed a model for CDF anomaly suggesting the existence of τ -pions. Since colored leptons are color octets, anomalous magnetic moment type coupling of form $\bar{L}Tr(F^{\mu\nu}\Sigma_{\mu\nu}L_8)$ (the trace is over the Lie-algebra generators of $SU(3)$ and $F^{\mu\nu}$ denotes color gauge field) between ordinary lepton, colored lepton and lepto-gluon is possible. The exchange of a virtual lepto-gluon allows lepto-pion to decay by lepto-strong interactions to electron-positron pairs. The decay rate is limited by the kinematics for the lightest state very near to the final state mass and might make decay rate to in this case very small. If the rate for the decay to electron-positron pair is comparable to that for the decay to two photons the production rate for electron-positron pairs could be of the same order of magnitude as lepto-pion production rate. The anomalous magnetic moment of electron however poses strong limitations on this coupling and it might be that the coupling is too small. This coupling could however induce the mixing of e_{ex} with e .

2.3.3 $\Gamma(\pi_L \rightarrow e + \bar{\nu}_e)$

The expression for the decay rate $\pi_L \rightarrow e + \bar{\nu}_e$ reads as

$$\begin{aligned}\Gamma(\pi_L^- \rightarrow e\nu_e) &= 8 G m_e^2 \frac{(1-x^2)^2}{2(1+x^2)} \frac{\sqrt{2}}{(2\pi)^5} m(\pi_L) , \\ &= \frac{4}{\cos^2(\theta_W)} \frac{(1-x^2)}{(1+x^2)(1-4x^2)} \Gamma(\pi_L^0 \rightarrow e^+e^-) ,\end{aligned}\tag{2.16}$$

and gives $\Gamma(\pi_L^- \rightarrow e\nu_e) \simeq 1/(3.6 \cdot 10^{-10} \text{ sec})$ for $m(\pi_L) = 1.062 \text{ MeV}$.

2.3.4 $\Gamma(\pi_L/\sigma_L \rightarrow e_{ex}\bar{e}_{ex})$ and $\Gamma(\pi_L/\sigma_L \rightarrow e_{ex}\bar{e})$

Sigma model predicts lepto-pion and lepto-sigma to have same coupling to lepto-nucleon e_{ex} pair so that in the sequel only lepto-pion decay rates are considered. One must consider also the

possibility that lepto-pion decay products are either $e_{ex}\bar{e}_{ex}$ or $e_{ex}\bar{e}$ pairs with e_{ex} having mass of near the mass of electron so that it could be misidentified as electron. If the mass of lepto-nucleon e_{ex} with quantum numbers of electron is smaller than $m(\pi_L)/2$ it can be produced in lepto-pion annihilation. One must also assume $m(e_{ex}) > m_e$: otherwise electrons would spontaneously decay to lepto-nucleons via photon emission. The production rate to lepto-nucleon pair can be written as

$$\begin{aligned}\Gamma(\pi_L \rightarrow e_{ex}^+ e_{ex}^-) &= \frac{1}{\sin^4(\theta_e)} \frac{(1-4y^2)}{(1-4x^2)} \Gamma(\pi_L \rightarrow e^+ e^-) , \\ y &= \frac{m(e_{ex})}{m(\pi_L)} .\end{aligned}\quad (2.17)$$

If $e - e_{ex}$ mass difference is sufficiently small the kinematic suppression does not differ significantly from that for e^+e^- pair. The limits from Bhabha scattering give no bounds on the rate of $\pi_L \rightarrow e_{ex}^+ e_{ex}^-$ decay. The decay rate $\Gamma \sim 10^{20}/sec$ implied by $\sin(\theta_e) \sim 10^{-4}$ implies decay width of order .1 MeV: the order of magnitude is the naively expected one and means that the decay to $e_{ex}^+ e_{ex}^-$ pairs dominates over the decay to gamma pairs except in the case of the lightest lepto-pion state for which the decay is kinematically forbidden.

The decay rate of the lepto-pion to $\bar{e}e_{ex}$ pair has sensible order of magnitude: for $\sin(\theta_e) = 1.2 \cdot 10^{-3}$, $m_{\sigma_L} = 1.8 MeV$ and $m_{e_{ex}} = 1.3 MeV$ one has $\Gamma \simeq 60 eV$ allowed by the experimental limits. This decay is kinematically possible only provided the mass of e_{ex} is in below $1.3 MeV$. These decays should dominate by a factor $1/\sin^2(\theta_e)$ over e^+e^- decays if kinematically allowed.

A signature of these events, if identified erratically as electron positron pairs, is the non-vanishing value of the energy difference in the cm frame of the pair: $E(e^-) - E(e^+) \simeq (m^2(e_{ex}) - m_e^2)/2E > 160 keV$ for $E = 1.8 MeV$. If the decay $e_{ex} \rightarrow e + \gamma$ takes place before the detection the energy asymmetry changes its sign. Energy asymmetry [C31] increasing with the rest energy of the decaying object has indeed been observed: the proposed interpretation has been that electron forms a bound state with the second nucleus so that its energy is lowered. Also a deviation from the momentum distribution implied by the decay of neutral particle to e^+e^- pair (momenta are opposite in the rest frame) results from the emission of photon. This kind of deviation has also been observed [C31]: the proposed explanation is that third object is involved in the decay. A possible alternative explanation for the asymmetries is the production mechanism ($\sigma_L \pi_L$ pairs instead of single particle states).

2.3.5 $\Gamma(e_{ex} \rightarrow e + \gamma)$

The decay to electron and photon would be a unique signature of e_{ex} . The general feature of fermion family mixing is that mixing takes place in charged currents. In present case mixing is of different type so that $e_{ex} \rightarrow e + \gamma$ might be allowed. If this is not the case then the decay takes place as weak decay via the emission of virtual W boson: $e_{ex} \rightarrow e + \nu_e + \bar{\nu}_e$ and is very slow due to the presence of mixing angle and kinematical suppression. The energy of the emitted photon is $E_\gamma = (m_{e_{ex}}^2 - m_e^2)/2m_e$. The decay rate $\Gamma(e_{ex} \rightarrow e + \gamma)$ is given by

$$\begin{aligned}\Gamma(e_{ex} \rightarrow e + \gamma) &= \alpha_{em} \sin^2(\theta_e) X m_e , \\ X &= \frac{(m_1 - m_e)^3 (m_1 + m_e) m_e}{(m_1^2 + m_e^2)^2 m_1} .\end{aligned}\quad (2.18)$$

For $m(e_{ex}) = 1.3 MeV$ the decay of order $1/(1.4 \cdot 10^{-12} sec)$ for $\sin(\theta_e) = 1.2 \cdot 10^{-3}$ so that lepto-nucleons would decay to electrons in the measurement volume. In the experiments positrons are identified via pair annihilation and since pair annihilation rate for \bar{e}_{ex} is by a factor $\sin^2(\theta_e)$ slower than for e^+ the particles identified as positrons must indeed be positrons. For sufficiently small mass difference $m(e_{ex}) - m_e$ the particles identified as electron could actually be e_{ex} . The decay of e_{ex} to electron plus photon before its detection seems however more reasonable alternative since it could explain the observed energy asymmetry [C31].

2.3.6 Some implications

The results have several implications as far as the decays of on mass shell states are considered:

1. For $m(e_{ex}) > 1.3 \text{ MeV}$ the only kinematically possible decay mode is the decay to e^+e^- pair. Production mechanism might explain the asymmetries [C31]. The decay rate of on mass shell π_L and σ_L (or η_L, ρ_L, \dots) is above the lower limit allowed by the detection in the measurement volume.
2. If the mass of e_{ex} is larger than $.9 \text{ MeV}$ but smaller than 1.3 MeV $e_{ex}\bar{e}$ decays dominate over e^+e^- decays. The decay $e_{ex} \rightarrow e + \gamma$ before detection could explain the observed energy asymmetry.
3. It will be found that the direct production of $e_{ex}\bar{e}$ pairs is also possible in the heavy ion collision but the rate is much smaller due to the smaller phase space volume in two-particle case. The annihilation rate of \bar{e}_{ex} in matter is by a factor $\sin^2(\theta_e)$ smaller than the annihilation rate of positron. This provides a unique signature of e_{ex} if e^+ annihilation rate in matter is larger than the decay rate of \bar{e}_{ex} . In lead the lifetime of positron is $\tau \sim 10^{-10} \text{ sec}$ and indeed larger than e_{ex} lifetime.

2.3.7 Karmen anomaly

A brief comment on the Karmen anomaly [C20] observed in the decays of π^+ is in order. The anomaly suggests the existence [C7] of new weakly interacting neutral particle x , which mixes with muon neutrino. Since $g = 1$ neutrino corresponds to charmed quark in hadron physics context having $k = 103$ rather than $k = 107$ as primary condensation level, a natural guess for its primary condensation level is $k = 113$, which would mean that the mass scale would be of order muon mass: the particle candidate indeed has mass of order 30 MeV. One class of solutions to laboratory constraints, which might evade also cosmological and astrophysical constraints, corresponds to object x mixing with muon type neutrino and decaying radiatively to $\gamma + \nu_\mu$ via the emission of virtual W boson. The value of the mixing parameter $U(\mu, x)$ describing $\nu_{mu} - x$ mixing satisfies $|U_{\mu,x}|^4 \simeq .8 \cdot 10^{-10}$.

The following naive PCAC argument gives order of magnitude estimate for $|U(\mu, x)| \sim \sin(\theta_\mu)$. The value of $g(\mu, \mu)$ is by a factor $m(\mu)/m_e$ larger than $g(e, e)$. If the lepto-hadronic couplings $g(\mu_{ex}, \mu_{ex})$ and $g(e_{ex}, e_{ex})$ are of same order of magnitude then one has $\sin(\theta_\mu) \leq .02$ (3 lepto-pion states and Op anomaly equal to $Op = 5 \cdot 10^{-3}$): the lower bound is 6.5 times larger than the value .003 deduced in [C7]. The actual value could be considerably smaller since e_{ex} mass could be larger than 1.3 MeV by a factor of order 10.

2.4 Lepto-Pions And Weak Decays

The couplings of lepto-meson to electro-weak gauge bosons can be estimated using PCAC and CVC hypothesis [B1]. The effective $m_{\pi_L} - W$ vertex is the matrix element of electro-weak axial current between vacuum and charged lepto-meson state and can be deduced using same arguments as in the case of ordinary charged pion

$$\langle 0 | J_A^\alpha | \pi_L^- \rangle = K m(\pi_L) p^\alpha, \quad (2.19)$$

where K is some numerical factor and p^α denotes the momentum of lepto-pion. For neutral lepto-pion the same argument gives vanishing coupling to photon by the conservation of vector current. This has the important consequence that lepto-pion cannot be observed as resonance in e^+e^- annihilation in single photon channel. In two photon channel lepto-pion should appear as resonance. The effective interaction Lagrangian is the ‘‘instanton’’ density of electromagnetic field giving additional contribution to the divergence of the axial current and was used to derive a model for lepto-pion production in heavy ion collisions.

2.4.1 Lepto-hadrons and lepton decays

The lifetime of charged lepto-pion is from PCAC estimates larger than 10^{-10} seconds by the previous PCAC estimates. Therefore lepto-pions are practically stable particles and can appear in the final states of particle reactions. In particular, lepto-pion atoms are possible and by Bose statistics have the peculiar property that ground state can contain many lepto-pions.

Lepton decays $L \rightarrow \nu_\mu + H_L$, $L = e, \mu, \tau$ via emission of virtual W are kinematically allowed and an anomalous resonance peak in the neutrino energy spectrum at energy

$$E(\nu_L) = \frac{m(L)}{2} - \frac{m_H^2}{2m(L)}, \quad (2.20)$$

provides a unique test for the lepto-hadron hypothesis. If lepto-pion is too light electrons would decay to charged lepto-pions and neutrinos unless the condition $m(\pi_L) > m_e$ holds true.

The existence of a new decay channel for muon is an obvious danger to the lepto-hadron scenario: large changes in muon decay rate are not allowed.

Consider first the decay $\mu \rightarrow \nu_\mu + \pi_L$ where π_L is on mass shell lepto-pion. Lepto-pion has energy $\sim m(\mu)/2$ in muon rest system and is highly relativistic so that in the muon rest system the lifetime of lepto-pion is of order $\frac{m(\mu)}{2m(\pi_L)}\tau(\pi_L)$ and the average length traveled by lepto-pion before decay is of order 10^8 meters! This means that lepto-pion can be treated as stable particle. The presence of a new decay channel changes the lifetime of muon although the rate for events using $e\nu_e$ pair as signature is not changed. The effective $H_L - W$ vertex was deduced above. The rate for the decay via lepto-pion emission and its ratio to ordinary rate for muon decay are given by

$$\begin{aligned} \Gamma(\mu \rightarrow \nu_\mu + H_L) &= \frac{G^2 K^2}{2^5 \pi} m^4(\mu) m^2(H_L) \left(1 - \frac{m^2(H_L)}{m^2(\mu)}\right) \frac{(m^2(\mu) - m^2(H_L))}{(m^2(\mu) + m^2(H_L))}, \\ \frac{\Gamma(\mu \rightarrow \nu_\mu + H_L)}{\Gamma(\mu \rightarrow \nu_\mu + e + \bar{\nu}_e)} &= 6 \cdot (2\pi^4) K^2 \frac{m^2(H_L)}{m^2(\mu)} \frac{(m^2(\mu) - m^2(H_L))}{(m^2(\mu) + m^2(H_L))}, \end{aligned} \quad (2.21)$$

and is of order $.93K^2$ in case of lepto-pion. As far as the determination of G_F or equivalently m_W^2 from muon decay rate is considered the situation seems to be good since the change introduced to G_F is of order $\Delta G_F/G_F \simeq 0.93K^2$ so that K must be considerably smaller than one. For the physical value of K : $K \leq 10^{-2}$ the contribution to the muon decay rate is negligible.

Lepto-hadrons can appear also as virtual particles in the decay amplitude $\mu \rightarrow \nu_\mu + e\nu_e$ and this changes the value of muon decay rate. The correction is however extremely small since the decay vertex of intermediate off mass shell lepto-pion is proportional to its decay rate.

2.4.2 Lepto-pions and beta decay

If lepto-pions are allowed as final state particles lepto-pion emission provides a new channel $n \rightarrow p + \pi_L$ for beta decay of nuclei since the invariant mass of virtual W boson varies within the range ($m_e = 0.511 \text{ MeV}$, $m_n - m_p = 1.293 \text{ MeV}$). The resonance peak for $m(\pi_L) \simeq 1 \text{ MeV}$ is extremely sharp due to the long lifetime of the charged lepto-pion. The energy of the lepto-pion at resonance is

$$E(\pi_L) = (m_n - m_p) \frac{(m_n + m_p)}{2m_n} + \frac{m(\pi_L)^2}{2m_n} \simeq m_n - m_p. \quad (2.22)$$

Together with long lifetime this lepto-pions escape the detector volume without decaying (the exact knowledge of the energy of charged lepto-pion might make possible its direct detection).

The contribution of lepto-pion to neutron decay rate is not negligible. Decay amplitude is proportional to superposition of divergences of axial and vector currents between proton and neutron states.

$$M = \frac{G}{\sqrt{2}} K m(\pi_L) (q^\alpha V_\alpha + q^\alpha A_\alpha) . \quad (2.23)$$

For exactly conserved vector current the contribution of vector current vanishes identically. The matrix element of the divergence of axial vector current at small momentum transfer (approximately zero) is in good approximation given by

$$\begin{aligned} \langle p | q^\alpha A_\alpha | n \rangle &= g_A (m_p + m_n) \bar{u}_p \gamma_5 u_n , \\ g_A &\simeq 1.253 . \end{aligned} \quad (2.24)$$

Straightforward calculation shows that the ratio for the decay rate via lepto-pion emission and ordinary beta decay rate is in good approximation given by

$$\begin{aligned} \frac{\Gamma(n \rightarrow p + \pi_L)}{\Gamma(n \rightarrow p + e + \bar{\nu}_e)} &= \frac{30\pi^2 g_A^2 K^2}{0.47 \cdot (1 + 3g_A^2)} \frac{m_{\pi_L}^2 (\Delta^2 - m_{\pi_L}^2)^2}{\Delta^6} , \\ \Delta &= m(n) - m(p) . \end{aligned} \quad (2.25)$$

Lepto-pion contribution is smaller than ordinary contribution if the condition

$$K < \left[\frac{.47 \cdot (1 + 3g_A^2)}{30\pi^2 g_A^2} \frac{\Delta^6}{(\Delta^2 - m_{\pi_L}^2)^2 m_{\pi_L}^2} \right]^{1/2} \simeq .28 , \quad (2.26)$$

is satisfied. The upper bound $K \leq 10^{-2}$ coming from the lepto-pion decay width and Op anomaly implies that the contribution of the lepto-pion to beta decay rate is very small.

2.5 Ortopositronium Puzzle And Lepto-Pion In Photon Photon Scattering

The decay rate of ortopositronium (Op) has been found to be slightly larger than the rate predicted by QED [C35, C17]: the discrepancy is of order $\Delta\Gamma/\Gamma \sim 10^{-3}$. For parapositronium no anomaly has been observed. Most of the proposed explanations [C17] are based on the decay mode $Op \rightarrow X + \gamma$, where X is some exotic particle. The experimental limits on the branching ratio $\Gamma(Op \rightarrow X + \gamma)$ are below the required value of order 10^{-3} . This explanation is excluded also by the standard cosmology [C17].

Lepto-pion hypothesis suggests an obvious solution of the Op-puzzle. The increase in annihilation rate is due to the additional contribution to $Op \rightarrow 3\gamma$ decay coming from the decay $Op \rightarrow \gamma V$ (V denotes “virtual”) followed by the decay $\gamma V \rightarrow \gamma + \pi_L^V$ followed by the decay $\pi_L^V \rightarrow \gamma + \gamma$ of the virtual lepto-pion to two photon state. $\gamma\gamma\pi_L$ vertices are induced by the axial current anomaly $\propto E \cdot B$. Also a modification of parapositronium decay rate is predicted. The first contribution comes from the decay $Op \rightarrow \pi_L^V \rightarrow \gamma + \gamma$ but the contribution is very small due the smallness of the coupling $g(e, e)$. The second contribution obtained from ortopositronium contribution by replacing one outgoing photon with a loop photon is also small. Since the production of a real lepto-pion is impossible, the mechanism is consistent with the experimental constraints.

The modification to the Op annihilation amplitude comes in a good approximation from the interference term between the ordinary e^+e^- annihilation amplitude F_{st} and lepto-pion induced annihilation amplitude F_{new} :

$$\Delta\Gamma \propto 2Re(F_{st}\bar{F}_{new}) , \quad (2.27)$$

and rough order of magnitude estimate suggests $\Delta\Gamma/\Gamma \sim K^2/e^2 = \alpha^2/4\pi \sim 10^{-3}$. It turns out that the sign and the order of magnitude of the new contribution are correct for $f(\pi_L) \sim 2 \text{ keV}$ deduced also from the anomalous e^+e^- production rate.

The new contribution to $e^+e^- \rightarrow 3\gamma$ decay amplitude is most easily derivable using for lepto-pion-photon interaction the effective action

$$\begin{aligned} L_1 &= K\pi_L F \wedge F , \\ K &= \frac{\alpha_{em}N_c}{8\pi f(\pi_L)} , \end{aligned} \quad (2.28)$$

where F is quantized electromagnetic field. The calculation of the lepto-pion contribution proceeds in manner described in [B1], where the expression for the standard contribution and an elegant method for treating the average over e^+e^- spin triplet states and sum over photon polarizations, can be found. The contribution to the decay rate can be written as

$$\begin{aligned} \frac{\Delta\Gamma}{\Gamma} &\simeq K_1 I_0 , \\ K_1 &= \frac{3\alpha N_c^2}{(\pi^2 - 9)2^9(2\pi)^3} \left(\frac{m_e}{f(\pi_L)}\right)^2 , \\ I_0 &= \int_0^1 \int_{-1}^{umax} \frac{f}{v+f-1-x^2} v^2 (2(f-v)u + 2 - v - f) dv du , \\ f &\equiv f(v, u) = 1 - \frac{v}{2} - \sqrt{\left(1 - \frac{v}{2}\right)^2 - \frac{1-v}{1-u}} , \\ u &= \bar{n}_1 \cdot \bar{n}_2 , \quad \bar{n}_i = \frac{\bar{k}_i}{\omega_i} , \quad umax = \frac{\left(\frac{v}{2}\right)^2}{\left(1 - \frac{v}{2}\right)^2} , \\ v &= \frac{\omega_3}{m_e} , \quad x = \frac{m_{\pi_L}}{2m_e} . \end{aligned} \quad (2.29)$$

ω_i and \bar{k}_i denote the energies of photons, u denotes the cosine of the angle between first and second photon and v is the energy of the third photon using electron mass as unit. The condition $\Delta\Gamma/\Gamma = 10^{-3}$ gives for the parameter $f(\pi_L)$ the value $f(\pi_L)(1.062 \text{ MeV}) \simeq N_c \cdot 7.9 \text{ keV}$. If there are several lepto-pion states, they contribute to the decay anomaly additively. If the four known resonances correspond directly to lepto-pions decaying to lepto-nucleon pairs and $f(\pi_L)$ is assumed to scale as $N_c m_{\pi_L}$, one obtains $f(\pi_L)(1.062 \text{ MeV}) \simeq N_c \cdot 14.7 \text{ keV}$. From the PCAC relation one obtains for $\sin(\theta_e)$ the upper bound $\sin(\theta_e) \leq x \cdot \sqrt{N_c} 10^{-4}$ assuming $m_{e\pi} \geq 1.3 \text{ MeV}$ (so that $e_{e\pi}\bar{e}$ decay is not possible), where $x = 1.2$ for single lepto-pion state and $x = 1.36$ for four lepto-pion states identified as the observed resonances.

Lepto-pion photon interaction implies also a new contribution to photon-photon scattering. Just at the threshold $E = m_{\pi_L}/2$ the creation of lepto-pion in photon photon scattering is possible and the appearance of lepto-pion as virtual particle gives resonance type behaviour to photon photon scattering near $s = m_{\pi_L}^2$. The total photon-photon cross section in zero decay width approximation is given by

$$\sigma = \frac{\alpha^4 N_c^2}{2^{14}(2\pi)^6} \frac{E^6}{f_{\pi_L}^4 \left(E^2 - \frac{m_{\pi_L}^2}{4}\right)^2} . \quad (2.30)$$

2.6 Spontaneous Vacuum Expectation Of Lepto-Pion Field As Source Of Lepto-Pions

The basic assumption in the model of lepto-pion and lepto-hadron production is the spontaneous generation of lepto-pion vacuum expectation value in strong nonorthogonal electric and magnetic fields. This assumption is in fact very natural in TGD ¹.

¹ "Instanton density" generates coherent state of lepto-pions just like classical em current generates coherent state of photons

Table 1: The dependence of various quantities on the number of lepto-pion type states and Op anomaly, whose value is varied assuming the proportionality $f(\pi_L) \propto N_c m_{\pi_L}$. N_c refers to the number of lepto-pion states in given representation and Op denotes lepto-pion anomaly.

N	$Op/10^{-3}$	$f(\pi_L)/(N_c keV)$	$\sin(\theta_e)(m_{ex}/1.3 MeV)^{1/2}$	$\Gamma(\pi_L)/keV$
1	1	7.9	$1.2 \cdot 10^{-4} \sqrt{N_c}$.51
3	1	14.7	$1.7 \cdot 10^{-4} \sqrt{N_c}$.13
3	5	6.5	$3.6 \cdot 10^{-4} \sqrt{N_c}$.73

1. The well known relation [B1] expressing pion field as a sum of the divergence of axial vector current and anomaly term generalizes to the case of lepto-pion

$$\pi_L = \frac{1}{f(\pi_L)m^2(\pi_L)} (\nabla \cdot j^A + \frac{\alpha_{em} N_c}{2\pi} E \cdot B) . \quad (2.31)$$

In the case of lepto-pion case the value of $f(\pi_L)$ has been already deduced from PCAC argument. Anomaly term gives rise to pion decay to two photons so that one obtains an estimate for the lifetime of the lepto-pion.

This relation is taken as the basis for the model describing also the production of lepto-pion in external electromagnetic field. The idea is that the presence of external electromagnetic field gives rise to a vacuum expectation value of lepto-pion field. Vacuum expectation is obtained by assuming that the vacuum expectation value of axial vector current vanishes.

$$\begin{aligned} \langle vac | \pi | vac \rangle &= KE \cdot B , \\ K &= \frac{\alpha_{em} N_c}{2\pi f(\pi_L)m^2(\pi_L)} . \end{aligned} \quad (2.32)$$

Some comments concerning this hypothesis are in order here:

- (a) The basic hypothesis making possible to avoid large parity breaking effects in atomic and molecular physics is that p-adic condensation levels with length scale $L(n) < 10^{-6} m$ are purely electromagnetic in the sense that nuclei feed their Z^0 charges on condensate levels with $L(n) \geq 10^{-6} m$. The absence of Z^0 charges does not however exclude the possibility of the classical Z^0 fields induced by the nonorthogonality of the ordinary electric and magnetic fields (if Z^0 fields vanish E and B are orthogonal in TGD).
 - (b) The non-vanishing vacuum expectation value of the lepto-pion field implies parity breaking in atomic length scales. This is understandable from basic principles of TGD since classical Z^0 field has parity breaking axial coupling to electrons and protons. The non-vanishing classical lepto-pion field is in fact more or less equivalent with the presence of classical Z^0 field.
2. The amplitude for the production of lepto-pion with four momentum $p = (p_0, \vec{p})$ in an external electromagnetic field can be deduced by writing lepto-pion field as sum of classical and quantum parts: $\pi_L = \pi_L(class) + \pi_L(quant)$ and by decomposing the mass term into interaction term plus c-number term and standard mass term:

$$\begin{aligned} \frac{m^2(\pi_L)\pi_L^2}{2} &= L_{int} + L_0 , \\ L_0 &= \frac{m^2(\pi_L)}{2} (\pi_L^2(class) + \pi_L^2(quant)) , \\ L_{int} &= m^2(\pi_L)\pi_L(class)\pi_L(quant) . \end{aligned} \quad (2.33)$$

Interaction Lagrangian corresponds to L_{int} linear in lepto-pion oscillator operators. Using standard LSZ reduction formula and normalization conventions of [B1] one obtains for the probability amplitude for creating lepto-pion of momentum p from vacuum the expression

$$\begin{aligned} A(p) &\equiv \langle a(p)\pi_L \rangle = (2\pi)^3 m^2(\pi_L) \int f_p(x) \langle vac | \pi | vac \rangle d^4x , \\ f_p &= e^{ip \cdot x} . \end{aligned} \quad (2.34)$$

The probability for the production of lepto-pion in phase space volume element d^3p is obtained by multiplying with the density of states factor $d^3n = V \frac{d^3p}{(2\pi)^3}$:

$$\begin{aligned} dP &= A|U|^2 V d^3p , \\ A &= \left(\frac{\alpha_{em} N_c^2 m^2(\pi_L)}{2\pi f(\pi_L)} \right)^2 , \\ U &= \int e^{ip \cdot x} E \cdot B d^4x . \end{aligned} \quad (2.35)$$

The first conclusion that one can draw is that nonstatic electromagnetic fields are required for lepto-pion creation since in static fields energy conservation forces lepto-pion to have zero energy and thus prohibits real lepto-pion production. In particular, the spontaneous creation lepto-pion in static Coulombic and magnetic dipole fields of nucleus is impossible.

2.7 Sigma Model And Creation Of Lepto-Hadrons In Electromagnetic Fields

2.7.1 Why sigma model approach?

For several reasons it is necessary to generalize the model for lepto-pion production to a model for lepto-hadron production.

1. Sigma model approach is necessary if one assumes that anomalous e^+e^- pairs are genuine e^+e^- pairs rather lepto-nucleon pairs produced in the decays of lepto-sigmas.
2. A model for the production of lepto-hadrons is obtained from an effective action describing the strong and electromagnetic interactions between lepto-hadrons. The simplest model is sigma model describing the interaction between lepto-nucleons, lepto-pion and a hypothetical scalar particle σ_L [B1]. This model realizes lepto-pion field as a divergence of the axial current and gives the standard relation between $f(\pi_L)$, g and m_{ex} . All couplings of the model are related to the masses of e_{ex} , π_L and σ_L . The generation of lepto-pion vacuum expectation value in the proposed manner takes place via triangle anomaly diagrams in the external electromagnetic field.
3. If needed the model can be generalized to contain terms describing also other lepto-hadrons. The generalized model should contain also vector bosons ρ_L and ω_L as well as pseudo-scalars η_L and η'_L and radial excitations of π_L and σ_L . An open question is whether also η and η' generate vacuum expectation value proportional to $E \cdot B$. Actually all these states appear as 3-fold degenerate for the minimal color representation content of the theory.
4. The following observations are useful for what follows.
 - (a) Ortopositronium decay width anomaly gives the estimate $f(\pi_L) \sim N_c \cdot 7.9 \text{ keV}$ and from this one can deduce an upper bound for lepto-pion production cross section in an external electromagnetic field. The calculation of lepto-pion production cross section shows that lepto-pion production cross section is somewhat smaller than the upper bound for the observed anomalous e^+e^- production cross section, even when one tunes the values of the various parameters. This is consistent with the idea that lepto-nucleon pairs, with lepto-nucleon mass being only slightly larger than electron mass, are in question.

- (b) Also the direct production of the lepto-nucleon pairs via the interaction term $g\cos(\theta_e)\bar{e}_{ex}\gamma_5 e_{ex}\pi_L(cl)$ is possible but gives rise to continuum mass squared spectrum rather than resonant structures. The direct production of the pairs via the interaction term $gsin(\theta_e)\bar{e}\gamma_5 e_{ex}\pi_L(cl)$ from is much slower process than the production via the meson decays and does not give rise to resonant structures since Also the production via the $\bar{e}e_{ex}$ decay of virtual lepto-pion created from classical field is slow process since it involves $sin^2(\theta_e)$.
- (c) e^+e^- production can also proceed also via the creation of many particle states. The simplest candidates are $V_L + \pi_L$ states created via $\partial_\alpha\pi_L V^\alpha\pi_L(class)$ term in action and $\sigma_L + \pi_L$ states created via the the $k\sigma_L\pi_L\pi_L(class)$ term in the sigma model action. The production cross section via the decays of vector mesons is certainly very small since the production vertex involves the inner product of vector boson 3 momentum with its polarization vector and the situation is non-relativistic.
- (d) If the strong decay of σ_L to lepto-mesons is kinematically forbidden (this is not suggested by the experience with the ordinary hadron physics), the production rate for σ_L meson is large since the coupling k turns out to be given by $k = (m_{\sigma_L}^2 - m_{\pi_L}^2)/2f(\pi_L)$ and is anomalously large for the value of $f(\pi_L) \geq 7.9 \cdot N_c \text{ keV}$ derived from orthopositronium anomaly: $k \sim 336m(\pi_L)/N_c$ for $f(\pi_L) \sim N_c \cdot 7.9 \text{ keV}$. The resulting additional factor in the production cross section compensates the reduction factor coming from two-particle phase space volume. Despite this the estimate for the production cross section of anomalous e^+e^- pairs is roughly by a factor $1/N_c^2$ smaller than the maximum experimental cross section. The radiative corrections are huge and should give the dominant contribution to the cross section. It is however questionable very the assumed small lepto-hadronic decay width and mass of σ_L is consistent with the extremely strong interactions of σ_L .

2.7.2 Simplest sigma model

A detailed description of the sigma model can be found in [B1] and it suffices to outline only the crucial features here.

1. The action of lepto-hadronic sigma model reads as

$$\begin{aligned}
 L &= L_S + c\sigma_L \ , \\
 L_S &= \bar{\psi}_L(i\gamma^k\partial_k + g(\sigma_L + i\pi_L \cdot \tau\gamma_5))\psi_L + \frac{1}{2}((\partial\pi_L)^2 + (\partial\sigma_L)^2) \\
 &\quad - \frac{\mu^2}{2}(\sigma_L^2 + \pi_L^2) - \frac{\lambda}{4}(\sigma_L^2 + \pi_L^2)^2 \ .
 \end{aligned} \tag{2.36}$$

π_L is isospin triplet and σ_L isospin singlet. ψ_L is isospin doublet with electro-weak quantum numbers of electron and neutrino (e_{ex} and ν_{ex}). The model allows $so(4)$ symmetry. Vector current is conserved but for $c \neq 0$ axial current generates divergence, which is proportional to pion field: $\partial^\alpha A_\alpha = -c\pi_L$.

2. The presence of the linear term implies that σ_L field generates vacuum expectation value $\langle 0|\sigma_L|0\rangle = v$. When the action is written in terms of new quantum field $\sigma'_L = \sigma_L - v$ one has

$$\begin{aligned}
 L &= \bar{\psi}_L(i\gamma^k\partial_k + m + g(\sigma'_L + i\pi_L \cdot \tau\gamma_5))\psi_L + \frac{1}{2}((\partial\pi_L)^2 + (\partial\sigma'_L)^2) \\
 &\quad - \frac{1}{2}m_{\sigma_L}^2(\sigma'_L)^2 - \frac{m_{\pi_L}^2}{2}\pi_L^2 \\
 &\quad - \lambda v\sigma'_L((\sigma'_L)^2 + \pi_L^2) - \frac{\lambda}{4}((\sigma'_L)^2 + \pi_L^2)^2 \ ,
 \end{aligned} \tag{2.37}$$

The masses are given by

$$\begin{aligned} m_{\pi_L}^2 &= \mu^2 + \lambda v^2 , \\ m_{\sigma_L}^2 &= \mu^2 + 3\lambda v^2 , \\ m &= -gv . \end{aligned} \tag{2.38}$$

These formulas relate the parameters μ, v, g to lepto-hadrons masses.

3. The requirement that σ'_L has vanishing vacuum expectation implies in Born approximation

$$c - \mu^2 v - \lambda v^3 = 0 , \tag{2.39}$$

which implies

$$\begin{aligned} f(\pi_L) &= -v = -\frac{c}{m^2(\pi_L)} , \\ m_{e_x} &= gf(\pi_L) . \end{aligned} \tag{2.40}$$

Note that e_{e_x} and ν_{e_x} are predicted to have identical masses in this approximation. The value of the strong coupling constant g of lepto-hadronic physics is indeed strong from $m_{e_x} > m_e$ and $f(\pi_L) < N_c \cdot 10$ keV.

4. A new feature is the generation of the lepto-pion vacuum expectation value in an external electromagnetic field (of course, this is possible for the ordinary pion field, too!). The vacuum expectation is generated via the triangle anomaly diagram in a manner identical to the generation of a non-vanishing photon-photon decay amplitude and is proportional to the instanton density of the electromagnetic field. By redefining the pion field as a sum $\pi_L = \pi_L(cl) + \pi'_L$ one obtains effective action describing the creation of the lepto-hadrons in strong electromagnetic fields.
5. As far as the production of $\sigma_L \pi_L$ pairs is considered, the interaction term $\lambda v \sigma'_L \pi_L^2$ is especially interesting since it leads to the creation of $\sigma_L \pi_L$ pairs via the interaction term $k \lambda v \sigma'_L \pi_L(qu) \pi_L(cl)$.

The coefficient of this term can be expressed in terms of the lepto-meson masses and $f(\pi_L)$:

$$\begin{aligned} k &\equiv 2\lambda v = \frac{m_{\sigma_L}^2 - m_{\pi_L}^2}{2f(\pi_L)} = x m_{\pi_L} , \\ x &= \frac{1}{2} \left(\frac{m_{\sigma_L}^2}{m_{\pi_L}^2} - 1 \right) \frac{m_{\pi_L}}{f(\pi_L)} . \end{aligned} \tag{2.41}$$

The large value of the coupling deriving from $f(\pi_L) = N_c \cdot 7.9$ keV) compensates the reduction of the production rate coming from the smallness of two-particle phase space volume as compared with single particle-phase space volume but fails to produce large enough production cross section. The large value of $g(\sigma_L, \sigma_L, \sigma_L) = g(\sigma_L, \pi_L, \pi_L)$ however implies that the radiative contribution to the production cross section coming from the emission of a virtual sigma in the production vertex is much larger than the lowest order production cross section and with a rather small value of the relative $\sigma_L - \pi_L$ mass difference correct order of magnitude of cross section should be possible.

2.8 Classical Model For Lepto-Pion Production

The nice feature of both quantum and classical model is that the production amplitudes associated with all lepto-hadron production reactions in external electromagnetic field are proportional to the lepto-pion production amplitude and apart from phase space volume factors production cross sections are expected to be given by lepto-pion production cross section. Therefore it makes sense to construct a detailed model for lepto-pion production despite the fact that lepto-pion decays probably contribute only a very small fraction to the observed e^+e^- pairs.

2.8.1 General considerations

Angular momentum barrier makes the production of lepto-mesons with orbital angular momentum $L > 0$ improbable. Therefore the observed resonances are expected to be $L = 0$ pseudo-scalar states. Lepto-pion production has two signatures which any realistic model should reproduce.

1. Data are consistent with the assumption that states are produced at rest in cm frame.
2. The production probability has a peak in a narrow region of velocities of colliding nucleus around the velocity needed to overcome Coulomb barrier in head on collision. The relative width of the velocity peak is of order $\Delta\beta/\beta \simeq \cdot 10^{-2}$ [C24]. In Th-Th system [J2] two peaks at projectile energies 5.70 MeV and 5.75 MeV per nucleon have been observed. This suggests that some kind of diffraction mechanism based on the finite size of nuclei is at work. In this section a model treating nuclei as point like charges and nucleus-nucleus collision purely classically is developed. This model yields qualitative predictions in agreement with the signature 1) but fails to reproduce the possible diffraction behavior although one can develop argument for understanding the behavior above Coulomb wall.

The general expression for the amplitude for creation of lepto-pion in external electric and magnetic fields has been derived in Appendix. Let us now specialize to the case of heavy ion collision. We consider the situation, where the scattering angle of the colliding nucleus is measured. Treating the collision completely classically we can assume that collision occurs with a well defined value of the impact parameter in a fixed scattering plane. The coordinates are chosen so that target nucleus is at rest at the origin of the coordinates and colliding nucleus moves in z-direction in $y=0$ plane with velocity β . The scattering angle of the scattered nucleus is denoted by α , the velocity of the lepto-pion by v and the direction angles of lepto-pion velocity by (θ, ϕ) .

The minimum value of the impact parameter for the Coulomb collision of point like charges is given by the expression

$$\begin{aligned} b &= \frac{b_0 \cot(\alpha/2)}{2} , \\ b_0 &= \frac{2Z_1 Z_2 \alpha_{em}}{M_R \beta^2} , \end{aligned} \quad (2.42)$$

where b_0 is the expression for the distance of the closest approach in head on collision. M_R denotes the reduced mass of the nucleus-nucleus system.

To estimate the amplitude for lepto-pion production the following simplifying assumptions are made.

1. Nuclei can be treated as point like charges. This assumption is well motivated, when the impact parameter of the collision is larger than the critical impact parameter given by the sum of radii of the colliding nuclei:

$$b_{cr} = R_1 + R_2 . \quad (2.43)$$

For scattering angles that are sufficiently large the values of the impact parameter do not satisfy the above condition in the region of the velocity peak. p-Adic considerations lead to the conclusion that nuclear condensation level corresponds to prime $p \sim 2^k$, $k = 113$ (k

is prime). This suggest that nuclear radius should be replaced by the size $L(113)$ of the p-adic convergence cube associated with nucleus (see the chapter “TGD and Nuclear Physics”: $L(113) \sim 1.7 \cdot 10^{-14} m$ implies that cutoff radius is $b_{cr} \sim 2L(113) \sim 3.4 \cdot 10^{-14} m$.

2. Since the velocities are non-relativistic (about $0.12c$) one can treat the motion of the nuclei non-relativistically and the non-retarded electromagnetic fields associated with the exactly known classical orbits can be used. The use of classical orbit doesn't take into account recoil effect caused by lepto-pion production. Since the mass ratio of lepto-pion and the reduced mass of heavy nucleus system is of order 10^{-5} the recoil effect is however negligible.
3. The model simplifies considerably, when the orbit is idealized with a straight line with impact parameter determined from the condition expressing scattering angle in terms of the impact parameter. This approximation is certainly well founded for large values of impact parameter. For small values of impact parameter the situation is quite different and an interesting problem is whether the contributions of long range radiation fields created by accelerating nuclei in head-on collision could give large contribution to lepto-pion production rate. On the line connecting the nuclei the electric part of the radiation field created by first nucleus is indeed parallel to the magnetic part of the radiation field created by second nucleus. In this approximation the instanton density in the rest frame of the target nucleus is just the scalar product of the Coulombic electric field E of the target nucleus and of the magnetic field B of the colliding nucleus obtained by boosting it from the Coulomb field of nucleus at rest.

2.8.2 Expression of the classical cross section

First some kinematical notations. lepto-pion four-momentum in the rest system of target nucleus is given by the following expression

$$\begin{aligned} p &= (p_0, \vec{p}) = m\gamma_1(1, v\sin(\theta)\cos(\phi), v\sin(\theta)\sin(\phi), v\cos(\theta)) \ , \\ \gamma_1 &= 1/(1-v^2)^{1/2} \ . \end{aligned} \quad (2.44)$$

The velocity and Lorentz boost factor of the projectile nucleus are denoted by β and $\gamma = 1/\sqrt{1-\beta^2}$.

The double differential cross section in the classical model can be written as

$$\begin{aligned} d\sigma &= dP 2\pi b db \ , \\ dP &= K |A(b, p)|^2 d^3n \ , \text{ per } d^3n = V \frac{d^3p}{(2\pi)^3} \ , \\ K &= (Z_1 Z_2)^2 (\alpha_{em})^4 \times N_c^2 \left(\frac{m(\pi_L)}{f(\pi_L)}\right)^2 \frac{1}{2\pi^{13}} \ , \\ A(b, p) &= N_0 \frac{4\pi}{Z_1 Z_2 \alpha_{em}} \times U(b, p) \ , \\ U(b, p) &= \int e^{ip \cdot x} E \cdot B d^4x \ , \\ N_0 &= \frac{(2\pi)^7}{i} \ . \end{aligned} \quad (2.45)$$

where b denotes impact parameter. The formula generalizes the classical formula for the cross section of Coulomb scattering. In the calculation of the total cross section one must introduce some cutoff radii and the presence of the volume factor V brings in the cutoff volume explicitly (particle in the box description for lepto-pions). Obviously the cutoff length must be longer than lepto-pion Compton length. Normalization factor N_0 has been introduced in order to extract out large powers of 2π .

From this one obtains differential cross section as

$$\begin{aligned}
d\sigma &= P2\pi b db , \\
P &= \int K|A(b,p)|^2 V \frac{d^3p}{(2\pi)^3} , .
\end{aligned}
\tag{2.46}$$

The first objection is the need to explicitly introduce the reaction volume: this obviously breaks manifest Lorentz invariance. The cross section was estimated in the earlier version of the model [K1] and turned to be too small by several orders of magnitude. This inspired the idea that constructive interference for the production amplitudes for different values of impact parameter could increase the cross section.

2.9 Quantum Model For Lepto-Pion Production

There are good reasons for considering the quantum model. First, the lepto-pion production cross section is by several orders of magnitude too small in classical model. Secondly, in Th-Th collisions there are indications about the presence of two velocity peaks with separation $\delta\beta/\beta \sim 10^{-2}$ [C24] and this suggests that quantum mechanical diffraction effects might be in question. These effects could come from the upper and/or lower length scale cutoff and from the de-localization of the wave function of incoming nucleus.

The question is what quantum model means. The most natural thing is to start from Coulomb scattering and multiply Coulomb scattering amplitude for a given impact parameter value b with the amplitude for lepto-pion production. This because the classical differential cross section given by $2\pi b db$ in Coulomb scattering equals to the quantum cross section. One might however argue that on basis of $S = 1 + T$ decomposition of S-matrix the lowest order contribution to lepto-pion production in quantum situation corresponds to the absence of any scattering. The lepto-pion production amplitude is indeed non-vanishing also for the free motion of nuclei. The resolution of what looks like a paradox could come from many-sheeted space-time concept (see **Fig.** <http://tgdtheory.fi/appfigures/manysheeted.jpg> or **Fig.** 9 in the appendix of this book): if no scattering occurs, the space-time sheets representing colliding nuclei do not touch and all and there is no interference of em fields so that there is no lepto-pion production. It turns however that lowest order contribution indeed corresponds to the absence of scattering in the model that works.

2.9.1 Two possible approaches

One can imagine two approaches to the construction of the model for production amplitude in quantum case.

The first approach is based on eikonal approximation [B2]. Eikonal approximation applies at high energy limit when the scattering angle is small and one can approximate the orbit of the projectile with a straight orbit.

The expression for the scattering amplitude in eikonal approximation reads as

$$\begin{aligned}
f(\theta, \phi) &= \frac{k}{2\pi i} \int d^2b \exp(-ik \cdot b) \exp(i\xi(b)) - 1 , \\
\xi(b) &= \frac{-m}{k\hbar^2} \int_{z=-\infty}^{z=\infty} dz V(z, b) , \\
\frac{d\sigma}{d\Omega} &= |f^2| .
\end{aligned}
\tag{2.47}$$

as one expands the exponential in lowest in spherically symmetric potential order one obtains the

$$f(\theta, \phi) \simeq -\frac{m}{2\pi\hbar^2} \int J_0(k_T b) \xi(b) b db .
\tag{2.48}$$

The challenge is to find whether it is possible to generalize this expression so that it applies to the production of lepto-pions.

1. The simplest guess is that one should multiply the eikonal amplitude with the dimensionless amplitude $A(b)$:

$$\begin{aligned} f(\theta, \phi) &\rightarrow f(\theta, \phi, p) = \frac{k}{2\pi i} \int d^2b \exp(-ik \cdot b) \exp(i\xi(b) - 1) A(b, p) \\ &\simeq -\frac{m}{2\pi\hbar^2} \int J_0(k_T b) \xi(b) A(b, p) b db . \end{aligned} \quad (2.49)$$

2. Amplitude squared must give differential cross section for lepto-pion production and scattering

$$\begin{aligned} d\sigma &= |f(\theta, \phi, p)|^2 d\Omega d^3n , \\ d^3n &= V d^3p . \end{aligned} \quad (2.50)$$

This requires an explicit introduction of a volume factor V via a spatial cutoff. This cutoff is necessary for the coordinate z in the case of Coulomb potential, and would have interpretation in terms of a finite spatio-temporal volume in which the space-time sheets of the colliding particles are in contact and fields interfere.

3. There are several objections against this approach. The loss of a manifest relativistic invariance in the density of states factor for lepto-pion does not look nice. One must keep count about the scattering of the projectile which means a considerable complication from the point of view of numerical calculations. In classical picture for orbits the scattering angle in principle is fixed once impact parameter is known so that the introduction of scattering angles does not look logical.

Second approach starts from the classical picture in which each impact parameter corresponds to a definite scattering angle so that the resulting amplitude describes lepto-pion production amplitude and says nothing about the scattering of the projectile. This approach is more in spirit with TGD since classical physics is exact part of quantum TGD and classical orbit is absolutely real from the point of view of lepto-pion production amplitude.

1. The counterpart of the eikonal exponent has interpretation as the exponent of classical action associated with the Coulomb interaction

$$S(b) = \int_{\gamma} V ds \quad (2.51)$$

along the orbit γ of the particle, which can be taken also as a real classical orbit but will be approximated with rectilinear orbit in sequel.

2. The first guess for the production amplitude is

$$\begin{aligned} f(p) &= \int d^2b \exp(-i\Delta k(b) \cdot b) \exp\left[\frac{i}{\hbar} S(b)\right] A(b, p) \\ &= \int J_0(k_T(b)b) \left(1 + \frac{i}{\hbar} \int_{z=-a}^{z=a} dz V(z, b) + \dots\right) A(b, p) . \end{aligned} \quad (2.52)$$

Δk is the change of the momentum in the classical scattering and in the scattering plane. The cutoff $|z| \leq a$ in the longitudinal direction corresponds to a finite imbedding space volume inside which the space-time sheets of target and projectile are in contact.

3. The production amplitude is non-trivial even if the interaction potential vanishes being given by

$$f(p) = \int d^2b \exp(-ik \cdot b) A(b, p) = 2\pi i \int J_0(k_T(b)b) \times A(b, p) b db . \quad (2.53)$$

This formula can be seen as a generalization of quantum formula in the sense that incoherent integral over production probabilities at various values of b is replaced by an integral over production amplitude over b so that interference effects become possible.

4. This result could be seen as a problem. On basis of $S = 1 + iT$ decomposition corresponding to free motion and genuine interaction, one could argue that since the exponent of action corresponds to S , $A(p, b)$ vanishes when the space-time sheets are not in contact. The improved guess for the amplitude is

$$\begin{aligned} f(p) &= \int d^2b \exp(-ik \cdot b) \exp\left(\frac{i}{\hbar} S(b)\right) A(b, p) \\ &= \int J_0(k_T(b)b) \left(\frac{i}{\hbar} \int_{z=-a}^{z=a} V(z, b) + \dots\right) A(b, p) . \end{aligned} \quad (2.54)$$

This would mean that there would be no classical limit when coherence is assumed to be lost. At this stage one must keep mind open for both options.

5. The dimension of $f(p)$ is L^2/\hbar

$$d\sigma = |f(p)|^2 \frac{d^3p}{2E_p (2\pi)^3} . \quad (2.55)$$

has correct dimension. This model will be considered in sequel. The earlier work in [K1] was however based on the first option.

2.9.2 Production amplitude

The Fourier transform of $E \cdot B$ can be expressed as a convolution of Fourier transforms of E and B and the resulting expression for the amplitude reduces by residue calculus (see APPENDIX) to the following general form

$$\begin{aligned} A(b, p) &\equiv N_0 \times \frac{4\pi}{Z_1 Z_2 \alpha_{em}} \times U(b, p) = 2\pi i (CUT_1 + CUT_2) , \\ N_0 &= \frac{(2\pi)^7}{i} . \end{aligned} \quad (2.56)$$

where nuclear charges are such that Coulomb potential is $1/r$. The motivation for the strange looking notation is to extract all powers of 2π so that the resulting amplitudes contain only factors of order unity.

The contribution of the first cut for $\phi \in [0, \pi/2]$ is given by the expression

$$\begin{aligned}
CUT_1 &= D_1 \times \int_0^{\pi/2} \exp\left(-\frac{b}{b_0} \cos(\psi)\right) A_1 d\psi , \\
D_1 &= -\frac{1}{2} \frac{\sin(\phi)}{\sin(\theta)} , \quad b_0 = \frac{\hbar \beta \gamma}{m \gamma_1} , \\
A_1 &= \frac{A + iB \cos(\psi)}{\cos^2(\psi) + 2iC \cos(\psi) + D} , \\
A &= \sin(\theta) \cos(\phi) , \quad B = K , \\
C &= K \frac{\cos(\phi)}{\sin(\theta)} , \quad D = -\sin^2(\phi) - \frac{K^2}{\sin^2(\theta)} , \\
K &= \beta \gamma \left(1 - \frac{v_{cm}}{\beta} \cos(\theta)\right) , \quad v_{cm} = \frac{2v}{1+v^2} .
\end{aligned} \tag{2.57}$$

The definitions of the various kinematical variables are given in previous formulas. The notation is tailored to express the facts that A_1 is rational function of $\cos(\psi)$ and that integrand depends exponentially on the impact parameter.

The expression for CUT_2 reads as

$$\begin{aligned}
CUT_2 &= D_2 \times \int_0^{\pi/2} \exp\left(i \frac{b}{b_1} \cos(\psi)\right) A_2 d\psi , \\
D_2 &= -\frac{\sin(\frac{\phi}{2})}{u \sin(\theta)} \times \exp\left(-\frac{b}{b_2}\right) , \\
b_1 &= \frac{\hbar \beta}{m \gamma_1} , \quad b_2 = \frac{\hbar}{mb \gamma_1 \times \sin(\theta) \cos(\phi)} \\
A_2 &= \frac{A \cos(\psi) + B}{\cos^2(\psi) + C \cos(\psi) + D} , \\
A &= \sin(\theta) \cos(\phi) u , \quad B = \frac{w}{v_{cm}} + \frac{v}{\beta} \sin^2(\theta) [\sin^2(\phi) - \cos^2(\phi)] , \\
C &= 2i \frac{\beta w \cos(\phi)}{uv_{cm} \sin(\theta)} , \quad D = -\frac{1}{u^2} \left(\frac{\sin^2(\phi)}{\gamma^2} + \beta^2 (v^2 \sin^2(\theta) - \frac{2vw}{v_{cm}}) \cos^2(\phi) \right) \\
&\quad + \frac{w^2}{v_{cm}^2 u^2 \sin^2(\theta)} + 2i \frac{\beta v}{u} \sin(\theta) \cos(\phi) , \\
u &= 1 - \beta v \cos(\theta) , \quad w = 1 - \frac{v_{cm}}{\beta} \cos(\theta) .
\end{aligned} \tag{2.58}$$

$$\tag{2.59}$$

The denominator X_2 has no poles and the contribution of the second cut is therefore always finite. Again the expression is tailored to make clear the functional dependence of the integrand on $\cos(\psi)$ and on impact parameter. Besides this the exponential damping makes in non-relativistic situation the integrand small everywhere except in the vicinity of $\cos(\Psi) = 0$ and for small values of the impact parameter.

Using the symmetries

$$\begin{aligned}
U(b, p_x, -p_y) &= -U(b, p_x, p_y) , \\
U(b, -p_x, -p_y) &= \bar{U}(b, p_x, p_y) ,
\end{aligned} \tag{2.60}$$

of the amplitude one can calculate the amplitude for other values of ϕ .

CUT_1 gives the singular contribution to the amplitude. The reason is that the factor X_1 appearing in denominator of cut term vanishes, when the conditions

$$\begin{aligned} \cos(\theta) &= \frac{\beta}{v_{cm}} , \\ \sin(\phi) &= \cos(\psi) , \end{aligned} \quad (2.61)$$

are satisfied. In forward direction this condition tells that z- component of the lepto-pion momentum in velocity center of mass coordinate system vanishes. In laboratory this condition means that the lepto-pion moves in certain cone defined by the value of its velocity. The condition is possible to satisfy only above the threshold $v_{cm} \geq \beta$.

For $K = 0$ the integral reduces to the form

$$CUT_1 = \frac{1}{2} \cos(\phi) \sin(\phi) \lim_{\varepsilon \rightarrow 0} \int_0^{\pi/2} \frac{\exp(-\frac{\cos(\psi)}{\sin(\phi_0)}) d\psi}{(\sin^2(\phi) - \cos^2\psi + i\varepsilon)} . \quad (2.62)$$

One can estimate the singular part of the integral by replacing the exponent term with its value at the pole. The integral contains two parts: the first part is principal value integral and second part can be regarded as integral over a small semicircle going around the pole of integrand in upper half plane. The remaining integrations can be performed using elementary calculus and one obtains for the singular cut contribution the approximate expression

$$\begin{aligned} CUT_1 &\simeq e^{-(b/a)(\sin(\phi)/\sin(\phi_0))} \left(\frac{\ln(X)}{2} + \frac{i\pi}{2} \right) , \\ X &= \frac{((1+s)^{1/2} + (1-s)^{1/2})}{((1+s)^{1/2} - (1-s)^{1/2})} , \\ s &= \sin(\phi) , \\ \sin(\phi_0) &= \frac{\beta\gamma}{\gamma_1 m(\pi_L) a} . \end{aligned} \quad (2.63)$$

The principal value contribution to the amplitude diverges logarithmically for $\phi = 0$ and dominates over ‘‘pole’’ contribution for small values of ϕ . For finite values of impact parameter the amplitude decreases exponentially as a function of ϕ .

If the singular term appearing in CUT_1 indeed gives the dominant contribution to the lepto-pion production one can make some conclusions concerning the properties of the production amplitude. For given lepto-pion cm velocity v_{cm} the production associated with the singular peak is predicted to occur mainly in the cone $\cos(\theta) = \beta/v_{cm}$: in forward direction this corresponds to the vanishing of the z-component of the lepto-pion momentum in velocity center of mass frame. Since the values of $\sin(\theta)$ are of order .1 the transversal momentum is small and production occurs almost at rest in cm frame as observed. In addition, the singular production cross section is concentrated in the production plane ($\phi = 0$) due to the exponential dependence of the singular production amplitude on the angle ϕ and impact parameter and the presence of the logarithmic singularity. The observed lepto-pion velocities are in the range $\Delta v/v \simeq 0.2$ [C24] and this corresponds to the angular width $\Delta\theta \simeq 34$ degrees.

2.9.3 Differential cross section in the quantum model

There are two options to consider depending on whether one uses $\exp(iS)$ or $\exp(iS) - 1$ to define the production amplitude.

1. For the $\exp(iS)$ option the expression for the differential cross section reads in the lowest order as

$$\begin{aligned}
d\sigma &= K|f_B|^2 \frac{d^3p}{2E_p} , \\
f_B &\simeq i \int \exp(-i\Delta k \cdot r)(CUT_1 + CUT_2)bdbdzd\phi , \\
K &= (Z_1 Z_2)^2 \alpha_{em}^4 N_c^2 \left(\frac{m(\pi_L)}{f(\pi_L)}\right)^2 \frac{1}{(2\pi)^{15}} .
\end{aligned} \tag{2.64}$$

Here Δk is the momentum exchange in Coulomb scattering and a vector in the scattering plane so that the above described formula is obtained for the linear orbits.

- For the $\exp(iS) - 1$ option the differential production cross section for lepto-pion is in the lowest non-trivial approximation for the exponent of action S given by the expression

$$\begin{aligned}
d\sigma &= K|f_B|^2 \frac{d^3p}{2E_p} , \\
f_B &\simeq \int \exp(-i\Delta k \cdot r)V(z, b)(CUT_1 + CUT_2)bdbdzd\phi , \\
V(z, b) &= \frac{1}{r} , \\
K &= (Z_1 Z_2)^4 \alpha_{em}^6 N_c^2 \left(\frac{m(\pi_L)}{f(\pi_L)}\right)^2 \frac{1}{(2\pi)^{15}} .
\end{aligned} \tag{2.65}$$

Effectively the Coulomb potential is replaced with the product of the Coulomb potential and lepto-pion production amplitude $A(b, p)$. Since α_{em} is assumed to correspond to relate to its standard value by a scaling \hbar_0/\hbar factor.

- Coulomb potential brings in an additional $(Z_1 Z_2 \alpha_{em})^2$ factor to the differential cross section, which in the case of heavy ion scattering increases the contribution to the cross section by a factor of order 3×10^3 but reduces it by a factor of order 5×10^{-5} in the case of proton-antiproton scattering. The increase of \hbar expected to be forced by the requirement that perturbation theory is not lost however reduces the contribution from higher orders in V . It should be possible to distinguish between the two options on basis of these differences.

The scattering amplitude can be reduced to a simpler form by using the defining integral representation

$$J_0(x) = \frac{1}{2\pi} \int_0^{2\pi} \exp(-ix \sin(\phi)) d\phi$$

of Bessel functions.

- For $\exp(iS)$ option this gives

$$\begin{aligned}
f_B &= 2\pi i \int J_0(\Delta k b)(CUT_1 + CUT_2)bdb , \\
\Delta k &= 2k \sin\left(\frac{\alpha}{2}\right) , \quad k = M_R \beta , \\
M_R &\simeq A_R m_p , \quad A_R = \frac{A_1 A_2}{A_1 + A_2} ,
\end{aligned} \tag{2.66}$$

where the length scale cutoffs in various integrations are not written explicitly. The value of α can be deduced once the value of impact parameter is known in the case of the classical Coulomb scattering.

2. For $\exp(iS) - 1$ option one has

$$\begin{aligned} f_B &= 2\pi i \int F(b) J_0(\Delta kb) (CUT_1 + CUT_2) b db , \\ F(b \geq b_{cr}) &= 2 \int dz \frac{1}{\sqrt{z^2 + b^2}} = \ln\left(\frac{\sqrt{a^2 - b^2} + a}{b}\right) , \end{aligned} \quad (2.67)$$

Note that the factors K appearing in the different cross section are different in these two cases.

2.9.4 Calculation of the lepto-pion production amplitude in the quantum model

The details related to the calculation of the production amplitude can be found in appendix and it suffices to describe only the general treatment here. The production amplitude of the quantum model contains integrations over the impact parameter and angle parameter ψ associated with the cut. The integrands appearing in the definition of the contributions CUT_1 and CUT_2 to the scattering amplitude have simple exponential dependence on impact parameter. The function F appearing in the definition of the scattering amplitude is a rather slow varying function as compared to the Bessel function, which allows trigonometric approximation and for small values of scattering angle equals to its value at origin. This motivates the division of the impact parameter range into pieces so that F can be approximated with its mean value inside each piece so that integration over cutoff parameters can be performed exactly inside each piece.

In Appendix the explicit expansion in power series with respect to impact parameter is derived by assuming $J_0(k_T b) \simeq 1$ and $F(b) = F = \text{constant}$. These formulas can be easily generalized by assuming a piecewise constancy of these two functions. This means that the only the integration over the lepto-pion phase space must be carried out numerically.

CUT_1 becomes also singular at $\cos(\theta) = \beta/v_{cm}$, $\cos(\psi) = \sin(\phi)$. The singular contribution of the production amplitude can be extracted by putting $\cos(\psi) = \sin(\phi)$ in the arguments of the exponent functions appearing in the amplitude so that one obtains a rational function of $\cos(\psi)$ and $\sin(\psi)$ integrable analytically. The remaining nonsingular contribution can be integrated numerically.

2.9.5 Formula for the production cross section

In the case of heavy ion collisions the rectilinear motion is not an excellent approximation since the anomalous events are observed near Coulomb wall and $\beta \simeq .1$ holds true. Despite this this can be taken as a first approximation.

The expression for the differential cross section for lepto-pion production in heavy ion collisions is given by

$$d\sigma = KF^2 \left| \int (CUT_1 + CUT_2) b db \right|^2 \frac{d^3p}{2E} , \quad (2.68)$$

This expression and also the expressions of the integrals of CUT_1 and CUT_2 are calculated explicitly as powers series of the impact parameter in the Appendix.

1. For $\exp(iS)$ option one has

$$\begin{aligned} K &= (Z_1 Z_2)^2 \alpha_{em}^4 N_c^2 \left[\frac{m(\pi_L)}{f(\pi_L)} \right]^2 \frac{1}{(2\pi)^{13}} , \\ F &= 1 . \end{aligned} \quad (2.69)$$

2. For $\exp(iS) - 1$ option one has

$$\begin{aligned} K &= (Z_1 Z_2)^4 \alpha_{em}^6 N_c^2 \left[\frac{m(\pi_L)}{f(\pi_L)} \right]^2 \frac{1}{(2\pi)^{13}} , \\ F &= 2 \langle \ln \left(\frac{\sqrt{a^2 - b^2} + a}{b} \right) \rangle . \end{aligned} \quad (2.70)$$

In the approximation that F is constant the two lowest order predictions are related by a scaling factor

$$R = (Z_1 Z_2 \alpha_{em})^2 F^2 . \quad (2.71)$$

It is interesting to get a rough order of magnitude feeling about the situation assuming that the contributions of CUT_1 and CUT_2 are of order unity. For $Z_1 = Z_2 = 92$ and $m(\pi_L)/f(\pi_L) \simeq 1.5$ -as in the case of ordinary pion- one obtains following results. It must be emphasized that these estimates are extremely sensitive to the over all scaling of f_B and to the choice of the cutoff parameter a and cannot be taken too seriously.

1. From $\beta \simeq .1$ one has $b_0 \simeq .1/m(\pi_L)$. One can argue that the impact parameter cutoff $a = x b_0$ should satisfy $a \geq 1/m_{\pi_L}$ so that $x \geq 10$ should hold true.
2. For $\exp(iS) - 1$ option one has $K = 4.7 \times 10^{-6}$. From the classical model the allowed phase space volume is of order $\frac{1}{3} \Delta v^3 \sim 10^{-4}$. By using $a = m(\pi_L)$ as a cutoff and $m(\pi_L) \simeq 2m_e$ one obtains $\sigma \sim 4 \mu\text{b}$, which is of same order of magnitude as the experimental estimate $5 \mu\text{b}$.
3. For $\exp(iS)$ option one has $K = 1.2 \times 10^{-9}$ and the estimate for cross section is 1.1 nb for $a = 1/m(\pi_L)$. A correct order of magnitude is obtained by assuming $a = 5.5/m(\pi_L)$ and that a^4 scaling holds true. At larger values of impact parameter a^2 scaling sets on and would require $a \sim 30/m(\pi_L)$ which would correspond to $.36 \text{ \AA}$ and to atomic length scale. It is not possible to distinguish between the two options.
4. The singular contribution near to production plane at the cone $v_{cm} \cos(\theta) = \beta$ is expected to enhance the total cross section. The strong sensitivity of the cross section to the choice of the cutoff parameter allows to reproduce the experimental findings easily and it would be important to establish strong bounds on the value of the impact parameter.

2.9.6 Dominating contribution to production cross section and diffractive effects

Consider now the behavior of the dominating singular contribution to the production amplitude at the cone $\cos(\theta) = \beta/v_{cm}$ depending on b via the exponent factor. This amplitude factorizes into a product

$$\begin{aligned} f_{B,sing} &= K_0 a^2 B(\Delta k) A_{sing}(b, p) , \\ B(\Delta k) &= \int F(ax) J_0(\Delta k a x) \exp\left(-\frac{\sin(\phi)}{\sin(\phi_0)} x\right) x dx , \\ &\sim \sqrt{\frac{2}{\pi \Delta k a}} \int F(ax) \cos\left(\Delta k a x - \frac{\pi}{4}\right) \exp\left(-\frac{\sin(\phi)}{\sin(\phi_0)} x\right) \sqrt{x} dx , \\ x &= \frac{b}{a} . \end{aligned} \quad (2.72)$$

The factor $A_{sing}(b, p) \equiv (4\pi/(Z_1 Z_2 \alpha_{em})) U_{sing}(b, p)$ is the analytically calculable singular and dominating part of the lepto-pion production amplitude (see appendix) with the exponential factor excluded. The factor B is responsible for diffractive effects. The contribution of the peak to the total production cross section is of same order of magnitude as the classical production cross section.

At the peak $\phi \sim 0$ the contribution the exponent of the production amplitude is constant at this limit one obtains product of the Fourier transform of Coulomb potential with cutoffs with the production amplitude. One can calculate the Fourier transform of the Coulomb potential analytically to obtain

$$\begin{aligned} f_{B,sing} &\simeq 4\pi K_0 \frac{(\cos(\Delta ka) - \cos(\Delta kb_{cr}))}{\Delta k^2} CUT_1 \\ \Delta k &= 2\beta \sin\left(\frac{\alpha}{2}\right) . \end{aligned} \quad (2.73)$$

One obtains oscillatory behavior as a function of the collision velocity in fixed angle scattering and the period of oscillation depends on scattering angle and varies in wide limits.

The relationship between scattering angle α and impact parameter in Coulomb scattering translates the impact parameter cutoffs to the scattering angle cutoffs

$$\begin{aligned} a &= \frac{Z_1 Z_2 \alpha_{em}}{M_R \beta^2} \cot(\alpha(min)/2) , \\ b_{cr} &= \frac{Z_1 Z_2 \alpha_{em}}{M_R \beta^2} \cot(\alpha(max)/2) . \end{aligned} \quad (2.74)$$

This gives for the argument Δkb of the Bessel function at lower and upper cutoffs the approximate expressions

$$\begin{aligned} \Delta ka &\simeq \frac{2Z_1 Z_2 \alpha_{em}}{\beta} \sim \frac{124}{\beta} , \\ \Delta kb_{cr} &\simeq x_0 \frac{2Z_1 Z_2 \alpha_{em}}{\beta} \sim \frac{124x_0}{\beta} . \end{aligned} \quad (2.75)$$

The numerical values are for $Z_1 = Z_2 = 92$ (U-U collision). What is remarkable that the argument Δka at upper momentum cutoff does not depend at all on the value of the cutoff length. The resulting oscillation at minimum scattering angle is more rapid than allowed by the width of the observed peak: $\Delta\beta/\beta \sim 3 \cdot 10^{-3}$ instead of $\Delta\beta/\beta \sim 10^{-2}$: of course, the measured value need not correspond to minimum scattering angle. The oscillation associated with the lower cutoff comes from $\cos(2M_R b_{cr} \beta \sin(\alpha/2))$ and is slow for small scattering angles $\alpha < 1/A_R \sim 10^{-2}$. For $\alpha(max)$ the oscillation is rapid: $\delta\beta/\beta \sim 10^{-3}$.

In the total production cross section integrated over all scattering angles (or finite angular range) diffractive effects disappear. This might explain why the peak has not been observed in some experiments [C24].

2.9.7 Cutoff length scales

Consider next the constraints on the upper cutoff length scale.

1. The production amplitude turns out to decrease exponentially as a function of impact parameter b unless lepto-pion is produced in scattering plane. The contribution of lepto-pions produced in scattering plane however gives divergent contribution to the total cross section integrated over all impact parameter values and upper cutoff length scale a is necessary. If one considers scattering with scattering angle between specified limits this is of course not a problem of classical model.
2. Upper cutoff length scale must be longer than the Compton length of lepto-pion.
3. Upper cutoff length scale a should be certainly smaller than the interatomic distance. For partially ionized atoms a more stringent upper bound for a is the size r of atom defined as the distance above which atom looks essentially neutral: a rough extrapolation from hydrogen atom gives $r \sim a_0/Z^{1/3} \sim 1.5 \cdot 10^{-11} m$ (a_0 is Bohr radius of hydrogen atom). Therefore cutoff scale would be between Bohr radius $a_0/Z \sim .5 \cdot 10^{-12} m$ and r . In the recent case however atoms are completely ionized so that cutoff length scale can be longer. It turns out that 10 A reproduces the empirical estimate for the cross section correctly.

2.9.8 Numerical estimate for the electro-pion production cross section

The numerical estimate for the electro-pion production cross section (see **Fig. 1**) is carried out for thorium with ($Z = 90, A = 232$). The value of the collision velocity of the incoming nucleus in the rest frame of the second nucleus is taken as $\beta = .1$. From the width $\delta v/v = .2$ of velocity distribution in the same frame the upper bound $\gamma \leq 1 + \delta$, $\delta \simeq 2 \times 10^{-3}$ for the Lorentz boost factor of electro-pion in cm system is deduced. The cutoff is necessary because energy conservation is not coded to the structure of the model.




Figure 1: Differential cross section $\sin^2(\theta) \times \frac{d^2\sigma}{2Ed^3p}$ for τ -pion production for $\gamma_1 = 1.0319 \times 10^3$ in the rest system of antiproton for $\delta = 1.5$. $m(\pi_\tau)$ defines the unit of energy and nb is the unit for cross section. The ranges of θ and ϕ are $(0, \pi)$ and $(0, \pi/2)$.

As expected, the singular contribution from the cone $v_{cm}\cos(\theta) = \beta$, $v_{cm} = 2v/(1 + v^2)$ gives the dominating contribution to the cross section. This contribution is proportional to the value of b_{max}^2 at the limit $\phi = 0$. Cutoff radius is taken to be $b_{max} = 150 \times \gamma_{cm}\hbar/m(\pi_e) = 1.04$ A. The numerical estimate for the cross section using the parameter values listed comes out as $\sigma = 5.6 \mu\text{b}$ to be compared with the rough experimental estimate of about $5 \mu\text{b}$. The interpretation would be that the space-time sheet associated with colliding nuclei during the collision has this transversal size in cm system. At this space-time sheet the electric and magnetic fields of the nuclei interfere.

From this one can cautiously conclude that lepto-pion model is consistent with both electro-pion production and τ -pion production in proton antiproton collisions. One can of course criticize the large value of impact parameter and a good justification for 1 Angstrom should be found. One could also worry about the singular character of the amplitude making the integration of total cross section somewhat risky business using the rather meager numerical facilities available. The rigorous method to calculate the contribution near the singularity relies on stepwise halving of the increment $\Delta\theta$ as one approaches the singularity. The calculation gives essentially the same result as that with constant value of $\Delta\theta$. Hence it seems that one can trust on the result of calculation.

Figure 2. gives the differential production cross section for $\gamma_1 = 1.0319$. Obviously the differential cross section is strongly concentrated at the cone due to singularity of the production amplitude for fixed b .

Table 2: Table summarizes lepto-pion lifetime and the upper bounds for lepto-pion (and lepto-nucleon pair) production cross sections for the lightest lepto-pion. N refers to the number of lepto-pion states and $Op = \Delta\Gamma/\Gamma$ refers to orthopositronium decay anomaly. The values of upper cutoff length a are in units of $10^{-10} m$.

N	$Op/10^{-3}$	$\Gamma(\pi_L)/keV$	$\sigma(\pi_L)/\mu b$	$\sigma(\pi_L)/\mu b$
			$a = .01$	$a = .1$
1	1	.51	.13	1.4
3	1	.13	.04	.41
3	5	.73	.19	2.1

The important conclusion is that the same model can reproduce the value of production cross section for both electro-pions explaining the old electron-positron anomaly of heavy ion collisions and τ -pions explaining the CDF anomaly of proton-antiproton collisions at cm energy $\sqrt{s} = 1.96$ TeV (to be discussed later) with essentially same and rather reasonable assumptions (do not however forget the large maximal value of the impact parameter!).

In the case of electro-pions one must notice that depending on situation the final states are gamma pairs for the electron-pion with mass very nearly equal to electron mass. In the case of neutral tau-pion the strong decay to three p-adically scaled down versions of τ -pion proceeds faster or at least rate comparable to that for the decay to gamma pair. For higher mass variants of electro-pion for which there is evidence (for instance, one with mass 1.6 MeV) the final states are dominated by electron-positron pairs. This is true if the primary decay products are electro-baryons of form (say) $e_{ex} = e_8\nu_8\nu_{c,8}$ resulting via electro-strong decays instead of electrons and having slightly larger mass than electron. Otherwise the decay to gamma pair would dominate also the decays of higher mass states. A small magnetic moment type coupling between e, e_{ex} and electro-gluon field made possible by the color octet character of colored leptons induces the mixing of e and e_{ex} so that e_{ex} can transform to e by the emission of photon. The anomalous magnetic moment of electron poses restrictions on the color magnetic coupling.

2.9.9 $e_{ex}^+e_{ex}^-$ pairs from lepto-pions or e^+e^- pairs from lepto-sigmas?

If one assumes that anomalous e^+e^- pairs correspond to lepto-nucleon pairs, then lepto-pion production cross section gives a direct estimate for the production rate of e^+e^- pairs. The results of the table 3 show that in case of 1.8 MeV state, the predicted cross section is roughly by a factor 5 smaller than the experimental upper bound for the cross section. Since this lepto-pion state is rather massive, positron decay width allows smaller $f(\pi_L)$ in this case and the production cross section could be larger than the estimate used by the $1/f(\pi_L)^2$ proportionality of the cross section. Both the simplicity and predictive power of this option and the satisfactory agreement with the experimental data suggest that this option provides the most plausible explanation of the anomalous e^+e^- pairs.

Table 2.

If one assumes that anomalous e^+e^- pairs result from the decays of lepto-sigmas, the value of e^+e^- production cross section can be estimated as follows. e^+e^- pairs are produced from via the creation of $\sigma_L\pi_L$ pairs from vacuum and subsequent decay σ_L to e^+e^- pairs. The estimate for (or rather for the upper bound of) $\pi_L\sigma_L$ production cross section is obtained as

$$\begin{aligned}
\sigma(e^+e^-) &\simeq X\sigma(\pi_L) , \\
X &= \frac{V_2}{V_1} \left(\frac{km_{\sigma_L}}{m_{\pi_L}^2} \right)^2 , \\
\frac{V_2}{V_1} &= V_{rel} = \frac{v_{12}^3}{3(2\pi)^2} \sim 1.1 \cdot 10^{-5} , \\
\frac{k}{m_{pi_L}} &= \frac{(m_{\sigma}^2 - m_{\pi_L}^2)}{2m_{\pi_L}f(\pi_L)} .
\end{aligned} \tag{2.76}$$

Here V_2/V_1 of two-particle and single particle phase space volumes. V_2 is in good approximation the product $V_1(cm)V_1(rel)$ of single particle phase space volumes associated with cm coordinate and relative coordinate and one has $V_2/V_1 \sim V_{rel} = \frac{v_{12}^3}{3(2\pi)^2} \simeq 1.1 \cdot 10^{-5}$ if the maximum value of the relative velocity is $v_{12} \sim .1$.

Situation is partially saved by the anomalously large value of $\sigma_L\pi_L\pi_L$ coupling constant k appearing in the production vertex $k\sigma_L\pi_L\pi_L(class)$. Production cross section is very sensitive to the value of $f(\pi_L)$ and Op anomaly $\Delta\Gamma/\Gamma = 5 \cdot 10^{-3}$ gives upper bound $2 \mu b/N_c^2$ for $a = 10^{-11} m$, which is considerably smaller than the experimental upper bound $5 \mu b$. The huge value of the $g(\pi_L, \pi_L, \sigma_L)$ and $g(\sigma_L, \sigma_L, \sigma_L)$, however implies that radiative corrections to the cross section given by σ exchange are much larger than the lowest order contribution to the cross section! If this is the case then lepto-sigma option might survive but perturbative approach probably would not make sense. On the other hand, one could argue that sigma model action should be regarded as an effective action giving only tree diagrams so that radiative corrections cannot save the situation. There are also purely physical counter arguments against lepto-sigma option: hadronic physics experience suggests that the mass of lepto-sigma is much larger than lepto-pion mass so that lepto-sigma becomes very wide resonance decaying strongly and having negligibly small branching ratio to e^+e^- pairs.

It must be emphasized that the estimates are very rough (the replacement of the integral over the angle α with rough upper bound, estimate for the phase space volume, the values of cutoff radii, the neglect of the velocity dependence of the production cross section, the estimate for the minimum scattering angle, ...). Also the measured production cross section is subject to considerable uncertainties (even the issue whether or not anomalous pairs are produced is not yet completely settled!).

2.9.10 Summary

The usefulness of the modelling lepto-pion production is that the knowledge of lepto-pion production rate makes it possible to estimate also the production rates for other lepto-hadrons and even for many particle states consisting of lepto-hadrons using some effective action describing the strong interactions between lepto-hadrons. One can consider two basic models for lepto-pion production. The models contain no free parameters unless one regards cutoff length scales as such. Classical model predicts the singular production characteristics of lepto-pion. Quantum model predicts several velocity peaks at fixed scattering angle and the distance between the peaks of the production cross section depends sensitively on the value of the scattering angle. Production cross section depends sensitively on the value of the scattering angle for a fixed collision velocity. In both models the reduction of the lepto-pion production rate above Coulomb wall could be understood as a threshold effect: for the collisions with impact parameter smaller than two times nuclear radius, the production amplitude becomes very small since $E \cdot B$ is more or less random for these collisions in the interaction region. The effect is visible for fixed sufficiently large scattering angle only. The value of the anomalous e^+e^- production cross section is of nearly the observed order of magnitude provided that e^+e^- pairs are actually lepto-nucleon pairs originating from the decays of the lepto-pions. Alternative mechanism, in which anomalous pairs originate from the creation of $\sigma_L\pi_L$ pairs from vacuum followed by the decay $\sigma_L \rightarrow e^+e^-$ gives too small production cross section by a factor of order $1/N_c^2$ in lowest order calculation. This alternative works only provided that radiative corrections give the dominant contribution to the production rate of $\pi_L\sigma_L$ pairs as is the case if $\pi_L\sigma_L$ mass difference is of order ten per cent. The existence of at least three colored leptons and family replication provide the most plausible explanation the appearance of several peaks.

The proposed models are certainly over idealizations: in particular the approximation that nuclear motion is free motion fails for those values of the impact parameter, which are most important in the classical model. To improve the models one should calculate the Fourier transform of $E \cdot B$ using the fields of nuclei for classical orbits in Coulomb field rather than free motion. The second improvement is related to the more precise modelling of the situation at length scales below b_{cr} , where nuclei do not behave like point like charges. A peculiar feature of the model from the point of view of standard physics is the appearance of the classical electromagnetic fields associated with the classical orbits of the colliding nuclei in the definition of the quantum model. This is in spirit with Quantum TGD: Quantum TGD associates a unique space-time surface (classical history)

to a given 3-surface (counterpart of quantum state).

3 Further Developments

This section represents further developments of lepto-hadron model which have emerged during years after the first version of the model published in International Journal of Theoretical Physics.

3.1 How To Observe Leptonic Color?

The most obvious argument against lepto-hadrons is that their production via the decay of virtual photons to lepto-mesons has not been observed in hadronic collisions. The argument is wrong. Anomalously large production of low energy e^+e^- pairs [C19, C28, C21, C47] in hadronic collisions has been actually observed. The most natural source for photons and e^+e^- pairs are lepto-hadrons. There are two possibilities for the basic production mechanism.

1. Colored leptons result directly from the decay of hadronic gluons. Internal consistency excludes this alternative.
2. Colored leptons result from the decay of virtual photons. This hypothesis is in accordance with the general idea that the QCD: s associated with different condensate levels of p-adic topological condensate do not communicate. More precisely, in TGD framework leptons and quarks correspond to different chiralities of WCW spinor s : this implies that baryon and lepton numbers are conserved exactly and therefore the stability of proton. In particular, leptons and quarks correspond to different Kac Moody representations: important difference as compared with typical unified theory, where leptons and quarks share common multiplets of the unifying group. The special feature of TGD is that there are several gluons since it is possible to associate to each Kac-Moody representation gluons, which are “irreducible” in the sense that they couple only to a single Kac Moody representation. It is clear that if the physical gluons are “irreducible” the world separates into different Kac Moody representations having their own color interactions and communicating only via electro-weak and gravitational interactions. In particular, no strong interactions between leptons and hadrons occur. Since colored lepton corresponds to colored ground state of Kac-Moody representations the gluonic color coupling between ordinary lepton and colored lepton vanishes.

If this picture is correct then lepto-hadrons are produced only via the ordinary electro-weak interactions: at higher energies via the decay of virtual photon to colored lepton pair and at low energies via the emission of lepto-pion by photon. Consider next various manners to observe the effects of lepton color.

1. Resonance structure in the photon-photon scattering and energy near lepto-pion mass is a unique signature of lepto-pion.
2. The production of lepto-mesons in strong classical electromagnetic fields (of nuclei, for example) is one possibility. There are several important constraints for the production of lepto-pions in this kind of situation.
 - i) The scalar product $E \cdot B$ must be large. Faraway from the source region this scalar product tends to vanish: consider only Coulomb field.
 - ii) The region, where $E \cdot B$ has considerable size cannot be too small as compared with lepto-pion de Broglie wavelength (large when compared with the size of nuclei for example). If this condition doesn't hold true the plane wave appearing in Fourier amplitude is essentially constant spatially and since the fields are approximately static the Fourier component of $E \cdot B$ is expressible as a spatial divergence, which reduces to a surface integral over a surface faraway from the source region. Resulting amplitude is small since fields in faraway region have essentially vanishing $E \cdot B$.
 - iii) If fields are exactly static, then energy conservation prohibits lepto-hadron production.

3. Also the production of $e_{ex}^+e_{ex}^-$ and $e^+e_{ex}^-$ pairs in nuclear electromagnetic fields with non-vanishing $E \cdot B$ is possible either directly or as decay products of lepto-pions. In the direct production, the predicted cross section is small due to the presence of two-particle phase space factor. One signature of e_{ex}^- is emission line accompanying the decay $e_{ex}^- \rightarrow e^- + \gamma$. The collisions of nuclei in highly ionized (perhaps astrophysical) plasmas provide a possible source of leptobaryons.
4. The interaction of quantized em field with classical electromagnetic fields is one experimental arrangement to come into mind. The simplest arrangement consisting of linearly polarized photons with energy near lepto-pion mass plus constant classical em field does not however work. The direct production of $\pi_L - \gamma$ pairs in rapidly varying classical electromagnetic field with frequency near lepto-pion mass is perhaps a more realistic possibility. An interesting possibility is that violent collisions inside astrophysical objects could lead to gamma ray bursts via the production of pions and lepto-pions in rapidly varying classical E and B fields.
5. In the collisions of hadrons, virtual photon produced in collision can decay to lepto-hadrons, which in turn produce lepto-pions decaying to leptonucleon pairs. As already noticed, anomalous production of low energy e^+e^- pairs (actually leptonucleon pairs!) [C19] in hadronic collisions has been observed.
6. $e-\nu_e$ and $e-\bar{\nu}_e$ scattering at energies below one MeV provide a unique signature of lepto-pion. In $e-\bar{\nu}_e$ scattering π_L appears as resonance.
7. If leptonic color coupling strength has sufficiently small value in the energy range at which lepto-hadronic QCD exists, e^+e^- annihilation at energies above few MeV should produce colored pairs and lepto-hadronic counterparts of the hadron jets should be observed. The fact that nothing like this has been observed, suggests that lepto-hadronic coupling constant evolution does not allow the perturbative QCD phase.

3.2 New Experimental Evidence

After writing this chapter astrophysical support for the notion of lepto-pions has appeared. There is also experimental evidence for the existence of colored muons

3.2.1 Could lepto-hadrons correspond to dark matter?

The proposed identification of cosmic strings (in TGD sense) as the ultimate source of both visible and dark matter discussed in [K3] does not exclude the possibility that a considerable portion of topologically condensed cosmic strings have decayed to some light particles. In particular, this could be the situation in the galactic nuclei.

The idea that lepto-hadrons might have something to do with the dark matter has popped up now and then during the last decade but for some reason I have not taken it seriously. Situation changed towards the end of the year 2003. There exist now detailed maps of the dark matter in the center of galaxy and it has been found that the density of dark matter correlates strongly with the intensity of monochromatic photons with energy equal to the rest mass of electron [E3].

The only explanation for the radiation is that some yet unidentified particle of mass very nearly equal to $2m_e$ decays to an electron positron pair. Electron and positron are almost at rest and this implies a high rate for the annihilation to a pair of gamma rays. A natural identification for the particle in question would be as a lepto-pion (or rather, electro-pion). By their low mass lepto-pions, just like ordinary pions, would be produced in high abundance, in lepto-hadronic strong reactions and therefore the intensity of the monochromatic photons resulting in their decays would serve as a measure for the density of the lepto-hadronic matter. Also the presence of lepto-pionic condensates can be considered.

These findings force to take seriously the identification of the dark matter as lepto-hadrons. This is however not the only possibility. The TGD based model for tetra-neutrons discussed in [K14] is based on the hypothesis that mesons made of scaled down versions of quarks corresponding to Mersenne prime M_{127} (ordinary quarks correspond to $k = 107$) and having masses around one MeV could correspond to the color electric flux tubes binding the neutrons to form a tetra-neutron. The same force would be also relevant for the understanding of alpha particles.

There are also good theoretical arguments for why lepto-hadrons should be dark matter in the sense of having a non-standard value of Planck constant.

1. Since particles with different Planck constant correspond to different pages of the book like structure defining the generalization of the imbedding space, the decays of intermediate gauge bosons to colored excitations of leptons would not occur and would thus not contribute to their decay widths.
2. In the case of electro-pions the large value of the coupling parameter $Z_1 Z_2 \alpha_{em} > 1$ combined with the hypothesis that a phase transition increasing Planck constant occurs as perturbative QFT like description fails would predict that electro-pions represent dark matter. Indeed, the power series expansion of the $exp(iS)$ term might well fail to converge in this case since S is proportional to $Z_1 Z_2$. For τ -pion production one has $Z_1 = -Z_2 = 1$ and in this case one can consider also the possibility that τ -pions are not dark in the sense of having large Planck constant. Contrary to the original expectations darkness does not affect the lowest order prediction for the production cross section of lepto-pion.

The proposed identification raises several questions.

1. Why the ratio of the lepto-hadronic mass density to the mass density of the ordinary hadrons would be so high, of order 7? Could an entire hierarchy of asymptotically non-free QCDs be responsible for the dark matter so that lepto-hadrons would explain only a small portion of the dark matter?
2. Under what conditions one can regard lepto-hadronic matter as a dark matter? Could short life-times of lepto-hadrons make them effectively dark matter in the sense that there would be no stable enough atom like structures consisting of say charged leptobaryons bound electromagnetically to the ordinary nuclei or electrons? But what would be the mechanism producing lepto-hadrons in this case (nuclear collisions produce lepto-pions only under very special conditions)?
3. What would be the role of the many-sheeted space-time: could lepto-hadrons and atomic nuclei reside at different space-time sheets so that leptobaryons could be long-lived? Could dark matter quite generally correspond to the matter at different space-time sheets and thus serve as a direct signature of the many-sheeted space-time topology?

3.2.2 Lightnings and lepto-pions

The latest discovery of Fermi space-telescope [C8] is the finding of .511 MeV gamma rays in the spectrum of photons associated with lightnings. It was discovered already years ago that lightnings are accompanied by X-rays [C43] and even gamma rays [C42]. For instance, the strong electric fields created by a positively charged region of cloud could accelerate electron from both downwards and upwards to this region. The problem is that atmosphere is not empty and dissipation would restrict the energies to be much lower than gamma ray energies which are in MeV range. Note that the temperatures in lightning are about 3×10^4 K and correspond to electron energy of 2.6 eV which is by a factor 10^5 smaller than electron mass and gamma ray energy scale!

Situation changes if dissipation is absent so that the electrons are accelerated without any energy losses. This is the case if the electrons reside in large \hbar quantum phase at magnetic flux tubes so that dissipative losses are small and electrons can reach relativistic energies. This is the explanation that I provided years ago for the [K4].

Fermi however observed also something completely new. There is also a peaking of gamma rays around energy .511 MeV. The decay of electro-pion is an obvious explanation for this peaking. If electro-pions are there, collisions of highly energetic particles lasting for time of about $\tau \sim \hbar/\text{MeV}$ are expected. The natural candidates for the colliding charged particles are electrons. The center of mass system -the system in which total momentum of colliding electron pair vanishes- should be in a good approximation at rest with respect to Fermi space telescope. Otherwise the energy of gamma rays would be higher or lower than .511 MeV.

The only possibility that I can imagine is that the second electron comes from below and second from above the positively charged region of the thunder cloud. Both arrive as dark electrons with

a large value of \hbar and are accelerated to relativistic energies since dissipation is very small. They could collide as dark electrons (the more probable option as will be found below) or suffer a phase transition transforming them to ordinary electrons before the collision. Electro-pion coherent state is created in the strong $E \cdot B$ created for a period of time of order $\tau \sim \hbar_0/\text{MeV}$. This state annihilates rapidly to pairs of gamma rays which are ordinary or transform to ordinary ones depending on whether electrons were dark or not.

What the phase transition of dark electrons to ordinary electrons means, needs some explaining. The generalized imbedding space is obtained by gluing almost copies of 8-D imbedding space $M^4 \times CP_2$ along their common back to get a book like structure. Particles at different pages of the book are dark with respect to each other in the sense that they have no local interactions. This is enough to explain what is actually known about dark matter. Particles at different pages can however interact via classical fields and photon exchange (for instance). The phase transition of electron from dark to visible form preceding the collision of dark electrons would simply mean the leakage from large \hbar page to the “visible” page with ordinary value of Planck constant.

Alert reader might be ready to ask the obvious question. Why not to test the hypothesis in laboratory? It should not be too difficult to allow two electrons to collide with a relativistic energy and find whether gamma pairs with energy .511 MeV are produced in rest system. Maybe gamma ray pairs have been missed for some reason? If not (the probable option), then colored electrons and lepto-pions are always dark. This would explain why the colored leptons do not contribute to the decay widths of weak gauge bosons which pose very strong constraints for the existence of light exotic particles.

3.2.3 Lightnings, dark matter, and lepto-pion hypothesis again

Lightnings have been found to involve phenomena difficult to understand in the framework of standard physics. Very high energy photons, even gamma rays and electrons and positrons with energies in gamma energy range, have been observed.

I learned recently about even more mysterious looking discovery (see <http://tinyurl.com/jucwhod>). Physicist Joseph Dwyer from University of New Hampshire and lightning scientists from the University of California at Santa Cruz and Florida Tech describe this discovery in a paper to be published in the Journal of Plasma Physics. In August 2009, Dwyer and colleagues were aboard a National Center for Atmospheric Research Gulfstream V when it inadvertently flew into the extremely violent thunderstorm, it turned out, through a large cloud of positrons, the antimatter opposite of electrons, that should not have been there. One would have expected that positrons would have been produced by annihilation of highly energetic gamma rays with energy above .5 MeV but no gamma rays were detected.

This looks rather mysterious from standard physics point of view. There are also earlier strange discoveries

1. Lightning strikes release powerful X-ray bursts [C43] (see “Lightning strikes release powerful X-ray bursts” at tinyurl.com/zqc7r7z).
2. Also high energy gamma rays and electrons accompany lightnings [C42] (see “Earth creates powerful gamma-ray flashes” at <http://tinyurl.com/juy8uj8>). The problem is that electrons should lose their energy while traversing through the atmosphere so that energies in even X ray range would be impossible.
3. The third strange discovery was made with Fermi telescope [C8] (see “Antimatter from lightning flashes the Fermi space telescope” at <http://tinyurl.com/p2z3n9p>): gamma rays with energies .511 MeV (electron mass) accompany lightnings as if something with mass of 2 electron masses would decay to gamma pairs.

Could TGD explain these findings.

1. A possible explanation for the finding of Fermi telescope is that in the strong magnetic field of colliding very high energy colliding electrons assignable to the dark magnetic flux tubes of Earth particles that I call electropions suggested by TGD are created [K16] (see <http://tinyurl.com/zvk3umn>). Also evidence for mu-pions and tau-pions exists. They

would have mass rather precisely 2 times the mass of electron and would be bound states of color excited electron and positron. Evidence for this kind of states was found already at seventies in heavy ion collisions around Coulomb wall producing electron positron pairs at total energy of 2 times electron mass but since they do not fit at all to the standard physics picture (too large decay width for weak bosons would be predicted) they have been put under the rug, so to say. The paradox is solved if these particles are dark in TGD sense.

2. If the annihilations of electropions give rise to dark electron-positron pairs and dark gamma rays, which then transform to ordinary particles, one could understand the absence of gamma rays in the situation described by Dwyer et al in terms of too slow transformation to ordinary particles. For instance, the strong electric fields created by a positively charged region of cloud could accelerate electron from both downwards and upwards to this region and leptopions would be generated in the strong magnetic fields generating strong electromagnetic instanton density $E \cdot B$ generating lepto-pion coherent state. If only positrons are observed, the absence of electrons could be due to different direction of accelerate motion for electrons. Since electrons are observed at the surface of Earth, this would suggest that electron positron pairs are created below the airplane.
3. But how it is possible to observe gamma rays and ultrahigh energy electrons at the surface of Earth? The problem is that atmosphere is not empty and dissipation would restrict the energies to be much lower than gamma ray energies which are in MeV range. Note that the temperatures in lightning are about 3×10^4 K and correspond to electron energy of 2.6 eV which is by a factor 10^5 smaller than electron mass and gamma ray energy scale! And how the electrons with energies above MeV range are created in thunder cloud? For years ago I proposed a model for high energy gamma rays and electrons associated with lightnings in terms of dark matter identified as $h_{eff} = n \times h$ phases. This model could provide answer to these questions.

First some background is needed.

1. I ended up to $h_{eff} = n \times h$ hypothesis [K6, K19] from the observations of Blackman and other pioneers of bio-electromagnetism [J1] about quantal effects of ELF em fields to vertebrate brain, which he explained in terms of cyclotron frequencies of Ca^+ ion in endoneous magnetic field $B_{end} = 0.2$ Gauss (2/5:th of the nominal value $B_E = .5$ Gauss of the Earth's magnetic field). Cyclotron energy $E = h \times f$ is however extremely low, much below the thermal energy in physiological temperature so that no quantal effects should be possible. This inspired the hypothesis $h_{eff} = n \times h$ scaling up the energy.
2. Nottale [E2] introduced originally the notion of gravitational Planck constant $\hbar_{gr} = GMm/v_0$ to explain the orbital radii of planets in solar system as Bohr orbits. The velocity parameter v_0 is different for inner and outer planets and quite recently I proposed v_0 is in constant ratio to the rotation velocity of the large mass M. The interpretation in TGD framework is that the magnetic flux tubes mediate gravitational interaction between M and m and the value of Planck constant is \hbar_{gr} at them. The proposal $h_{eff} = \hbar_{gr}$ at flux tubes is very natural sharpening of the original hypothesis [K19, K20]. The predictions of the model do not depend on whether m is taken to be the mass of the planet or any elementary particle associated with it and the gravitational Compton length $\lambda_{gr} = GMc/v_0$ does not depend on the mass of the particle as is proportional to the Schwarzschild radius $2GM$ of Sun.
3. This hypothesis can be generalized to apply also to Earth (see <http://tinyurl.com/ht4pwy7>). For the strength $B_{gal} \sim 1$ nT for galactic magnetic field assumed to mediate Earth's gravitational interaction cyclotron frequency 10 Hz in alpha band is mapped to cyclotron frequency scale of 72 minutes. Scaled EEG range corresponds to cyclotron periods varying up to 12 hours for B_{gal} . For $M = M_E$ and B_{gal} the cyclotron energy corresponds to about 1 eV at the lower end of visible photon energies.
4. What about the interpretation of ordinary EEG in terms of cyclotron frequencies assuming that the corresponding energies are in visible and UV range corresponding to the variation of B_{end} ? M_E is certainly too large to give a spectrum of cyclotron energies in this range

suggested by Blackman to explain the findings about quantal effects of ELF radiation on brain not possible in standard quantum theory because the energy is much below the thermal threshold. $M_D = .5 \times 10^{-4} M_E$ would be needed. I have proposed that M_D corresponds to a mass assignable to a spherical layer at distance of Moon's orbital radius and there are independent pieces of evidence for the existence of this layer. B_{end} would represent the lower bound for the value range of the magnetic field varying at least by 7 octaves would give the highest UV energies around 124 eV. The transformation of dark photons to ordinary photons would yield biophotons with energies in visible and UV range. Also B_{gal} would have some variation range.

5. This has a connection to quantum biology and neuroscience. The proposal is that dark cyclotron photons with energies in visible and UV range associated with flux tubes of magnetic field of appropriate strength serve as a communication tool making biological body (BB) to communicate sensory data to magnetic body (MB) and allow BB to control BB. The recent model involves

Consider now the model for how electrons and gamma rays accompanying lightnings can travel to the surface of Earth without dissipating their energies and how the collisions of electrons with gamma ray energies generating electropions are possible.

1. What happens if one replaces M_D with M_E meaning that also Earth's gravitons would reside at the flux tubes of B_{end} ? The energies get scale up by a factor $M_E/M_1 = 2 \times 10^4$ and this scales up the 1-100 eV range .02-2 MeV so that also gamma ray energies would be obtained.
2. The earlier proposal was that electrons and gamma rays associated with lightning arrive to the surface of Earth along dark magnetic flux tubes so that by macroscopic quantum coherence in scale of λ_{gr} they do not dissipate their energy.

3.2.4 Experimental evidence for colored muons

Also μ and τ should possess colored excitations. About fifteen years after this prediction was made. Direct experimental evidence for these states finally emerges (the year I am adding this comment is 2007) [C38, C39]. The mass of the new particle, which is either scalar or pseudo-scalar, is 214.4 MeV whereas muon mass is 105.6 MeV. The mass is about 1.5 per cent higher than two times muon mass. The proposed interpretation is as a light Higgs. I do not immediately resonate with this interpretation although p-adically scaled up variants of also Higgs bosons live happily in the fractal Universe of TGD. The most natural TGD inspired interpretation is as a pion like bound state of colored excitations of muon completely analogous to lepto-pion (or rather, electro-pion).

Scaled up variants of QCD appear also in nuclear string model [K14, L1], [L1], where scaled variant of QCD for exotic quarks in p-adic length scale of electron is responsible for the binding of 4He nuclei to nuclear strings. One cannot exclude the possibility that the fermion and anti-fermion at the ends of color flux tubes connecting nucleons are actually colored leptons although the working hypothesis is that they are exotic quark and anti-quark. One can of course also turn around the argument: could it be that lepto-pions are "leptonuclei", that is bound states of ordinary leptons bound by color flux tubes for a QCD in length scale considerably shorter than the p-adic length scale of lepton.

3.3 Evidence For τ -Hadrons

The evidence for τ -leptons came in somewhat funny but very pleasant manner. During my friday morning blog walk, the day next to my birthday October 30, I found that Peter Woit had told in his blog about a possible discovery of a new long-lived particle by CDF experiment [C44] emphasizing how revolutionary finding is if it is real. There is a detailed paper [C10] with title *Study of multi-muon events produced in p-pbar collisions at $\sqrt{s} = 1.96$ TeV* by CDF collaboration added to the ArXiv October 29 - the eve of my birthday. I got even second gift posted to arXiv the very same day and reporting an anomalously high abundance of positrons in cosmic ray radiation [C15]. Both of these article give support for basic predictions of TGD differentiating between TGD and standard model and its generalizations.

3.3.1 The first gift

A brief summary of Peter Woit about the finding gives good idea about what is involved.

The article originates in studies designed to determine the b - b bar cross-section by looking for events, where a b - b bar pair is produced, each component of the pair decaying into a muon. The b -quark lifetime is of order a picosecond, so b -quarks travel a millimeter or so before decaying. The tracks from these decays can be reconstructed using the inner silicon detectors surrounding the beam-pipe, which has a radius of 1.5 cm. They can be characterized by their impact parameter, the closest distance between the extrapolated track and the primary interaction vertex, in the plane transverse to the beam.

If one looks at events where the b -quark vertices are directly reconstructed, fitting a secondary vertex, the cross-section for b - b bar production comes out about as expected. On the other hand, if one just tries to identify b -quarks by their semi-leptonic decays, one gets a value for the b - b bar cross-section that is too large by a factor of two. In the second case, presumably there is some background being misidentified as b - b bar production.

The new result is based on a study of this background using a sample of events containing two muons, varying the tightness of the requirements on observed tracks in the layers of the silicon detector. The background being searched for should appear as the requirements are loosened. It turns out that such events seem to contain an anomalous component with unexpected properties that disagree with those of the known possible sources of background. The number of these anomalous events is large (tens of thousands), so this cannot just be a statistical fluctuation.

One of the anomalous properties of these events is that they contain tracks with large impact parameters, of order a centimeter rather than the hundreds of microns characteristic of b -quark decays. Fitting this tail by an exponential, one gets what one would expect to see from the decay of a new, unknown particle with a lifetime of about 20 picoseconds. These events have further unusual properties, including an anomalously high number of additional muons in small angular cones about the primary ones.

The lifetime is estimated to be considerably longer than b quark life time and below the lifetime 89.5 ps of $K_{0,s}$ mesons. The fit to the tail of “ghost” muons gives the estimate of 20 picoseconds.

3.3.2 The second gift

In October 29 also another remarkable paper [C15] had appeared in arXiv. It was titled *Observation of an anomalous positron abundance in the cosmic radiation*. PAMELA collaboration finds an excess of cosmic ray positron at energies 10 \rightarrow 50 GeV. PAMELA anomaly is discussed in Resonaances blog [C1]. ATIC collaboration in turn sees an excess of electrons and positrons going all the way up to energies of order 500-800 GeV [C37].

Also Peter Woit refers to these cosmic ray anomalies and also to the article *LHC Signals for a SuperUnified Theory of Dark Matter* by Nima Arkadi-Hamed and Neal Weiner [C6], where a model of dark matter inspired by these anomalies is proposed together with a prediction of lepton jets with invariant masses with mass scale of order GeV. The model assumes a new gauge interaction for dark matter particles with Higgs and gauge boson masses around GeV. The prediction is that LHC should detect “lepton jets” with smaller angular separations and GeV scale invariant masses.

3.3.3 Explanation of the CDF anomaly

Consider first the CDF anomaly. TGD predicts a fractal hierarchy of QCD type physics. In particular, colored excitations of leptons are predicted to exist. Neutral lepto-pions would have mass only slightly above two times the charged lepton mass. Also charged lepto-pions are predicted and their masses depend on what is the p -adic mass scale of neutrino and it is not clear whether it is much longer than that for charge colored lepton as in the case of ordinary leptons.

1. There exists a considerable evidence for colored electrons as already found. The anomalous production of electron positron pairs discovered in heavy ion collisions can be understood in terms of decays of electro-pions produced in the strong non-orthogonal electric and magnetic fields created in these collisions. The action determining the production rate would

be proportional to the product of the lepto-pion field and highly unique “instanton” action for electromagnetic field determined by anomaly arguments so that the model is highly predictive.

2. Also the 511 MeV emission line [C22, C34] from the galactic center can be understood in terms of decays of neutral electro-pions to photon pairs. Electro-pions would reside at magnetic flux tubes of strong galactic magnetic fields. It is also possible that these particles are dark in TGD sense.
3. There is also evidence for colored excitations of muon and muo-pion [C38, C39]. Muo-pions could be produced by the same mechanism as electro-pions in high energy collisions of charged particles when strong non-orthogonal magnetic and electric fields are generated.

Also τ -hadrons are possible and CDF anomaly can be understood in terms of a production of higher energy τ -hadrons as the following argument demonstrates.

1. τ -QCD at high energies would produce “lepton jets” just as ordinary QCD. In particular, muon pairs with invariant energy below $2m(\tau) \sim 3.6$ GeV would be produced by the decays of neutral τ -pions. The production of monochromatic gamma ray pairs is predicted to dominate the decays. Note that the space-time sheet associated with both ordinary hadrons and τ lepton correspond to the p-adic prime $M_{107} = 2^{107} - 1$.
2. The model for the production of electro-pions in heavy ion collisions suggests that the production of τ -pions could take place in higher energy collisions of protons generating very strong non-orthogonal magnetic and electric fields. This This would reduce the model to the quantum model for electro-pion production.
3. One can imagine several options for the detailed production mechanism.
 - (a) The decay of *virtual* τ -pions created in these fields to pairs of leptobaryons generates lepton jets. Since colored leptons correspond to color octets, leptobaryons could correspond to states of form LLL or $L\bar{L}\bar{L}$.
 - (b) The option inspired by a blog discussion with Ervin Goldfein is that a coherent state of τ -pions is created first and is then heated to QCD plasma like state producing the lepton jets like in QCD. The linear coupling to $E \cdot B$ defined by em fields of colliding nucleons would be analogous to the coupling of harmonic oscillator to constant force and generate the coherent state.
 - (c) The option inspired by CDF model [C25] is that a p-adically scaled up variant of *on mass shell* neutral τ -pion having $k = 103$ and 4 times larger mass than $k = 107$ τ -pion is produced and decays to three $k = 105$ τ -pions with $k = 105$ neutral τ -pion in turn decaying to three $k = 107$ τ -pions.
4. The basic characteristics of the anomalous muon pair prediction seems to fit with what one would expect from a jet generating a cascade of τ -pions. Muons with both charges would be produced democratically from neutral τ -pions; the number of muons would be anomalously high; and the invariant masses of muon pairs would be below 3.6 GeV for neutral τ -pions and below 1.8 GeV for charged τ -pions if colored neutrinos are light.
5. The lifetime of 20 ps can be assigned with charged τ -pion decaying weakly only into muon and neutrino. This provides a killer test for the hypothesis. In absence of CKM mixing for colored neutrinos, the decay rate to lepton and its antineutrino is given by

$$\Gamma(\pi_\tau \rightarrow L + \bar{\nu}_L) = \frac{G^2 m(L)^2 f^2(\pi) (m(\pi_\tau)^2 - m(L)^2)^2}{4\pi m^3(\pi_\tau)} . \quad (3.1)$$

The parameter $f(\pi_\tau)$ characterizing the coupling of pion to the axial current can be written as $f(\pi_\tau) = r(\pi_\tau)m(\pi_\tau)$. For ordinary pion one has $f(\pi) = 93$ MeV and $r(\pi) = .67$. The decay rate for charged τ -pion is obtained by simple scaling giving

$$\begin{aligned}\Gamma(\pi_\tau \rightarrow L + \bar{\nu}_L) &= 8x^2u^2y^3(1-z^2)\frac{1}{\cos^2(\theta_c)}\Gamma(\pi \rightarrow \mu + \bar{\nu}_\mu) , \\ x &= \frac{m(L)}{m(\mu)} , \quad y = \frac{m(\tau)}{m(\pi)} , \quad z = \frac{m(L)}{2m(\tau)} , \quad u = \frac{r(\pi_\tau)}{r(\pi)} .\end{aligned}\tag{3.2}$$

If the p-adic mass scale of the colored neutrino is same as for ordinary neutrinos, the mass of charged lepto-pion is in good approximation equal to the mass of τ and the decay rates to τ and electron are for the lack of phase space much slower than to muons so that muons are produced preferentially.

6. For $m(\tau) = 1.8$ GeV and $m(\pi) = .14$ GeV and the same value for f_π as for ordinary pion the lifetime is obtained by scaling from the lifetime of charged pion about 2.6×10^{-8} s. The prediction is 3.31×10^{-12} s to be compared with the experimental estimate about 20×10^{-12} s. $r(\pi_\tau) = .41r_\pi$ gives a correct prediction. Hence the explanation in terms of τ -pions seems to be rather convincing unless one is willing to believe in really nasty miracles.
7. Neutral τ -pion would decay dominantly to monochromatic pairs of gamma rays. The decay rate is dictated by the product of τ -pion field and ‘‘instanton’’ action, essentially the inner product of electric and magnetic fields and reducing to total divergence of instanton current locally. The rate is given by

$$\begin{aligned}\Gamma(\pi_\tau \rightarrow \gamma + \gamma) &= \frac{\alpha_{em}^2 m^3(\pi_\tau)}{64\pi^3 f(\pi_\tau)^2} = 2x^{-2}y \times \Gamma(\pi \rightarrow \gamma + \gamma) , \\ x &= \frac{f(\pi_\tau)}{m(\pi_\tau)} , \quad y = \frac{m(\tau)}{m(\pi)} . \Gamma(\pi \rightarrow \gamma + \gamma) = 7.37 \text{ eV} .\end{aligned}\tag{3.3}$$

The predicted lifetime is 1.17×10^{-17} seconds.

8. Second decay channel is to lepton pairs, with muon pair production dominating for kinematical reasons. The invariant mass of the pairs is 3.6 GeV if no other particles are produced. Whether the mass of colored neutrino is essentially the same as that of charged lepton or corresponds to the same p-adic scale as the mass of the ordinary neutrino remains an open question. If colored neutrino is light, the invariant mass of muon-neutrino pair is below 1.78 GeV.

3.3.4 PAMELA and ATIC anomalies

TGD predicts also a hierarchy of hadron physics assignable to Mersenne primes. The mass scale of M_{89} hadron physics is by a factor 512 higher than that of ordinary hadron physics. Therefore a very rough estimate for the nucleons of this physics is 512 GeV. This suggests that the decays of M_{89} hadrons are responsible for the anomalous positrons and electrons up to energies 500-800 GeV reported by ATIC collaboration. An equally naive scaling for the mass of pion predicts that M_{89} pion has mass 72 GeV. This could relate to the anomalous cosmic ray positrons in the energy interval 10-50 GeV reported by PAMELA collaboration. Be as it may, the prediction is that M_{89} hadron physics exists and could make itself visible in LHC.

The surprising finding is that positron fraction (the ratio of flux of positrons to the sum of electron and positron fluxes) increases above 10 GeV. If positrons emerge from secondary production during the propagation of cosmic ray-nuclei, this ratio should decrease if only standard physics is involved with the collisions. This is taken as evidence for the production of electron-positron pairs, possibly in the decays of dark matter particles.

Leptohadron hypothesis predicts that in high energy collisions of charged nuclei with charged particles of matter it is possible to produce also charged electro-pions, which decay to electrons or

positrons depending on their charge and produce the electronic counterparts of the jets discovered in CDF. This proposal - and more generally lepto-hadron hypothesis - could be tested by trying to find whether also electronic jets can be found in proton-proton collisions. They should be present at considerably lower energies than muon jets. I decided to check whether I have said something about this earlier and found that I have noticed years ago that there is evidence for the production of anomalous electron-positron pairs in hadronic reactions [C19, C28, C21, C47]: some of it dates back to seventies.

The first guess is that the center of mass energy at which the jet formation begins to make itself visible is in a constant ratio to the mass of charged lepton. From CDF data this ratio satisfies $\sqrt{s}/m_\tau = x < 10^3$. For electro-pions the threshold energy would be around $10^{-3}x \times .5$ GeV and for muo-pions around $10^{-3}x \times 100$ GeV.

3.3.5 Comparison of TGD model with the model of CDF collaboration

Few days after the experimental a theoretical paper by CDF collaboration proposing a phenomenological model for the CDF anomaly appeared in the arXiv [C25], and it is interesting to compare the model with TGD based model (or rather, one of them corresponding to the third option mentioned above).

The paper proposes that three new particles are involved. The masses for the particles - christened h_3 , h_2 , and h_1 - are assumed to be 3.6 GeV, 7.3 GeV, and 15 GeV. h_1 is assumed to be pair produced and decay to h_2 pair decaying to h_3 pair decaying to a τ pair.

h_3 is assumed to have mass 3.6 GeV and life-time of 20×10^{-12} seconds. The mass is same as the TGD based prediction for neutral τ -pion mass, whose lifetime however equals to 1.12×10^{-17} seconds ($\gamma + \gamma$ decay dominates). The correct prediction for the lifetime provides a strong support for the identification of long-lived state as charged τ -pion with mass near τ mass so that the decay to μ and its antineutrino dominates. Hence the model is not consistent with lepto-hadronic model.

p-Adic length scale hypothesis predicts that allowed mass scales come as powers of $\sqrt{2}$ and these masses indeed come in good approximation as powers of 2. Several p-adic scales appear in low energy hadron physics for quarks and this replaces Gell-Mann formula for low-lying hadron masses. Therefore one can ask whether the proposed masses correspond to neutral tau-pion with $p = M_k = 2^k - 1$, $k = 107$, and its p-adically scaled up variants with $p \simeq 2^k$, $k = 105$, and $k = 103$ (also prime). The prediction for masses would be 3.6 GeV, 7.2 GeV, 14.4 GeV.

This co-incidence cannot of course be taken too seriously since the powers of two in CDF model have a rather mundane origin: they follow from the assumed production mechanism producing 8 τ -leptons from h_1 . One can however spend some time by looking whether it could be realized somehow allowing p-adically scaled up variants of τ -pion.

1. The proposed model for the production of muon jets is based on production of $k=103$ neutral τ -pion (or several of them) having 4 times larger mass than $k=107$ τ -pion in strong EB background of the colliding proton and antiproton and decaying via weak boson and gluon exchanges to $k=105$ and $k=107$ τ -pions. The simplest decays are parity breaking $1 \rightarrow 2$ decays and must involve exchange of virtual W or Z boson. Three-pion coupling λ with dimensions of mass determines the decay rates for neutral τ -pions appearing in the cascade. For the four-pion decay the coupling is dimensionless. Rates are proportional to phase space-volumes, which are rather small by kinetic reasons and also reduced by weak coupling.
2. For a neutral initial state the first step could be one of the following ones:

$$\begin{aligned}
 \pi_\tau^0(103) &\rightarrow \pi_\tau^+(105) + \pi_\tau^-(105) \\
 \pi_\tau^0(103) &\rightarrow \pi_\tau^0(105) + \pi_\tau^0(105) \\
 \pi_\tau^0(103) &\rightarrow 2\gamma \\
 \pi_\tau^0(103) &\rightarrow \pi_\tau^+(105) + \pi_\tau^-(107) + \pi_\tau^0(107)
 \end{aligned}$$

In the last decay permutations of the final state charges are possible. Since the last reaction is parity conserving and governed by strong interactions it dominates. This step is not kinematically possible if masses are obtained by exact scaling and if $m(\pi_\tau^0) < m(\pi_\tau^\pm)$ holds true as for ordinary pion. p-Adic mass formulas do not however predict exact scaling. In the

case that reaction is not kinematically possible, it must be replaced with a reaction in which one final state pion is virtual.

3. At the second step charged pion would decay to two pions

$$\pi_{\tau}^{\pm}(105) \rightarrow \pi_{\tau}^0(107) + \pi_{\tau}^{\pm}(107) \ ,$$

Neutral pion could decay to two gammas or to two pions

$$\pi_{\tau}^0(105) \rightarrow 2\gamma \text{ or } \pi_{\tau}^+(107) + \pi_{\tau}^-(107) \text{ or } \pi_{\tau}^0(107) + \pi_{\tau}^0(107) \ ,$$

Here second charged pion also can be virtual and decay weakly, and the weak decays of the $\pi_{\tau}^{\pm}(105)$ with mass $2m(\tau)$ to lepton pairs. The rates for these are obtained from previous formulas by scaling. For neutral pion the decay to two gammas dominates now.

4. The last step would involve the decays of both charged and neutral $\pi_{\tau}(107)$. The signature of the mechanism would be anomalous γ pairs with invariant masses $2^k \times m(\tau)$, $k = 1, 2, 3$ coming from the decays of neutral τ -pions.

The total cross section for producing single lepto-pion can be estimated by using the quantum model for lepto-pion production. Production amplitude is essentially Coulomb scattering amplitude for a given value of the impact parameter b for colliding proton and anti-proton multiplied by the amplitude $U(b, p)$ for producing on mass shell $k = 103$ lepto-pion with given four-momentum in the fields E and B and given essentially by the Fourier transform of $E \cdot B$. The replacement of the motion with free motion should be a good approximation.

UV and IR cutoffs for the impact parameter appear in the model and are identifiable as appropriate p-adic length scales. UV cutoff could correspond to the Compton size of nucleon ($k = 107$) and IR cutoff to the size of the space-time sheets representing topologically quantized electromagnetic fields of colliding nucleons (perhaps $k = 113$ corresponding to nuclear p-adic length scale and size for color magnetic body of constituent quarks or $k = 127$ for the magnetic body of current quarks with mass scale of order MeV). If one has $\hbar/\hbar_0 = 2^7$ one could also guess that the IR cutoff corresponds to the size of dark em space-time sheet equal to $2^7 L(113) = L(127)$ (or $2^7 L(127) = L(141)$), which corresponds to electron's p-adic length scale. These are of course rough guesses.

Quantitatively the jet-likeness of muons means that the additional muons are contained in the cone $\theta < 36.8$ degrees around the initial muon direction. If the decay of $\pi_{\tau}^0(k)$ can occur to on mass shell $\pi_{\tau}^0(k+2)$, $k = 103, 105$, it is possible to understand jets as a consequence of the decay kinematics forcing the pions resulting as decay products to be almost at rest.

1. Suppose that the decays to three pions can take place as on mass shell decays so that pions are very nearly at rest. The distribution of decay products $\mu\bar{\nu}$ in the decays of $\pi^{\pm}(105)$ is spherically symmetric in the rest frame and the energy and momentum of the muon are given by

$$[E, p] = [m(\tau) + \frac{m^2(\mu)}{4m(\tau)}, m(\tau) - \frac{m^2(\mu)}{4m(\tau)}] \ .$$

The boost factor $\gamma = 1/\sqrt{1-v^2}$ to the rest system of muon is $\gamma = \frac{m(\tau)}{m(\mu)} + \frac{m(\mu)}{4m(\tau)} \sim 18$.

2. The momentum distribution for μ^+ coming from π_{τ}^+ is spherically symmetric in the rest system of π^+ . In the rest system of μ^- the momentum distribution is non-vanishing only for when the angle θ between the direction of velocity of μ^- is below a maximum value of given by $\tan(\theta_{max}) = 1$ corresponding to a situation in which the momentum μ^+ is orthogonal to the momentum of μ^- (the maximum transverse momentum equals to $m(\mu)v\gamma$ and longitudinal momentum becomes $m(\mu)v\gamma$ in the boost). This angle corresponds to 45 degrees and is not too far from 36.8 degrees.

3. At the next step the energy of muons resulting in the decays of $\pi^\pm(103)$

$$[E, p] = \left[\frac{m(\tau)}{2} + \frac{m^2(\mu)}{2m(\tau)}, \frac{m(\tau)}{2} - \frac{m^2(\mu)}{2m(\tau)} \right],$$

and the boost factor is $\gamma_1 = \frac{m(\tau)}{2m(\mu)} + \frac{m(\mu)}{2m(\tau)} \sim 9$. θ_{max} satisfies the condition $\tan(\theta_{max}) = \gamma_1 v_1 / \gamma v \simeq 1/2$ giving $\theta_{max} \simeq 26.6$ degrees.

If on mass shell decays are not allowed the situation changes since either of the charged pions is off mass shell. In order to obtain similar result the virtual should occur dominantly via states near to on mass shell pion. Since four-pion coupling is just constant, this option does not seem to be realized.

Quantitatively the jet-likeness of muons means that the additional muons are contained in the cone $\theta < 36.8$ degrees around the initial muon direction. If the decay of $\pi_\tau^0(k)$ can occur to on mass shell $\pi_\tau^0(k+2)$, $k = 103, 105$, it is possible to understand jets as a consequence of the decay kinematics forcing the pions resulting as decay products to be almost at rest.

1. Suppose that the decays to three pions can take place as on mass shell decays so that pions are very nearly at rest. The distribution of decay products $\mu\bar{\nu}$ in the decays of $\pi^\pm(105)$ is spherically symmetric in the rest frame and the energy and momentum of the muon are given by

$$[E, p] = \left[m(\tau) + \frac{m^2(\mu)}{4m(\tau)}, m(\tau) - \frac{m^2(\mu)}{4m(\tau)} \right].$$

The boost factor $\gamma = 1/\sqrt{1-v^2}$ to the rest system of muon is $\gamma = \frac{m(\tau)}{m(\mu)} + \frac{m(\mu)}{4m(\tau)} \sim 18$.

2. The momentum distribution for μ^+ coming from π_τ^+ is spherically symmetric in the rest system of π^+ . In the rest system of μ^- the momentum distribution is non-vanishing only for when the angle θ between the direction of velocity of μ^- is below a maximum value of given by $\tan(\theta_{max}) = 1$ corresponding to a situation in which the momentum μ^+ is orthogonal to the momentum of μ^- (the maximum transverse momentum equals to $m(\mu)v\gamma$ and longitudinal momentum becomes $m(\mu)v\gamma$ in the boost). This angle corresponds to 45 degrees and is not too far from 36.8 degrees.
3. At the next step the energy of muons resulting in the decays of $\pi^\pm(103)$

$$[E, p] = \left[\frac{m(\tau)}{2} + \frac{m^2(\mu)}{2m(\tau)}, \frac{m(\tau)}{2} - \frac{m^2(\mu)}{2m(\tau)} \right],$$

and the boost factor is $\gamma_1 = \frac{m(\tau)}{2m(\mu)} + \frac{m(\mu)}{2m(\tau)} \sim 9$. θ_{max} satisfies the condition $\tan(\theta_{max}) = \gamma_1 v_1 / \gamma v \simeq 1/2$ giving $\theta_{max} \simeq 26.6$ degrees.

If on mass shell decays are not possible, the situation changes since either of the charged pions is off mass shell. In order to obtain similar result the virtual should occur dominantly via states near to on mass shell pion. Since four-pion coupling is just constant, this option does not seem to be realized.

3.3.6 Numerical estimate for the production cross section

The numerical estimate of the cross section involves some delicacies. The model has purely physical cutoffs which must be formulated in a precise manner.

1. Since energy conservation is not coded into the model, some assumption about the maximal τ -pion energy in cm system expressed as a fraction ϵ of proton's center of mass energy is necessary. Maximal fraction corresponds to the condition $m(\pi_\tau) \leq m(\pi_\tau)\gamma_1 \leq \epsilon m_p \gamma_{cm}$ in cm system giving $[m(\pi_\tau)] / (m_p \gamma_{cm}) \leq \epsilon \leq 1$. γ_{cm} can be deduced from the center of mass energy

of proton as $\gamma_{cm} = \sqrt{s}2m_p$, $\sqrt{s} = 1.96$ TeV. This gives $1.6 \times 10^{-2} < \epsilon < 1$ in a reasonable approximation. It is convenient to parameterize ϵ as

$$\epsilon = (1 + \delta) \times \frac{m(\pi_\tau)}{m_p} \times \frac{1}{\gamma_{cm}} .$$

The coordinate system in which the calculations are carried out is taken to be the rest system of (say) antiproton so that one must perform a Lorentz boost to obtain upper and lower limits for the velocity of τ -pion in this system. In this system the range of γ_1 is fixed by the maximal cm velocity fixed by ϵ and the upper/lower limit of γ_1 corresponds to a direction parallel/opposite to the velocity of proton.

2. By Lorentz invariance the value of the impact parameter cutoff b_{max} should be expressible in terms τ -pion Compton length and the center of mass energy of the colliding proton and the assumption is that $b_{max} = \gamma_{cm} \times \hbar/m(\pi_\tau)$, where it is assumed $m(\pi_\tau) = 8m(\tau)$. The production cross section does not depend much on the precise choice of the impact parameter cutoff b_{max} unless it is un-physically large in which case b_{max}^2 proportionality is predicted.

The numerical estimate for the production cross section involves some delicacies.

1. The power series expansion of the integral of CUT_1 using partial fraction representation does not converge since that roots c_\pm are very large in the entire integration region. Instead the approximation $A_1 \simeq iB\cos(\psi)/D$ simplifying considerably the calculations can be used. Also the value of b_1L is rather small and one can use stationary phase approximation for CUT_2 . It turns out that the contribution of CUT_2 is negligible as compared to that of CUT_1 .
2. Since the situation is singular for $\theta = 0$ and $\phi = 0$ and $\phi = \pi/2$ (by symmetry it is enough to calculate the cross section only for this kinematical region), cutoffs

$$\theta \in [\epsilon_1, (1 - \epsilon_1)] \times \pi \quad , \quad \phi \in [\epsilon_1, (1 - \epsilon_1)] \times \pi/2 \quad , \quad \epsilon_1 = 10^{-3} .$$

The result of the calculation is not very sensitive to the value of the cutoff.

3. Since the available numerical environment was rather primitive (MATLAB in personal computer), the requirement of a reasonable calculation time restricted the number of intervals in the discretization for the three kinematical variables γ, θ, ϕ to be below $N_{max} = 80$. The result of calculation did not depend appreciably on the number of intervals above $N = 40$ for γ_1 integral and for θ and ϕ integrals even $N = 10$ gave a good estimate.

The calculations were carried for the $exp(iS)$ option since in good approximation the estimate for $exp(iS) - 1$ model is obtained by a simple scaling. $exp(iS)$ model produces a correct order of magnitude for the cross section whereas $exp(iS) - 1$ variant predicts a cross section, which is by several orders of magnitude smaller by downwards α_{em}^2 scaling. As I asked Tommaso Dorigo for an estimate for the production cross section in his first blog posting [C49], he mentioned that authors refer to a production cross section is 100 nb which looks to me suspiciously large (too large by three orders of magnitude), when compared with the production rate of muon pairs from b-bbar. $\delta = 1.5$ which corresponds to τ -pion energy 36 GeV gives the estimate $\sigma = 351$ nb. The energy is suspiciously high.

In fact, in the recent blog posting of Tommaso Dorigo [C48] a value of order.1 nb for the production cross section was mentioned. Electro-pions in heavy ion collisions are produced almost at rest and one has $\Delta v/v \simeq .2$ giving $\delta = \Delta E/m(\pi) \simeq 2 \times 10^{-3}$. If one believes in fractal scaling, this should be at least the order of magnitude also in the case of τ -pion. This would give the estimate $\sigma = 1$ nb. For $\delta = \Delta E/m(\pi) \simeq 10^{-3}$ a cross section $\sigma = .16$ nb would result.

One must of course take the estimate cautiously but there are reasons to hope that large systematic errors are not present anymore. In any case, the model can explain also the order of magnitude of the production cross section under reasonable assumptions about cutoffs (see **Fig. 2**).




Figure 2: Differential cross section $\sin^2(\theta) \times \frac{d^2\sigma}{2E d^3p}$ for τ -pion production for $\gamma_1 = 1.090 \times 10^3$ in the rest system of antiproton for $\delta = 1.5$. $m(\pi_\tau)$ defines the unit of energy and nb is the unit for cross section. The ranges of θ and ϕ are $(0, \pi)$ and $(0, \pi/2)$.

3.3.7 Does the production of lepto-pions involve a phase transition increasing Planck constant?

The critical argument of Tommaso Dorigo in his blog inspired an attempt to formulate more precisely the hypothesis $\sqrt{s}/m_\tau > x < 10^3$. This led to the realization that a phase transition increasing Planck constant might happen in the production process as also the model for the production of electro-pions requires.

Suppose that the instanton coupling gives rise to *virtual* neutral lepto-pions which ultimately produce the jets (this is first of the three models that one can imagine). E and B could be associated with the colliding proton and antiproton or quarks.

1. The amplitude for lepto-pion production is essentially Fourier transform of $E \cdot B$, where E and B are the non-orthogonal electric and magnetic fields of the colliding charges. At the level of scales one has $\tau \sim \hbar/E$, where τ is the time during which $E \cdot B$ is large enough during collision and E is the energy scale of the virtual lepto-pion giving rise to the jet.
2. In order to have jets one must have $m(\pi_\tau) \ll E$. If the scaling law $E \propto \sqrt{s}$ hold true, one indeed has $\sqrt{s}/m(\pi_\tau) > x < 10^3$.
3. If proton and antiproton would move freely, τ would be of the order of the time for proton to move through a distance, which is 2 times the Lorentz contracted radius of proton: $\tau_{free} = 2 \times \sqrt{1 - v^2} R_p / v = 2\hbar/E_p$. This would give for the energy scale of virtual τ -pion the estimate $E = \hbar/\tau_{free} = \sqrt{s}/4$. $x = 4$ is certainly quite too small value. Actually $\tau > \tau_{free}$ holds true but one can argue that without new physics the time for the preservation of $E \cdot B$ cannot be by a factor of order 2^8 longer than for free collision.
4. For a colliding quark pair one would have $\tau_{free} = 4\hbar/\sqrt{s_{pair}(s)}$, where $\sqrt{s_{pair}(s)}$ would be the typical invariant energy of the pair which is exponentially smaller than \sqrt{s} . Somewhat paradoxically from classical physics point of view, the time scale would be much longer for the collision of quarks than that for proton and antiproton.

The possible new physics relates to the possibility that lepto-pions are dark matter in the sense that they have Planck constant larger than the standard value.

1. Suppose that the produced lepto-pions have Planck constant larger than its standard value \hbar_0 . Originally the idea was that larger value of \hbar would scale up the production cross section. It turned out that this is not the case. For $exp(iS)$ option the lowest order contribution is not affected by the scaling of \hbar and for $exp(iS) - 1$ option the lowest order contribution scales down as $1/\hbar^2$. The improved formulation of the model however led to a correct order of magnitude estimates for the production cross section.
2. Assume that a phase transition increasing Planck constant occurs during the collision. Hence τ is scaled up by a factor $y = \hbar/\hbar_0$. The inverse of the lepto-pion mass scale is a natural candidate for the scaled up dark time scale. $\tau(\hbar_0) \sim \tau_{free}$, one obtains $y \sim \sqrt{s_{min}}/4m(\pi_\tau) \leq 2^8$ giving for proton-antiproton option the first guess $\sqrt{s}/m(\pi_\tau) > x < 2^{10}$. If the value of y does not depend on the type of lepto-pion, the proposed estimates for muo- and electro-pion follow.
3. If the fields E and B are associated with colliding quarks, only colliding quark pairs with $\sqrt{s_{pair}(s)} > (>)m(\pi_\tau)$ contribute giving $y_q(s) = \sqrt{s_{pair}(s)}/s \times y$.

If the τ -pions produced in the magnetic field are on-mass shell τ -pions with $k = 113$, the value of \hbar would satisfy $\hbar/\hbar_0 < 2^5$ and $\sqrt{s}/m(\pi_\tau) > x < 2^7$.

3.3.8 Tau-pions again but now as dark matter candidate in galactic center

The standard view about dark matter is that it has only gravitational interactions with ordinary matter so that high densities of dark matter are required to detect its signatures. On the average the density of dark matter is about 80 per cent of ordinary matter. Clearly, Milky Way's center is an excellent place for detecting the signatures of dark matter. The annihilation of pairs of dark matter particles to gamma rays is one possible signature and one could study the anomalous features of gamma ray spectrum from the galactic center (a region with radius about 100 light years).

Europe's INTEGRAL satellite launched in 2002 indeed found bright gamma ray radiations coming from the center of galaxy with energy of 511 MeV, which is slightly above electron mass (see the references below). The official interpretation is that the gammas are produced in the annihilations of particles of positrons and electrons in turn created in dark matter annihilations. TGD suggests much simpler mechanism. Gamma rays would be produced in the decay of what I call electro-pions having mass which is slightly larger than $m = 2m_e$.

The news of the day [C45] was that the data from Fermi Gamma Ray telescope give analyzed by Dan Hooper and Lisa Goodenough [C40] gives evidence for a dark matter candidate with mass between 7.3-9.2 GeV decaying predominantly into a pair of τ leptons. The estimate for the mass region is roughly 4 times τ mass. What puts bells ringing that a mass of a charged lepton appears again!

1. Explanation in TGD framework

The new finding fits nicely to a bigger story based on TGD.

1. TGD predicts that both quarks and leptons should have colored excitations devoted to the lepto-hadron model). In the case of leptons lowest excitations are color octets. In the case of electro-pion this hypothesis finds support from the anomalous production of electron positron pairs in heavy ion collisions discovered already at seventies but forgotten for long ago since the existence of light particle at this mass scale simply was in total complete with standard model and what was known about the decay widths of intermediate gauge bosons. Also orthopositronium decay width anomaly -forgotten also-has explanation in terms of lepto-pion hypothesis [C35, C17].
2. The colored leptons would be dark in TGD sense, which means that they live in dark sector of the "world of classical worlds" (WCW) meaning that they have no direct interactions (common vertices of Feynman diagrams) with ordinary matter. They simply live at different

space-time sheets. A phase transition which is geometrically a leakage between dark sector and ordinary sector are possible and make possible interactions between ordinary and dark matter based on exchanged particles suffering this phase transition. Therefore the decay widths of intermediate gauge bosons do not kill the model. TGD based model of dark matter in terms of hierarchy of values of Planck constants coming as multiples of its smallest possible value (the simplest option) need not to be postulated separately and can be regarded as a prediction of quantum TGD reflecting directly the vacuum degeneracy and extreme non-linearity of Kähler action (Maxwell action for induced CP_2 Kähler form).

3. CDF anomaly which created a lot of discussion in blogs for two years ago can be understood in terms of taupion. Taupion and its p-adically scaled up versions with masses about $2^k m_\tau$, $k = 1, 2, 3$ and $m_\tau \simeq 1.8$ GeV explains the findings reported by CDF in TGD framework. The masses of taupions would be 3.6 GeV, 7.2 GeV, and 14.2 GeV in good approximation and come as octaves of the mass of tau-lepton pair.

2. Predictions

The mass estimate for the dark matter particle suggested by Fermi Gamma Ray telescope corresponds to $k = 2$ octave for taupion and the predicted mass is about 7.2 GeV which at the lower boundary of the range 7.3-9.2 GeV. Also dark matter particles decaying to tau pairs and having masses 3.6 GeV and 14.2 GeV should be found.

Also muo-pion should exist there and should have mass slightly above $2m_\mu = 210.4$ MeV so that a gamma rays peak slightly above the energy $m_\mu = 105.2$ MeV should be discovered. Also octaves of this mass can be imagined. There is also evidence also for the existence of muo-pion [C38, C39].

LHC should provide excellent opportunities to test tau-pion and muo-pion hypothesis. Electropion was discovered in heavy ion collisions and also at LHC they study heavy ion collisions but at much higher energies generating the required very strong non-orthogonal electric and magnetic fields for which the “instanton density” defined as the inner product of electric and magnetic fields is large and rapidly varying. As an optimist I hope that muo-pion and tau-pion could be discovered despite the fact that their decay signatures are very different from those for ordinary particles and despite that fact that at these energies one must know precisely what one is trying to find in order to disentangle it from the enormous background.

3. Also DAMA, CoGeNT, and PAMELA give indications for tau-pion

Note that also DAMA experiment [C14] suggests the existence of dark matter particle in this mass range but it is not clear whether it can have anything to do with tau-pion state. One could of course imagine that dark tau-pions are created in the collisions of highly energetic cosmic rays with the nuclei of atmosphere. Also Coherent Germanium Neutrino Technology (CoGeNT) experiment [C12] has released data that are best explained in terms of a dark matter particle with mass in the range 7-11 GeV.

The decay of tau-pions produce lepton pairs, mostly tau but also muons and electrons. The subsequent decays of tau-leptons to muons and electrons produce also electrons and positrons. This relates interestingly to the positron excess reported by PAMELA collaboration [C15] at the same time as CDF anomaly was reported. The anomaly started at positron energy about 3.6 GeV, which is one just one half of 7.2 GeV for tau-pion mass! What was remarkable that no antiproton excess predicted by standard dark matter candidates was observed. Therefore the interpretation as decay products of tau-pions seems to make sense!

3.3.9 Could it have been otherwise?

To sum up, the probability that a correct prediction for the lifetime of the new particle using only known lepton masses and standard formulas for weak decay rates follows by accident is extremely low. Throwing billion times coin and getting the same result every time might be something comparable to this. Therefore my sincere hope is that colleagues would be finally mature to take TGD seriously. If TGD based explanation of the anomalous production of electron positron pairs in heavy ion collisions would have been taken seriously for fifteen years ago, particle physics might look quite different now.

3.4 Dark Matter Puzzle

Sean Carroll has explained in Cosmic Variance (<http://tinyurl.com/c2r2cv>) the latest rather puzzling situation in dark matter searches. Some experiments support the existence of dark matter particles with mass of about 7 GeV, some experiments exclude them. The following arguments show that TGD based explanation might allow to understand the discrepancy.

3.4.1 How to detect dark matter and what's the problem?

Consider first the general idea behind the attempts to detect dark matter particles and how one ends up with the puzzling situation.

1. Galactic nucleus serves as a source of dark matter particles and these one should be able to detect. There is an intense cosmic ray flux of ordinary particles from galactic center which must be eliminated so that only dark matter particles interacting very weakly with matter remain in the flux. The elimination is achieved by going sufficiently deep underground so that ordinary cosmic rays are shielded but extremely weakly interacting dark matter particles remain in the flux. After this one can in the ideal situation record only the events in which dark matter particles scatter from nuclei provided one eliminates events such as neutrino scattering.
2. DAMA experiment does not detect dark matter events as such but annual variations in the rate of events which can include besides dark matter events and other kind of events. DAMA finds an annual variation interpreted as dark matter signal since other sources of events are not expected to have this kind of variation [C13]. Also CoGENT has reported the annual variation with 2.8 sigma confidence level [C46]. The mass of the dark matter particle should be around 7 GeV rather than hundreds of GeVs as required by many models. An unidentified noise with annual variation having nothing to do with dark matter could of course be present and this is the weakness of this approach.
3. For a few weeks ago we learned that XENON100 experiment detects no dark matter [C16] (<http://tinyurl.com/y9t3gxxu>). Also CDMS has reported a negative result [C11]. According to Sean Carroll, the detection strategy used by XENON100 is different from that of DAMA: individual dark matter scatterings on nuclei are detected. This is a very significant difference which might explain the discrepancy since the theory laden prejudices about what dark matter particle scattering can look like, could eliminate the particles causing the annual variations. For instance, these prejudices are quite different for the habitants of the main stream Universe and TGD Universe.

3.4.2 TGD based explanation of the DAMA events and related anomalies

I have commented earlier the possible interpretation of DAMA events in terms of tau-pions (<http://tinyurl.com/ycgkvuy9>). The spirit is highly speculative.

1. Tau-pions would be identifiable as the particles claimed by Fermi Gamma Ray telescope with mass around 7 GeV and decaying into tau pairs so that one could cope with several independent observations instead of only single one.
2. Recall that the CDF anomaly gave for two and half years ago support for tau-pions whereas earlier anomalies dating back to seventies give support for electro-pions and mu-pions. The existence of these particles is purely TGD based phenomenon and due to the different view about the origin of color quantum numbers. In TGD colored states would be partial waves in CP_2 and spin like quantum numbers in standard theories so that leptons would not have colored excitations.
3. Tau-pions are of course highly unstable and would not come from the galactic center. Instead, they would be created in cosmic ray events at the surface of Earth and if they can penetrate the shielding eliminating ordinary cosmic rays they could produce events responsible for the annual variation caused by that for the cosmic ray flux from galactic center.

Can one regard tau-pion as dark matter in some sense? Or must one do so? The answer is affirmative to both questions on both theoretical and experimental grounds.

1. The existence of colored variants of leptons is excluded in standard physics by intermediate gauge boson decay widths. They could however appear as states with non-standard value of Planck constant and therefore not appearing in same vertices with ordinary gauge bosons so that they would not contribute to the decay widths of weak bosons. In this minimal sense they would be dark and this is what is required in order to understand what we know about dark matter.

Of course, all particles can in principle appear in states with non-standard value of Planck constant so that tau-pion would be one special instance of dark matter. For instance, in living matter the role of dark variants of electrons and possibly also other stable particles would be decisive. To put it bluntly: in mainstream approach dark matter is identified as some exotic particle with ad hoc properties whereas in TGD framework dark matter is outcome of a generalization of quantum theory itself.

2. DAMA experiment requires that the tau-pions behave like dark matter: otherwise they would never reach the strongly shielded detector. The interaction with the nuclei of detector would be preceded by a transformation to a particle-tau-pion or something else- with ordinary value of Planck constant.

3.4.3 TGD based explanation for the dark matter puzzle

The criteria used in experiments to eliminate events which definitely are not dark matter events - according to the prevailing wisdom of course - dictates to high degree what interactions of tau pions with solid matter detector are used as a signature of dark matter event. It could well be that the criteria used in XENON100 do not allow the scatterings of tau-pions with nuclei. This is indeed the case. The clue comes from the comments of Jester in Resonaances (see <http://tinyurl.com/yd3vs7x5>). From a comment of Jester one learns that CoGENT - and also DAMA utilizing the same detections strategy - “does not cut on ionization fraction”. Therefore, if dark matter mimics electron recoils (as Jester says) or if dark matter produced in the collisions of cosmic rays with the nuclei of the atmosphere decays to charged particles one can understand the discrepancy.

The TGD based model [K16] explaining the more than two years old CDF anomaly [C10, C27] indeed explains also the discrepancy between XENON100 and CDMS on one hand and DAMA and CoGENT on the other hand. The TGD based model for the CDF anomaly can be found in [K16].

1. To explain the observations of CDF [C10, C27] one had to assume that tau-pions and therefore also color excited tau-leptons inside them appear as several p-adically scaled up variants so that one would have several octaves of the ground state of tau-pion with masses in good approximation equal to 3.6 GeV (two times the tau-lepton mass), 7.2 GeV, 14.4 GeV. The 14.4 GeV tau-pion was assumed to decay in a cascade like manner via lepto-strong interactions to lighter tau-pions- both charged and neutral- which eventually decayed to ordinary charged leptons and neutrinos.
2. Also other decay modes -say the decay of neutral tau-pions to gamma pair and to a pair of ordinary leptons- are possible but the corresponding rates are much slower than the decay rates for cascade like decay via multi-tau-pion states proceeding via lepto-strong interactions.
3. Just this cascade would take place also now after the collision of the incoming cosmic ray with the nucleus of atmosphere. The mechanism producing the neutral tau-pions -perhaps a coherent state of them- would degenerate in the collision of charged cosmic ray with nucleus generating strong non-orthogonal electric and magnetic fields and the production amplitude would be essentially the Fourier transform of the “instanton density” $E \cdot B$. The decays of 14 GeV neutral tau-pions would produce 7 GeV charged tau-pions, which would scatter from the protons of nuclei and generate the events excluded by XENON100 but not by DAMA and Cogent.

4. In principle the model predicts to a high degree quantitatively the rate of the events. The scattering rates are proportional to an unknown parameter characterizing the transformation probability of tau-pion to a particle with ordinary value of Planck constant and this allows to perform some parameter tuning. This parameter would correspond to a mass insertion in the tau-pion line changing the value of Planck constant and have dimensions of mass squared.

The overall conclusion is that the discrepancy between DAMA and XENON100 might be interpreted as favoring TGD view about dark matter and it is fascinating to see how the situation develops. This confusion is not the only confusion in recent day particle physics. All believed-to-be almost-certainties are challenged.

3.4.4 Has Fermi observed dark matter?

Resonaances (<http://tinyurl.com/y79x4v3y>) reports about a possible dark matter signal at Fermi satellite [C9]. Also Lubos Motl (<http://tinyurl.com/yd7xsfxo1>) has a posting about the finding and mentions that the statistical significance is 3.3 sigma.

The proposed dark matter interpretation for the signal would be pair of monochromatic photons with second one detected at Earth. The interpretation would be that dark matter particles with mass m nearly at rest in galactic center annihilate to a pair of photons so that one obtains a pair of photons with energy equal to the cm energy which is in a good approximation the sum $E = 2 \times m$ for the masses of the particles. The mass value would be around $m=130$ GeV if the final state involves only 2 photons.

In TGD framework I would consider as a first guess a pion like state decaying to two photons with standard coupling given by the coupling to the “instanton density” $E \cdot B$ of electromagnetic field. The mass of this particle would be 260 GeV, in reasonable approximation 2 times the mass $m=125$ GeV of the Higgs candidate.

1. Similar coupling was assumed to [K16]. The anomaly would have been produced by tau-pions, which are pionlike states formed by pairs of colored excitations of tau and its antiparticle (or possibly their super-partners). What was remarkable that the mass had three values coming as powers of two: $M = 2^k \times 2m(\tau)$, $k = 0, 1, 2$. The interpretation in terms of p-adic length scale hypothesis would be obvious: also the octaves of the basic state are there. The constraint from intermediate gauge boson decay widths requires that these states are dark in TGD sense and therefore correspond to a non-standard value of Planck constant coming as an integer multiple of the standard value.
2. Also the explanation of the findings of PAMELA discussed in this chapter require octaves of tau-pion produced in Earth’s atmosphere.
3. Even ordinary pion should have 2-adic octaves. But doesn’t this kill the hypothesis? We “know” that pion does not have any octaves! Maybe not, there is recent evidence for satellites of ordinary pion with energy scale of 40 MeV interpreted in terms of IR Regge trajectories assignable to the color magnetic flux tubes assignable to pion. There has been several wrong alarms about Higgs: at 115 GeV and 155 GeV at least. Could it be that there there is something real behind these wrong alarms: the scale for IR Regge trajectories would be about 20 GeV now!

So: could the dark matter candidates with mass around 260 GeV correspond to the first octave of M_{89} pion with mass around 125 GeV, the particle that colleagues want to call Higgs boson although its decay signatures suggest something different?

1. In this case it does not seem necessary to assume that the Planck constant has non-standard value although this is possible.
2. This particle should be produced in M_{89} strong interactions in the galactic center. This would require the presence of matter consisting of M_{89} nucleons emitting these pions in strong interactions. Galactic center (<http://tinyurl.com/y9clyfs>) is very exotic place and believed to contain even super-massive black hole. Could this environment accommodate also a scaled up copy of hadron physics? Presumably this would require very high temperatures

with thermal energy of order 5 TeV correspond to the mass of M_{89} proton to make possible the presence of M_{89} matter. Or could M_{89} pion be produced in ultrastrong non-orthogonal electric and magnetic fields in the galactic center by the coupling to the instanton density. The needed field strengths would be extremely high. I have indeed proposed long time ago an explanation of very high energy cosmic rays in terms of the decay products of scaled up hadron physics (see “Cosmic Rays and Mersenne primes” in this chapter).

One can of course imagine that the photon pair is produced in the annihilation of M_{89} pions with opposite charges via standard electromagnetic coupling. Also the annihilation of M_{89} spions consisting of squark pair can be considered in TGD framework where squarks could have same mass scale as quarks. In this case mass would be near 125 GeV identified as mass of neutral M_{89} pion. By scaling up the mass difference 139.570-134.976 MeV of the ordinary charged and neutral pion by the ratio of the pion M_{89} and M_{107} pion masses equal to $(125/140) \times 10^3$ one obtains that the charged M_{89} pion should have mass equal to 129.6 MeV to be compared with the 130 GeV mass suggested by experimental evidence.

The story did not end here as so often when observations cannot be replicated. The Estonian researchers Elmo Tempel, Andi Hektora and Martti Raidala have found a confirmation for the 130 GeV Fermi excess in gamma radiation from galactic center discovered by Cristoph Weniger [E1]. An important conclusion of these researchers is that best fit is obtained if the dark matter candidates decay by two-body annihilation to photons and have mass 145 GeV. The reason for why the gamma peak is at 130 GeV rather than 145 GeV would be due to the emission light particle pairs by the photons. There are also indications for a peak at 111 GeV: this could be assigned to γZ final state of two-body decay.

In TGD framework the annihilating particles with mass about 145 GeV mass could be charged pion-like states of M_{89} hadron physics. They could be dark in the sense of having large value of Planck constant but it is not clear whether this is necessarily so. The TGD based on view about galactic dark matter locates in cosmic string like objects containing galaxies as pearls in necklace and no halo is needed to explain galactic rotation spectrum [K3]. An ultrahigh temperature would be needed to excite M_{89} hadron physics and if there is giant blackhole in galactic nucleus, there are hopes about this. M_{89} hadron physics could also produce ultrahigh energy cosmic rays as described in this chapter.

It is amusing that also CDF found for a couple of years ago evidence for a bump at the same 145 GeV energy (this has been forgotten long time ago by bloggers in 125 GeV Higgs hysteria). Estonians propose that also a particle with 290 GeV (mass would twice that of 145 GeV state) is needed. This brings further support for the idea about mass octaves of ground state of pionlike states needed to explain various anomalies (see this chapter and [K16]).

If one takes seriously the evidence for 125 GeV state and its identification as Euclidian pion together with the evidence for galactic pionlike state with mass of 145 GeV identified as M_{89} , one has a nice support for the overall TGD based view about situation described in this chapter. The small splitting between pionlike states has possible counterpart in the ordinary hadron physics: there is evidence for satellites of pion, mesons, and baryons in 20-40 MeV scale for mass splittings and in TGD framework they would correspond to IR Regge trajectories with the scale of 10-20 GeV mass splittings (see this chapter).

We are living exciting times!

3.4.5 Two different lifetimes for neutron as evidence for dark protons

I found a popular article (see <http://tinyurl.com/hqsaqok>) about very interesting finding related to neutron lifetime [C36] (see <http://tinyurl.com/h88n57j>). Neutron lifetime turns out to be by about 8 seconds shorter when measured by looking what fraction of neutrons disappears via decays in a box than by measuring the number of protons produced in beta decays for a neutron beam travelling through a given volume. The lifetime of neutron is about 15 minutes so that relative lifetime difference is about $8/(15 \times 60) \simeq .8$ per cent. The statistical significance is 4 sigma: 5 sigma is accepted as the significance for a finding acceptable as discovery.

How could one explain the finding? The difference between the methods is that the beam experiment measures only the disappearances of neutrons via beta decays producing protons whereas box measurement detects the outcome from all possible decay modes. The experiment suggests two alternative explanations.

1. Neutron has some other decay mode or modes, which are not detected in the box method since one measures the number of neutrons in initial and final state. For instance, in TGD framework one could think that the neutrons can transform to dark neutrons with some rate. But it is extremely improbable that the rate could be just about 1 per cent of the decay rate. Why not 1 millionth? Beta decay should be involved with the process.

Could some fraction of neutrons decay to dark proton, electron, and neutrino so that dark protons would not be detected in beam experiment? No, if one takes seriously the basic assumption of TGD that particles with different value of $h_{eff}/h = n$ do not appear in the same vertex. Neutron should first transform to dark proton but then also the disappearance could take place also without the beta decay of dark proton and the discrepancy would be larger.

2. The proton produced in the ordinary beta decay of neutron can however transform to dark proton not detected in the beam experiment! This would automatically predict that the rate is some reasonable fraction of the beta decay rate. About 1 percent of the resulting protons would transform to dark protons. This makes sense and would give strong hold about the rate for ordinary-dark transition rate. The observation of decays of neutron to electron, and neutrino but not proton would provide a support for the hypothesis. Both neutrino and proton would represent missing mass!

Dark matter as hierarchy of phases of ordinary matter is now a basic prediction of adelic TGD and $h_{eff}/h = n$ has in terms of number theory and space-time topology [K21, K19]. What is so nice is that the transformation of protons to dark protons is indeed the basic mechanism of TGD inspired quantum biology [K20, K19]! For instance, it would occur in Pollack effect [?] in with infrared irradiation of water bounded by gel phase generates so called exclusion zone (EZ), which is negatively charged, and creates what Pollack calls fourth phase of water. TGD explanation is that some fraction of protons transforms to dark protons at magnetic flux tubes outside the system. Negative charge of DNA and cell could be due to this mechanism. One also ends up to a model of genetic code with the analogs of DNA, RNA, tRNA and amino-acids represented as triplets of dark protons associated with magnetic flux tubes parallel to DNA strands [K17] [L4]. The model predicts correctly the numbers of DNAs coding a given amino-acid. Besides quantum biology, the model has applications to cold fusion [L3], and various phenomena referred to as “free energy phenomena” [K18, K2].

3.5 Has Pamela Observed Evidence For The Non-Dark Electro-PionOf M_{89} Lepto-Hadron Physics?

Resonaances (see <http://tinyurl.com/ybtqfgfn1> tells that the Fermi collaboration confirms the claim of PAMELA collaboration about anomalous e^+e^- pairs in cosmic ray radiation (see that abstract *Consistency of fermi-lat and pameala cosmic ray lepton measurement* by P. Grandi et al at (see <http://tinyurl.com/y7mc59gx>).

The announcement of Pamela was my second birthday gift at October 30 for two and half years ago. The first gift was CDF anomaly which found a beautiful explanation in terms of tau-pions and the p-adically scaled up variants with color tau- lepton having mass scale by power of two. The tau-pion of mass about 14 GeV decaying in cascade like manner to lower octaves of basic tau-pion explained elegantly the observations reported by CDF.

For some time ago the dilemma posed by the contradictory claims of DAMA and Cogent collaborations on one hand and XENON100 collaboration on one hand finds also nice solution in terms of 14 GeV taupion decaying to charged taupions with mass about 7 GeV [K16].

The decays of electro-pions to gamma pair can explain the observed anomalous gammas from galactic nucleus with energy very nearly to electron rest mass. Could one understand also the anomalous positrons reported by PAMELA as decay products of lepto-pion like states, say tau-pions? Intriguingly, the first figures of the article by Alessandro Strumia (see <http://tinyurl.com/y9jp333p>) [C5] discussing the constraints on the possible explanations of the PAMELA anomaly show that the anomalous positron excess starts around 10 GeV, possible it starts already at 7 GeV. It is not possible to say anything certain below 10 GeV since the measurements are affected by the solar activity below 10 GeV. What is however clear is that the excess cannot be

explained by taupion decays with 14 GeV mass since the excess would be localized around energy of about 7 GeV. Higher mass is required.

The article by Alessandro Strumia summarizes various theoretical constraints on the new particle explaining positron and electron excesses. The conclusions are following.

1. DM should result in a decay of quite a narrow particle with a mass very near to $2M$, which is nearly at rest. What narrow means quantitatively is not clear to me.
2. DM should carry a charge mediating long range interaction with the mediating boson which is must lighter than the particle itself: photon is the obvious candidate. Electromagnetically charged dark matter is however in conflict with the standard prejudices about dark matter and actually in dramatic conflict with its basic property of being invisible. Hierarchy of Planck constants is the only solution to the paradox of charged invisible dark matter.
3. DM must prefer the decays to leptons since otherwise there would be also antiproton and proton excess which has not been observed.
4. The mass of DM should be above 100 GeV.

These conditions encourage the identification of DM as a decay product of lepto-pion like state but with mass considerably higher than the 14 GeV mass. Tau-pions could of course be present but would not contribute to the anomaly at energies not too much above 7 GeV. Tau-pions would also give muon pair anomaly. Heavier lepto-pion like states are required and electro-pion would be the most natural candidate.

1. If a scaled up variant of ordinary hadron physics characterized by M_{89} is there as the recent bumps having interpretation as mesons of this physics suggest, there is no deep reason preventing the presence of also the scale variant of lepto-hadron physics in this scale. Even more, one can argue that colored leptons must appear as both dark and ordinary variants. Dark variants with non-standard value of Planck constant can have masses of ordinary leptons plus possibly their octaves as in the case of tau at least. The decay widths of intermediate gauge bosons require ordinary colored leptons to have mass higher than 45 GeV.
2. The mass of scaled up electro-pion would be obtained by scaling the mass of the dark electro-pion which for M_{89} electro-pion physics is in a good approximation $2m_e=1$ MeV by a factor $2^{(127-89)/2} = 2^{19}$. This gives electro-pion mass equal to 500 GeV. Ordinary colored electron would therefore have mass of 250 GeV consistent with the lower bound. The conclusion would be rather ironic: we would have seen dark colored electron (in TGD sense) already at seventies and covered it carefully under the rug and would be seeing now the ordinary colored electron and stubbornly trying to identify it as DM without caring about the fact that if dark matter is invisible in the standard sense it cannot be electromagnetically charged!
3. By stretching one's imagination one might play with the thought that superpartners of colored leptons with mass scale of order 100 GeV could form pion like states. The superpartners decay to partner and neutrino since R-parity is not exact invariance in TGD and all depends on how fast this process occurs.
4. Skeptic could wonder why the counterparts for colored excitations of quarks are not there and induce the increase of proton and antiproton fluxes.

To summarize, entire Zoo of not only new particles but even of new physics could be waiting for us at LHC energies if we live in TGD Universe!

3.6 Positron anomaly nine years later

The old PAMELA experiment [C15] (see <http://tinyurl.com/y7y9maac>) and perhaps newer ones by Fermi-LAT and AMS-02 have discovered lots of positrons in the cosmic rays, whose flux is generally higher than expected. The energies of positrons show steady rise in the range [10,100] GeV and presumably the rise will continue. Such positrons may originate from dark matter and could amount to an “almost direct detection” of the particles that make up dark matter. There are also other interpretations.

3.6.1 Dark matter explanations for the positron excess

Consider first new physics explanations postulating dark matter.

1. Dark spin 1 particles could decay to electron positron pair. The energy spectrum of energies is however discrete for dominating decay modes. For instance, vector mesons of new hadron physics could produce these events. Many neutral vector mesons say (Ψ/J) were discovered in electron-positron annihilation.
2. Pion-like spin 0 pseudoscalars decaying to electron-positron pairs and gamma rays predicts continuous spectrum. In the case of ordinary pion most decays are to gamma pairs. The decay to electron-positron pair and gamma ray has quite reasonable branching ratio .01. The reason is that the diagram describing this process is diagram for the decay to gamma pair with second gamma decaying to e^+e^- so that the rate is roughly the rate for the decay to gamma pair multiplied by $\alpha_{em} \sim 1/137$. This relation is expected to hold true for the decays of all pion-like states. For the decay to electron positron pair branching ratio about 6.5×10^{-8} . For pion-like states X the decay $X \rightarrow e^+e^- + \gamma$ for pion-like state could give a continuous spectrum. The mass of X should be of the order of 100 GeV for this option.

3.6.2 Standard physics explanation for the positron excess

One of the standard physics explanations is that the positrons emerge from pulsars. The beams from pulsars contain electrons accelerated to very high energies in the gigantic magnetic field of pulsar. This beam collides with the matter surrounding the pulsar and both gamma rays and positrons are generated in these interactions.

The standard physics proposal has been put to a test. One can predict the intensity of gamma rays coming from pulsars using standard model physics and deduce from it the density of electrons needed to generate it. Both positrons and gamma rays would be created when electrons from the pulsar are accelerated to very high energies in enormous magnetic field of the pulsar and collide with surrounding matter. This is like particle accelerator. The energies of the produced gamma rays and also positrons extend to TeV range, which corresponds to the energy range for LHC.

It turns out that the flux of electrons implied by the gamma ray intensity is too low to explain the flux of positrons detected by PAMELA [C15] and some other experiments. See the popular article at <http://tinyurl.com/ycw6xjs6> and the research article "Extended gamma-ray sources around pulsars constrain the origin of the positron flux at Earth" in Science [C18] (see <http://tinyurl.com/yach5g83>).

3.6.3 TGD based model for positron excess

Also TGD suggests an explanation for the positron excess (I learned about PAMELA experiment at my birth day nine years ago and it was an excellent birthday present!). TGD allows a hierarchy of scaled up copies of hadron physics labelled by ordinary Mersenne primes $M_n = 2^n - 1$ or by Gaussian Mersennes $M_{G,n} = (1 + i)^n - 1$. Ordinary hadron physics would correspond to M_{107} .

1. M_{89} hadron physics would have mass scale which is 512 times higher than that for ordinary hadron physics: the size scale of these hadrons is by factor 1/512 shorter than that for ordinary hadrons. There are indications for the copies also in other scales: M_{79} for instance [K10]. X boson provides indication for $M_{G,113}$ pions in nuclear scale. Even copies of hadron physics in biologically important length scales labelled by Gaussian Mersennes $M_{G,k}$, $k = 151, 157, 163, 167$ could exist and play key role in living matter [K8]. By the way, the appearance of four Gaussian Mersennes in this length scale range is number theoretical miracle.
2. M_{89} hadrons can also appear as dark states with Planck constant $h_{eff} = n \times h$. For $n = 512$ they would have the size of ordinary hadrons. This could explain the strange anomalies observed at RHIC and later at LHC and hinting about the presence of string like structures in what was expected to be color deconfinement phase transition predicting thermal spectrum should have been observed instead of strong correlations suggesting for quantum criticality characterized by long range correlations and fluctuations for which $h_{eff}/h = n$ would be an explanation.

3. A large number of bumps, whose masses correspond to the masses of ordinary hadron physics scaled up by factor 512, have been reported at LHC. Unfortunately these bumps cannot be explained by SUSY and other main stream models so that they have been forgotten.

TGD based model could be combined with the pulsar model for the positron excess. The collisions between protons from the pulsar accelerated in its magnetic field and the matter surrounding the pulsar would be analogous to those taking place between proton beams at LHC. If the collision energy is high enough (as it seems since gamma rays up to TeV range have been observed) they could produce dark M_{89} mesons, in particular pions, which then decay to gamma rays and lepton pairs, in particular electron-positron pairs. Similar collisions could occur also in the atmosphere of Earth between ultrahigh energy cosmic rays and nuclei of atmosphere and be responsible for the exotic cosmic ray events like Centauro challenging standard model physics [K11].

3.6.4 Other evidence for dark pion like states

There is also other evidence for pion-like states dark in TGD sense.

1. There is an old observation that gamma ray pairs with energy essential that of electrons rest must come from the center of Milky Way presumably resulting in decays of a particle with mass slightly larger than two times the mass of electron. These particles would also decay to electron positron pairs and the resulting electrons and positrons would be accelerated in the magnetic field of say pulsar to high energies. The rate for the decay to electron positron pairs is quite too slow as compared to the decay rate to gamma pairs. Therefore this mechanism cannot explain positron surplus.
2. The TGD model for the pion-like states decaying to gamma pairs is as leptopion [K16], which would be a pion-like bound state of color excitations of electrons predicted to be possible in TGD Universe. “Electropion” like states were discovered experimentally in CERN already at seventies and later evidence also for the muopions and taupions has emerged but since they did not fit with standard model, their existence was forgotten. This has been the fate of many other anomalies in particle physics. In nuclear physics there are century old forgotten anomalies re-discovered several times only to be “forgotten” again. The laws of Nature are not discovered nowadays as in good old days: they are decided by the hegemony, which happens to be in power. SUSY, superstring models, and M-theory already disappearing in the sands of time are basic examples of this new political science.
3. The reason for not accepting the existence of leptopion like states was that in standard model intermediate bosons should decay to them and their decay widths would be larger than their experimental values. However, if leptopions are dark matter in TGD sense having non-standard value of Planck constant $h_{eff}/h = n$, the problem can be circumvented.

3.7 Could Lepto-Hadrons Be Replaced With Bound States Of Exotic Quarks?

Can one then exclude the possibility that electron-hadrons correspond to colored quarks condensed around $k = 127$ hadronic space-time sheet: that is M_{127} hadron physics? There are several objections against this identification.

1. The recent empirical evidence for the colored counterpart of μ and τ supports the view that colored excitations of leptons are in question.
2. The octet character of color representation makes possible the mixing of leptons with lepton-baryons of form $L\nu_L\bar{\nu}_L$ by color magnetic coupling between leptogluons and ordinary and colored lepton. This is essential for understanding the production of electron-positron pairs.
3. In the case CDF anomaly also the assumption that colored variant of τ neutrino is very light is essential. In the case of colored quarks this assumption is not natural.

3.8 About The Masses Of Lepto-Hadrons

The progress made in understanding of dark matter hierarchy [K6] and non-perturbative aspects of hadron physics [K13, K10] allow to sharpen also the model of lepto-hadrons.

The model for the masses of ordinary hadrons [K13] applies also to the scaled up variants of the hadron physics. The two contributions to the hadron mass correspond to quark contribution and a contribution from super-symplectic bosons. For quarks labeled identical p-adic primes mass squared is additive and for quarks labeled by different primes mass is additive. Quark contribution is calculable once the p-adic primes of quarks are fixed.

super-symplectic contribution comes from super-symplectic bosons at hadronic space-time sheet labeled by Mersenne prime and is universal if one assumes that the topological mixing of the super-symplectic bosons is universal. If this mixing is same as for U type quarks, hadron masses can be reproduced in an excellent approximation if the super-symplectic boson content of hadron is assumed to correlate with the net spin of quarks.

In the case of baryons and pion and kaon one must assume the presence of a negative color conformal weight characterizing color binding. The value of this conformal weight is same for all baryons and super-symplectic contribution dominates over quark contribution for nucleons. In the case of mesons binding conformal weight can be assumed to vanish for mesons heavier than kaon and one can regard pion and kaon as Golstone bosons in the sense that quark contribution gives the mass of the meson.

This picture generalizes to the case of lepto-hadrons.

1. By the additivity of the mass squared leptonic contribution to lepto-pion mass would be $\sqrt{2}m_e(k)$, where k characterizes the p-adic length scale of colored electron. For $k = 127$ the mass of lepto-pion would be .702 MeV and too small. For $k = 126$ the mass would be $2m_e = 1.02$ MeV and is very near to the mass of the lepto-pion. Note that for ordinary hadrons quarks can appear in several scaled up variants inside hadrons and the value of k depends on hadron. The prediction for the mass of lepto ρ would be $m_{\pi_L} + \sqrt{7}m_{127} \simeq 1.62$ MeV ($m_{127} = m_e/\sqrt{5}$).
2. The state consisting of three colored electrons would correspond to leptonic variant of Δ_{++} having charge $q = -3$. The quark contribution to the mass of $\Delta_L \equiv \Delta_{L,3-}$ would be by the additivity of mass squared $\sqrt{3} \times m_e(k = 126) = 1.25$ MeV. If super-symplectic particle content is same as for Δ_L , super-symplectic contribution would be $m_{SC} = 5 \times m_{127}$, and equal to $m_{SC} = .765$ MeV so that the mass of Δ_L would be $m_{\Delta_L} = 2.34$ MeV. If colored neutrino corresponds to the same p-adic prime as colored electron, also lepto-proton has mass in MeV scale.

3.9 Do X And Y Mesons Provide Evidence For Color Excited Quarks Or Squarks?

Now and then come the days when head is completely empty of ideas. One just walks around and gets more and more frustrated. One can of course make authoritative appearances in blog groups and express strong opinions but sooner or later one is forced to look for web if one could find some problem. At this time I had good luck. By some kind of divine guidance I found myself immediately in Quantum Diaries and found a blog posting with title *Who ordered that?! An X-traordinary particle?* (see <http://tinyurl.com/3k9pts5>) [C3].

Not too many unified theorists take meson spectroscopy seriously. Although they are now accepting low energy phenomenology (*the physics for the rest of us*) as something to be taken seriously, meson physics is for them a totally uninteresting branch of botany. They could not care less. As a crackpot I am however not well-informed about what good theoretician should do and shouldn't do and got interested. Could this give me a problem that my poor crackpot brain is crying for?

The posting told me that in the spectroscopy of $c\bar{c}$ type mesons is understood except for some troublesome mesons christened imaginatively with letters X and Y plus brackets containing their mass in MeVs. $X(3872)$ is the firstly discovered troublemaker and what is known about it can be found in the blog posting and also in Particle Data Tables (see <http://tinyurl.com/y7x23br5>) [C2]. The problem is that these mesons should not be there. Their decay widths seem to be

narrow taking into account their mass and their decay characteristics are strange: in particular the kinematically allow decays to DD dominating the decays of $\Psi(3770)$ with branching ratio 93 per cent has not been observed whereas the decay to $DD\pi^0$ occurs with a branching fraction $> 3.2 \times 10^{-3}$. Why the pion is needed? $X(3872)$ should decay to photon and charmonium state in a predictable way but it does not.

3.9.1 Could these be the good questions?

TGD predicts a lot of exotic physics and I of course started to exclude various alternatives. First one must however try to invent a good question. Maybe the following questions might satisfy the criterion of goodness.

1. Why these exotic states appear only for mesons made of heavy quark and antiquark? Why not for light mesons? Why not for mesons containing one heavy quark and light quark? Could it be that also $b\bar{b}$ mesons could have exotic partners not yet detected? Could it be that also exotic $b\bar{c}$ type mesons could be there? Why the presence of light quark would eliminate the exotic partner from the spectrum?
2. Do the decays obey some selection rules? There is indeed this kind of rule: the numbers of c and \bar{c} quarks in the final state are equal to one.
 - (a) If c and \bar{c} exist in the initial state and the decay involves only strong interactions, the rule holds true.
 - (b) If c and \bar{c} are not present in the initial state the only option that one can imagine is the exchange of two W bosons transforming d type quarks to c type quarks must be present. If this were the case the initial state should correspond to $d\bar{d}$ like state rather than $c\bar{c}$ and this looks very strange from the standard physics point of view. Also the rate for this kind of decays would be very small and it seems that this option cannot make sense.

3.9.2 Both leptons and quarks have color excitations in TGD Universe

TGD predicts that both leptons and quarks have color excitations [K16]. For leptons they correspond to color octets and there is a lot of experimental evidence for them. Why we do not have any evidence for color excited quarks? Or do we actually have?! Could these strange X : s and Y : s provide this evidence?

Ordinary quarks correspond to triality one color triplet partial waves in CP_2 . The higher color partial waves would also correspond to triality one states but in higher color partial waves in CP_2 . The representations of the color group are labelled by two integers (p, q) and the dimension of the representation is given by

$$d = \frac{(p+1)(q+1)(p+q+2)}{2} .$$

A given $t = \pm 1$ representation is accompanied by its conjugate with the same dimension and opposite triality $t = \mp 1$. $t = 1$ representations satisfy $p - q = 1$ modulo 3 and come as $(1, 0)$, $(0, 2)$, $(3, 0)$, $(2, 1)$, with dimensions 3, 6, 10, 15, ... The simplest candidate for the color excitations would correspond to the representation $\bar{6}$. It does not correspond directly the a solution of the Dirac equation in CP_2 since physical states involve also color Kac-Moody generators [K9].

Some remarks are in order:

1. The tensor product of gluon octet with $t = 1$ with color triplet representation contains $8 \times 3 = 24$ states and decomposes into $t = 1$ representations as $3 \oplus \bar{6} \oplus 15$. The coupling of gluons by Lie algebra action can couple given representation only with itself. The coupling between triplet and $\bar{6}$ and 15 is therefore not by Lie algebra action. The coupling constant between quarks and color excited quarks is *assumed* to be proportional to color coupling.
2. The existence of this kind of coupling would explain the selection rules elegantly. If this kind of coupling is not allowed then only the annihilation of exotic quark to gluon decaying to quark pair can transform exotic mesons to ordinary ones and I have not been able to explain selection rules using this option.

The basic constraint applying to all variants based on exotic states of quarks comes from the fact that the decay widths of intermediate gauge bosons do not allow new light particles. This objection is encountered already in the model of lepto-hadrons [K16]. The solution is that the light exotic states are possible only if they are dark in TGD sense having therefore non-standard value of Planck constant and behaving as dark matter. The value of Planck constant is only effective and has purely geometric interpretation in TGD framework. This implies that a phase transition transforming quarks and gluons to their dark counterparts is the key element of the model. After this a phase transition a gluon exchange would transform the quark pair to an exotic quark pair.

3.9.3 Also squarks could explain exotic charmonium states

Supersymmetry provides an alternative mechanism. Right-handed neutrino generates super-symmetries in TGD Universe and quarks are accompanied by squarks consisting in a well-defined sense of quark and right-handed neutrino. Super-symmetry would allow completely standard couplings to gluons by adding to the spectrum squarks and gluinos. Exactly the same selection rules result if these new states are mesonlike states from squark and anti-squark and the exchange of gluino after the \hbar changing phase transition transforms exotic meson to ordinary one and vice versa.

In the sequel it will be shown that the existence of color excited quarks or of their superpartners could indeed allow to understand the origin of X and Y mesons and also the absence of analogous states accompanying mesons containing light quarks or antiquarks.

This picture would lead to a completely new view about detection of squarks and gluinos.

1. In the standard scenario the basic processes are production of squark and gluino pair. The creation of squark-antisquark pair is followed by the decay of squark (anti-squark) to quark (antiquark) and neutralino or chargino. If R-parity is conserved, the decay chain eventually gives rise to at least two hadron jets and lightest neutralinos identifiable as missing energy. Gluinos in turn decay to quark and anti-squark (squark and antiquark) and squark (anti-squark) in turn to quark (anti-quark) and neutralino or chargino. At least four hadron jets and missing energy is produced. In TGD framework neutralinos would decay eventually to zinos or photinos and right-handed neutrino transforming to ordinary neutrino (R-parity is not conserved). This process might be however slow.
2. In the recent case quite different scenario relying on color confinement and “shadronization” suggests itself. By definition smesons consist of squarks and antisquark. Sbaryons could consist of two squarks containing right-handed neutrino and its antineutrino ($\mathcal{N} = 2$ SUSY) and one quark and thus have same quantum numbers as baryon. Note that the squarks are dark in TGD sense.

Also now dark squark or gluino pair would be produced at the first step and would require \hbar changing phase transition of gluon. These would shadronize to form a dark shadron. One can indeed argue that the required emission of winos and zinos and photinos is too slow a process as compared to shadronization. Shadrons (mostly smesons) would in turn decay to hadrons by the exchange of gluinos between squarks. No neutralinos (missing energy) would be produced. This would explain the failure to detect squarks and gluinos at LHC.

This mechanism does not however apply to sleptons so that it seems that the p-adic mass scale of sleptons must be much higher for sleptons than that for squarks as I have indeed proposed.

3.9.4 Could exotic charmonium states consist of color excited c and \bar{c} or of their spartners?

Could one provide answers to the questions presented in the beginning assuming that exotic charmonium states consists of dark color excited c and \bar{c} : or more generally, a mixture of ordinary charmonium and exotic charmonium state? The mixing is expected since \hbar changing phase transition followed by a gluon exchange can transform these meson states to each other. Also annihilation to gluon and back to quark pair can induce this mixing. The mixing is however small for heavy quarks for which $\alpha_s \simeq .1$ holds true. Exactly the same arguments apply to the meson like bound states of squarks and in the following only the first option will be discussed.

1. In the case of charged leptons colored excitations have have same p-adic mass scale: for τ however several p-adic mass scales appear as the model if the two year old CDF anomaly is taken seriously [K16]. Assume that p-adic mass scales - but not necessarily masses- are the same also now. This assumption might be non-sensical since also light mesons would have exotic counterparts and somehow they should disappear from the spectrum. To simplify the estimates one could even assume even that the masses are same.
2. In the presence of small mixing the decay amplitude would come solely from the small contribution of the ordinary $c\bar{c}$ state present in the state dominated by color excited pair. The two manners to see the situation should give essentially the same answer.
3. The decays would take place via strong interactions.

The challenge is to understand why the dominating decays to $D\bar{D}$ with branching fraction of 93 per cent are not allowed whereas $D\bar{D}\pi^0$ takes place. Why the pion is needed? The second challenge is to understand why X does not decay to charmonium and photon.

1. For ordinary charmonium the decay to $D\bar{D}$ could take place by the emission of gluon from either c or \bar{c} which then decays to light quark pair whose members combine with c and \bar{c} to form D and \bar{D} . Now this mechanism does not work. At least *two* gluons must be emitted to transform colored excited $c\bar{c}$ to ordinary $c\bar{c}$. If these gluons decay to light quark pairs one indeed obtains an additional pion in hadronization. The emission of two gluons instead of only one is expected to reduce the rate roughly by $\alpha_s^2 \simeq 10^{-2}$ factor.
2. Also ordinary decays are predicted to occur but with a slower rate. The first step would be an exchange of gluon transforming color excited charmed quark pair to an ordinary charmed quark pair. After the transformation to off mass shell $c\bar{c}$ pair, the only difference to the decays of charmonium states would be due to the fact that charmonium would be replaced with $c\bar{c}$ pair. The exchange of the gluon preceding this step could reduce the decay rate with respect to charmonium decay rates by a factor of order $\alpha_s^2 \simeq 10^{-2}$. Therefore also the ordinary decay modes should be there but with a considerably reduced rate.
3. Why the direct decays to photon and charmonium state do not occur in the manner predicted by the model of charmonium? For ordinary charmonium the decay proceeds by an emission of photon by either quark or antiquark. Same mechanism applies for exotic charmonium states but leads to final state which consists of *exotic* charmonium and photon. In the case of $X(3872)$ there exists no lighter exotic charmonium state so that the decay is forbidden in this order of perturbation theory. Heavier exotic charmonium states can however decay to photon plus exotic charmonium state in this order of perturbation theory if discrete symmetries favor this.

Essentially identical arguments go through if c and \bar{c} are replaced with their dark partners and exchange of gluon by the emission of gluino. The transformation of gluon to its dark variants is an essential element in the process.

3.9.5 Why the color excitations/spartners of light quarks would be effectively absent?

Can one understand the effective absence of mesons consisting of color excited light quarks or squarks if the excitations have same mass scale and even mass as the light quarks? The following arguments are for color excited quarks but they apply also to squarks.

1. Suppose that the mixing induced by \hbar changing phase transition followed by a gluon exchange and annihilation is described by mass squared matrix containing besides diagonal components $M_1^2 = M_2^2$ also non-diagonal component $M_{12}^2 = \overline{M_{21}^2}$. The eigenstates of the mass squared matrix correspond to the physical states which are mixtures of states consisting of ordinary quark pair and pair of color excited quarks. The non-diagonal elements of the mass squared matrix corresponds to gluon exchange and since color interactions get very strong at low energy scales, one expects that these elements get very large. In the degenerate case $M_1^2 = M_2^2$ the mass squared eigen values are given by

$$M_{\pm}^2 = M_0^2 \pm |M_{12}|^2 . \quad (3.4)$$

2. Suppose that $M_0^2 = 0$ holds true in accordance with approximate pseudo Goldstone nature of pion and more generally all light pseudo-scalar mesons. In fact assume that this is the case before color magnetic spin-spin splitting has taken place so that in this approximation pion and ρ would have same mass $m_{\pi}^2 = m_{\rho}^2 = M_0^2$. In TGD based model for color magnetic spin-spin splitting M_0^2 energy is replaced with mass squared [K13] and M_0^2 is obtained in terms of physical masses of π and ρ from the basic formulas

$$\begin{aligned} m_{\pi}^2 &= M_0^2 - \frac{1}{4}\Delta , \quad m_{\rho}^2 = M_0^2 + \frac{3}{4}\Delta , \\ M_0^2 &= \frac{m_{\rho}^2 + 3m_{\pi}^2}{2} , \quad \Delta = m_{\rho}^2 - m_{\pi}^2 . \end{aligned} \quad (3.5)$$

The exotic π and ρ would have masses

$$\begin{aligned} m_{\pi_{ex}}^2 &= -M_0^2 - \frac{1}{4}\Delta = m_{\pi}^2 - 2M_0^2 , \\ m_{\rho_{ex}}^2 &= -M_0^2 + \frac{3}{4}\Delta = m_{\rho}^2 - 2M_0^2 . \end{aligned} \quad (3.6)$$

For $m_{\pi} = 140 \text{ MeV}$ and $m_{\rho} = 770 \text{ MeV}$ the calculation gives $m_{\pi_{ex}} = i \times 685 \text{ MeV}$ so a tachyon would be in question. For ρ one would have $m_{\rho_{ex}} = 323 \text{ MeV}$ so that the mass would not be tachyonic.

One can try to improve the situation by allowing $M_1^2 \neq M_2^2$ giving additional flexibility and hopes about tachyonicity of the exotic ρ .

1. In this case one obtains the equations

$$\begin{aligned} m_{\pi}^2 &= M_+^2 - \frac{1}{4}\Delta , \quad m_{\rho}^2 = M_+^2 + \frac{3}{4}\Delta \\ m_{\pi_{ex}}^2 &= M_-^2 - \frac{1}{4}\Delta , \quad m_{\rho_{ex}}^2 = M_-^2 + \frac{3}{4}\Delta , \\ M_+^2 &= \frac{M_1^2 + M_2^2}{2} + \sqrt{\left(\frac{M_1^2 + M_2^2}{2}\right)^2 + M_{12}^4} = \frac{m_{\rho}^2 + 3m_{\pi}^2}{2} , \\ M_-^2 &= \frac{M_1^2 + M_2^2}{2} - \sqrt{\left(\frac{M_1^2 + M_2^2}{2}\right)^2 + M_{12}^4} = M_+^2 - 2\sqrt{\left(\frac{M_1^2 + M_2^2}{2}\right)^2 + M_{12}^4} . \end{aligned} \quad (3.7)$$

2. The condition that ρ_{ex} is tachyonic gives

$$m_{\rho_{ex}}^2 = M_-^2 + \frac{3}{4}\Delta < 0 , \quad (3.8)$$

giving

$$\begin{aligned} m_{\rho}^2 &< 2\sqrt{\left(\frac{M_1^2 + M_2^2}{2}\right)^2 + M_{12}^4} , \\ M_+^2 &= \frac{M_1^2 + M_2^2}{2} + \sqrt{\left(\frac{M_1^2 + M_2^2}{2}\right)^2 + M_{12}^4} = \frac{m_{\rho}^2 + 3m_{\pi}^2}{2} , \end{aligned} \quad (3.9)$$

3. In the parameterization $(m_1^2, m_2^2, M_{12}^2) = (x, y, z)m_\rho^2$ one obtains the conditions

$$D \equiv \sqrt{(x+y)^2 + z^2} > 1/2 ,$$

$$\frac{x+y}{2} + D = \frac{1}{2} + \frac{3}{2} \frac{m_\pi^2}{m_\rho^2} . \quad (3.10)$$

4. These equations imply the conditions

$$x+y < 3 \frac{m_\pi^2}{m_\rho^2} \simeq .099 ,$$

$$.490 < z < .599 . \quad (3.11)$$

The first condition implies $\sqrt{m_1^2 + m_2^2} < 242.7$ MeV. Second condition gives $339 < M_{12}/MeV < 595.9$ so that rather stringent bounds on the parameters are obtained. The simplest solution to the conditions corresponds to $x = y = 0$ and $z = .599$. This solution would mean vanishing masses in the absence of mixing and spin-spin splitting and could be defended by the Golstone boson property of pions mass degenerate with ρ mesons.

This little calculation encourages to consider the possibility that all exotic counterparts of light mesons are tachyonic and that this due the very large mixing induced by gluon exchange (gluon exchange squark option) at low energies. It would be nice if also mesons containing only single heavy quark were tachyonic and this could be the case if the p-adic length scale defining the strength of color interactions corresponds to that of the light quark so that the mass matrix has large enough non-diagonal component. Here one must be however very cautious since experimental situation is far from clear.

The model suggests that ordinary charmonium states and their exotic partners are in 1-1 correspondence. If so then many new exotic states are waiting to be discovered.

3.9.6 The option based on heavy color excitations/spartners of light quarks

An alternative option is that color excitations/spartners of light quarks have large mass: this mass should not be however larger than the mass of c quarks if we want to explain X : s and Y : s as pairs of color excitations of light quarks. Suppose that the p-adic mass scale is same as that for c quarks or near it (not that the scales come as powers of $\sqrt{2}$). This raises the question whether exotic $c\bar{c}$ mesons really consist of exotic c and \bar{c} : why not color excitations of u, d, s and their antiquarks? As a matter fact, we cannot be sure about the quark content of X and Y mesons. Could these states be $d\bar{d}$ and $u\bar{u}$ states for their color excitations? It however seems that the presence of two W exchanges makes the decay rate quite too low so that this option seems to be out of question.

One can however consider the option in which the squarks associated with light quarks are heavy. This option is indeed realized in standard SUSY were the mass scales of particles families are inverted so that stop and sbottom are the lightest squarks and super-partners of u and d the heaviest ones. This would predict that the smesons associated with \bar{t} and $b\bar{b}$ are lighter than X and Y (s)mesons. This option does not look at all natural in TGD but of course deserves experimental checking.

3.9.7 How to test the dark squark option?

The identification of X and Y as dark smesons looks like a viable option and explains the failure to find SUSY at LHC if shadronization is a fast process as compared to the selectro-weak decays. The option certainly deserves an experimental testing. One could learn a lot about SUSY in TGD sense (or maybe in some other sense!) by just carefully scanning the existing data at lower energies. For instance, one could try to answer the following questions by analyzing the already existing experimental data.

1. Are X and Y type mesons indeed in 1-1 correspondence with charmonium states? One could develop numerical models allowing to predict the precise masses of charmonium states and their decay rates to various final states and test the predictions experimentally.
2. Do $b\bar{b}$ mesons have smesonic counterparts with the same mass scale? What about B_c type smesons containing two heavy squarks?
3. Do the mesons containing one heavy quark and one light quark have smesonic counterparts? My light-hearted guess that this is not the case is based on the assumption that the general mass scale of the mass squared matrix is defined by the p-adic mass scale of the heavy quark and the non-diagonal elements are proportional to the color coupling strength at p-adic length scale associated with the light quark and therefore very large: as a consequence the second mass eigenstate would be tachyonic.
4. What implications the strong mixing of light mesons and smesons would have for CP breaking? CP breaking amplitudes would be superpositions of diagrams representing CP breaking for mesons *resp.* smesons. Could the presence of smesonic contributions perhaps shed light on the poorly understood aspects of CP breaking?

3.9.8 Objection against covariantly constant neutrinos as SUSY generators

TGD SUSY in its simplest form assumes that covariantly constant right-handed neutrino generates SUSY. The second purely TGD based element is that squarks would correspond to the same p-adic mass scale as partners.

This looks nice but there are objections.

1. The first objection relates to the tachyonicity needed to get rid of double degeneracy of light mesons consisting of u, d, and s quarks. Mesons and smesons consisting of squark pair mix and for large α_s the mixing is large and can indeed make second eigenvalue of the mass squared matrix negative. If so, these states disappear from spectrum. At least to me this looks however somewhat unaesthetic.

Luckily, the transformation of second pion-like state to tachyon and disappearance from spectrum is not the only possibility. After a painful search I found experimental work (see <http://tinyurl.com/ybq323yy>) [C50] claiming the existence of states analogous to ordinary pion with masses 60, 80, 100, 140, MeV. Also nucleons have this kind of satellite states. Could it be that one of these states is spion predicted by TGD SUSY for ordinary hadrons? But what about other states? They are not spartners: what are they?

2. The second objection relates to the missing energy. SUSY signatures involving missing energy have not been observed at LHC. This excludes standard SUSY candidates and could do the same in the case of TGD. In TGD framework the missing energy would be eventually right handed neutrinos resulting from the decays of sfermions to fermion and sneutrino in turn decaying to neutrino and right handed neutrino. The naive argument is that shadronization would be much faster process than the decay of squarks to quarks and spartners of electro-weak gauge bosons and missing energy so that these events would not be observed. Shadrons would in turn decay to hadrons by gluino exchanges. The problem with this argument is that the weak decays of squarks producing right handed neutrinos as missing energy are still there!

This objection forces to consider the possibility that covariantly constant right handed neutrino which generates SUSY is replaced with a color octet. Color excitations of leptons of lepto-hadron hypothesis [K16] would be sleptons which are color octets so that SUSY for leptons would have been seen already at seventies in the case of electron. The whole picture would be nicely unified. Sleptons and squark states would contain color octet right handed neutrino the same wormhole throats as their em charge resides. In the case of squarks the tensor product $3 \otimes 8 = 3 + \bar{6} + 15$ would give several colored exotics. Triplet squark would be like ordinary quark with respect to color.

Covariantly constant right-handed neutrino as such would represent pure gauge symmetry, a super-generator annihilating the physical states. Something very similar can occur in the

reduction of ordinary SUSY algebra to sub-algebra familiar in string model context. By color confinement missing energy realized as a color octet right handed neutrino could not be produced and one could overcome the basic objections against SUSY by LHC.

What about the claimed anomalous trilepton events at LHC interpreted in terms of SUSY, which however breaks either the conservation of lepton or baryon number. I have proposed TGD based interpretation [K10] in terms of the decays of W to \tilde{W} and \tilde{Z} , which in turn decay and produce the three lepton signature. Suppose that \tilde{W} and \tilde{Z} are color octets and that sleptons replace the color octet excitations of leptons responsible for lepto-hadron physics [K16]. One possible decay chain would involve the decays $\tilde{W}^+ \rightarrow \tilde{L}^+ + \bar{\nu}_L$ and $\tilde{Z} \rightarrow L^+ + \tilde{L}^-$. Color octet sleptons pair combine to form lepto-pion which decays to lepton pair. This decay cascade would produce missing energy as neutrino and this seems to be the case for other options too. One could overcome the basic objections against SUSY by LHC.

This view about TGD SUSY clearly represents a hybrid of the two alternative views about X and Y bosons as composites of either color excitations of quarks or of squarks and is just one possibility. The situation is not completely settled and one must keep mind open.

4 Appendix

4.1 Evaluation Of Lepto-Pion Production Amplitude

4.1.1 General form of the integral

The amplitude for lepto-pion production with four momentum

$$\begin{aligned} p &= (p_0, \vec{p}) = m\gamma_1(1, v\sin(\theta)\cos(\phi), v\sin(\theta)\sin(\phi), v\cos(\theta)) , \\ \gamma_1 &= 1/(1-v^2)^{1/2} , \end{aligned} \quad (4.1)$$

is essentially the Fourier component of the instanton density

$$U(b, p) = \int e^{ip \cdot x} E \cdot B d^4x \quad (4.2)$$

associated with the electromagnetic field of the colliding nuclei.

In order to avoid cumbersome numerical factors, it is convenient to introduce the amplitude $A(b, p)$ as

$$\begin{aligned} A(b, p) &= N_0 \times \frac{4\pi}{Z_1 Z_2 \alpha_{em}} \times U(b, p) , \\ N_0 &= \frac{(2\pi)^7}{i} \end{aligned} \quad (4.3)$$

Coordinates are chosen so that target nucleus is at rest at the origin of coordinates and colliding nucleus moves along positive z direction in $y = 0$ plane with velocity β . The orbit is approximated with straight line with impact parameter b .

Instanton density is just the scalar product of the static electric field E of the target nucleus and magnetic field B the magnetic field associated with the colliding nucleus, which is obtained by boosting the Coulomb field of static nucleus to velocity β . The flux lines of the magnetic field rotate around the direction of the velocity of the colliding nucleus so that instanton density is indeed non vanishing.

The Fourier transforms of E and B for nuclear charge 4π (chose for convenience) giving rise to Coulomb potential $1/r$ are given by the expressions

$$\begin{aligned} E_i(k) &= N\delta(k_0)k_i/k^2 , \\ B_i(k) &= N\delta(\gamma(k_0 - \beta k_z))k_j\epsilon_{ijz}e^{ik_x b}/((\frac{k_z}{\gamma})^2 + k_T^2) , \\ N &= \frac{1}{(2\pi)^2} . \end{aligned} \quad (4.4)$$

The normalization factor corresponds to momentum space integration measure d^4p . The Fourier transform of the instanton density can be expressed as a convolution of the Fourier transforms of E and B.

$$\begin{aligned} A(b, p) &\equiv = N_0 N_1 \int E(p-k) \cdot B(k) d^4k , \\ N_1 &= \frac{1}{(2\pi)^4} . \end{aligned} \quad (4.5)$$

Where the fields correspond to charges $\pm 4\pi$. In the convolution the presence of two delta functions makes it possible to integrate over k_0 and k_z and the expression for U reduces to a two-fold integral

$$\begin{aligned} A(b, p) &= \beta\gamma \int dk_x dk_y \exp(ik_x b) (k_x p_y - k_y p_x) / AB , \\ A &= (p_z - \frac{p_0}{\beta})^2 + p_T^2 + k_T^2 - 2k_T \cdot p_T \\ B &= k_T^2 + (\frac{p_0}{\beta\gamma})^2 , \\ p_T &= (p_x, p_y) . \end{aligned} \quad (4.6)$$

To carry out the remaining integrations one can apply residue calculus.

1. k_y integral is expressed as a sum of two pole contributions
2. k_x integral is expressed as a sum of two pole contributions plus two cut contributions.

4.1.2 k_y -integration

Integration over k_y can be performed by completing the integration contour along real axis to a half circle in upper half plane (see **Fig. ??**).

The poles of the integrand come from the two factors A and B in denominator and are given by the expressions

$$\begin{aligned} k_y^1 &= i(k_x^2 + (\frac{p_0}{\beta\gamma})^2)^{1/2} , \\ k_y^2 &= p_y + i((p_z - \frac{p_0}{\beta})^2 + p_x^2 + k_x^2 - 2p_x k_x)^{1/2} . \end{aligned} \quad (4.7)$$

One obtains for the amplitude an expression as a sum of two terms

$$A(b, p) = 2\pi i \int e^{ik_x b} (U_1 + U_2) dk_x , \quad (4.8)$$

corresponding to two poles in upper half plane.

The explicit expression for the first term is given by

$$\begin{aligned} U_1 &= RE_1 + iM_1 , \\ RE_1 &= (k_x \frac{p_0}{\beta} y - p_x r e_1 / 2) / (r e_1^2 + i m_1^2) , \\ IM_1 &= (-k_x p_y r e_1 / 2 K_1^{1/2} - p_x p_y K_1^{1/2}) / (r e_1^2 + i m_1^2) , \\ r e_1 &= (p_z - \frac{p_0}{\beta})^2 + p_T^2 - (\frac{p_0}{\beta\gamma})^2 - 2p_x k_x , \\ i m_1 &= -2K_1^{1/2} p_y , \\ K_1 &= k_x^2 + (\frac{p_0}{\beta\gamma})^2 . \end{aligned} \quad (4.9)$$

The expression for the second term is given by

$$\begin{aligned}
U_2 &= RE_2 + iIM_2 , \\
RE_2 &= -((k_x p_y - p_x p_y) p_y + p_x r e_2 / 2) / (r e_2^2 + i m_2^2) , \\
IM_2 &= -(k_x p_y - p_x p_y) r e_2 / 2 K_2^{1/2} + p_x p_y K_2^{1/2} / (r e_2^2 + i m_2^2) , \\
r e_2 &= -(p_z - \frac{p_0}{\beta})^2 + (\frac{p_0}{\beta \gamma})^2 + 2 p_x k_x + \frac{p_0}{\beta} y - \frac{p_0}{\beta} x , \\
i m_2 &= 2 p_y K_2^{1/2} , \\
K_2 &= (p_z - \frac{p_0}{\beta})^2 + \frac{p_0}{\beta} x + k_x^2 - 2 p_x k_x .
\end{aligned} \tag{4.10}$$

A little inspection shows that the real parts cancel each other: $RE_1 + RE_2 = 0$. A further useful result is the identity $i m_1^2 + r e_1^2 = r e_2^2 + i m_2^2$ and the identity $r e_2 = -r e_1 + 2 p_y^2$.

4.1.3 k_x -integration

One cannot perform k_x -integration completely using residue calculus. The reason is that the terms IM_1 and IM_2 have cuts in complex plane. One can however reduce the integral to a sum of pole terms plus integrals over the cuts.

The poles of U_1 and U_2 come from the denominators and are in fact common for the two integrands. The explicit expressions for the pole in upper half plane, where integrand converges exponentially are given by

$$\begin{aligned}
r e_i^2 + i m_i^2 &= 0 , \quad i = 1, 2 , \\
k_x &= (-b + i(-b^2 + 4ac)^{1/2}) / 2a , \\
a &= 4 p_T^2 , \\
b &= -4((p_z - \frac{p_0}{\beta})^2 + p_T^2 - (\frac{p_0}{\beta \gamma})^2) p_x \text{ per} , \\
c &= ((p_z - \frac{p_0}{\beta})^2 + p_T^2 - (\frac{p_0}{\beta \gamma})^2)^2 + 4(\frac{p_0}{\beta \gamma})^2 p_y^2 .
\end{aligned} \tag{4.11}$$

A straightforward calculation using the previous identities shows that the contributions of IM_1 and IM_2 at pole have opposite signs and the contribution from poles vanishes identically!

The cuts associated with U_1 and U_2 come from the square root terms K_1 and K_2 . The condition for the appearance of the cut is that K_1 (K_2) is real and positive. In case of K_1 this condition gives

$$k_x = it, \quad t \in (0, \frac{p_0}{\beta \gamma}) . \tag{4.12}$$

In case of K_2 the same condition gives

$$k_x = p_x + it, \quad t \in (0, \frac{p_0}{\beta} - p_z) . \tag{4.13}$$

Both cuts are in the direction of imaginary axis.

The integral over real axis can be completed to an integral over semi-circle and this integral in turn can be expressed as a sum of two terms (see **Fig. 4**).

$$A(b, p) = 2\pi i (CUT_1 + CUT_2) . \tag{4.14}$$

The first term corresponds to contour, which avoids the cuts and reduces to a sum of pole contributions. Second term corresponds to the addition of the cut contributions.

In the following we shall give the expressions of various terms in the region $\phi \in [0, \pi/2]$. Using the symmetries

$$\begin{aligned} A(b, p_x, -p_y) &= -A(b, p_x, p_y) , \\ A(b, -p_x, -p_y) &= \bar{A}(b, p_x, p_y) . \end{aligned} \quad (4.15)$$

of the amplitude one can calculate the amplitude for other values of ϕ .

The integration variable for cuts is the imaginary part t of complexified k_x . To get a more convenient form for cut integrals one can perform a change of the integration variable

$$\begin{aligned} \cos(\psi) &= \frac{t}{\left(\frac{p_0}{\beta\gamma}\right)} , \\ \cos(\psi) &= \frac{t}{\left(\frac{p_0}{\beta} - p_z\right)} , \\ \psi &\in [0, \pi/2] . \end{aligned} \quad (4.16)$$

1. The contribution of the first cut

By a painstaking calculation one verifies that the expression for the contribution of the first cut is given by

$$\begin{aligned} CUT_1 &= D_1 \times \int_0^{\pi/2} \exp\left(-\frac{b}{b_0} \cos(\psi)\right) A_1 d\psi , \\ D_1 &= -\frac{1}{2} \frac{\sin(\phi)}{\sin(\theta)} , \quad b_0 = \frac{\hbar \beta\gamma}{m \gamma_1} , \\ A_1 &= \frac{A + iB \cos(\psi)}{\cos^2(\psi) + 2iC \cos(\psi) + D} , \\ A &= \sin(\theta) \cos(\phi) , \quad B = K , \\ C &= K \frac{\cos(\phi)}{\sin(\theta)} , \quad D = -\sin^2(\phi) - \frac{K^2}{\sin^2(\theta)} , \\ K &= \beta\gamma \left(1 - \frac{v_{cm}}{\beta} \cos(\theta)\right) , \quad v_{cm} = \frac{2v}{1+v^2} . \end{aligned} \quad (4.17)$$

The definitions of the various kinematical variables are given in previous formulas. The notation is tailored to express that A_1 is rational function of $\cos(\psi)$.

1. The exponential $\exp(-b \cos(\psi)/b_0)$ is very small in the condition

$$\cos(\psi) \geq \cos(\psi_0) \equiv \frac{\hbar \beta\gamma}{mb \gamma_1 \cos(\phi)} \quad (4.18)$$

holds true. Here $\hbar = 1$ convention has been given up to make clear that the increase of the Compton length of lepto-pion due to the scaling of \hbar increase the magnitude of the contribution. If the condition $\cos(\psi_0) \ll 1$ holds true, the integral over ψ receives contributions only from narrow range of values near the upper boundary $\psi = \pi/2$ plus the contribution corresponding to the pole of X_1 . The practical condition is in terms of critical parameter b_{max} above which exponential approaches zero very rapidly.

2. For $\cos(\psi_0) \ll 1$, that is for $b > b_{max}$ and in the approximation that the function multiplying the exponent is replaced with its value for $\psi = \pi/2$, one obtains for CUT_1 the expression

$$\begin{aligned} CUT_1 &\simeq D_1 A_1(\psi = \pi/2) \frac{\hbar}{mb} \\ &= \frac{1}{2} \times \frac{\beta\gamma}{\gamma_1} \times \frac{\hbar}{mb} \times \frac{\sin^2(\theta) \cos(\phi) \sin(\phi)}{\sin^2(\theta) \sin^2(\phi) + K^2} . \end{aligned} \quad (4.19)$$

3. For $\cos(\psi_0) \gg 1$ exponential factor can be replaced by unity in good approximation and the integral reduces to an integral of rational function of $\cos(\psi)$ having the form

$$D_1 \frac{A + iB\cos(\psi)}{\cos^2(\psi) + 2iC \times \cos(\psi) + D} \quad (4.20)$$

which can be expressed in terms of the roots c_{\pm} of the denominator as

$$D_1 \times \sum_{\pm} \frac{A \mp iBc_{\pm}}{\cos(\psi) - c_{\pm}} , \quad c_{\pm} = -iC \pm \sqrt{-C^2 - D} \quad (4.21)$$

Integral reduces to an integral of rational function over the interval $[0, 1]$ by the standard substitution $\tan(\psi/2) = t$, $d\psi = 2dt/(1+t^2)$, $\cos(\psi) = (1-t^2)/(1+t^2)$, $\sin(\psi) = 2t/(1+t^2)$.

$$I = 2D_1 \sum_{\pm} \int_0^1 dt \frac{A \mp iBc_{\pm}}{1 - c_{\pm} - (1 + c_{\pm})t^2} \quad (4.22)$$

This gives

$$I = 2D_1 \sum_{\pm} \frac{A \mp iBc_{\pm}}{s_{\pm}} \times \arctan\left(\frac{1 + c_{\pm}}{1 - c_{\pm}}\right) \quad (4.23)$$

s_{\pm} is defined as $\sqrt{1 - c_{\pm}^2}$ and one must be careful with the signs. This gives for CUT_1 the approximate expression

$$\begin{aligned} CUT_1 &= D_1 \sum_{\pm} \frac{\sin(\theta)\cos(\phi) \mp iKc_{\pm}}{s_{\pm}} \times \arctan\left(\frac{1 + c_{\pm}}{1 - c_{\pm}}\right) , \\ c_{\pm} &= \frac{-iK\cos(\phi) \pm \sin(\phi)\sqrt{\sin^2(\theta) + K^2}}{\sin(\theta)} \quad (4.24) \end{aligned}$$

Arcus tangent function must be defined in terms of logarithm functions since the argument is complex.

4. In the intermediate region, where the exponential differs from unity one can use expansion in Taylor polynomial to sum over integrals of rational functions of $\cos(\psi)$ and one obtains the expression

$$\begin{aligned} CUT_1 &= D_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \left(\frac{b}{b_0}\right)^n I_n , \\ I_n &= \sum_{\pm} (A \mp iBc_{\pm} I_n(c_{\pm})) , \\ I_n(c) &= \int_0^{\pi/2} \frac{\cos^n(\psi)}{\cos(\psi) - c} \quad (4.25) \end{aligned}$$

$I_n(c)$ can be calculated explicitly by expanding in the integrand $\cos(\psi)^n$ to polynomial with respect to $\cos(\psi) - c$, $c \equiv c_{\pm}$

$$\frac{\cos^n(\psi)}{\cos(\psi) - c} = \sum_{m=0}^{n-1} \binom{n}{m} c^m (\cos(\psi) - c)^{n-m-1} + \frac{c^n}{\cos(\psi) - c} . \quad (4.26)$$

After the change of the integration variable the integral reads as

$$\begin{aligned} I_n(c) &= \sum_{m=0}^{n-1} \sum_{k=0}^{n-m-1} \binom{n}{m} \binom{n-m-1}{k} (-1)^k (1-c)^{n-m-1-k} (1+c)^k c^m I(k, n-m) \\ &+ \frac{c^n}{1-c} \times \log \left[\frac{\sqrt{1-c} + \sqrt{1+c}}{\sqrt{1-c} - \sqrt{1+c}} \right] , \\ I(k, n) &= 2 \int dt \frac{t^{2k}}{(1+t^2)^n} . \end{aligned} \quad (4.27)$$

Partial integration for $I(k, n)$ gives the recursion formula

$$I(k, n) = -\frac{2^{-n+1}}{n-1} + \frac{2k-1}{2(n-1)} \times I(k-1, n-1) . \quad (4.28)$$

The lowest term in the recursion formula corresponds to $I(0, n-k)$, can be calculated by using the expression

$$\begin{aligned} (1+t^2)^{-n} &= \sum_{k=0}^n c(n, k) [(1+it)^{-k} + (1-it)^{-k}] , \\ c(n, k) &= \sum_{l=0}^{n-k-1} c(n-1, k+l) 2^{-l-2} + c(n-1, n-1) 2^{-n+k-1} . \end{aligned} \quad (4.29)$$

The formula is deducible by assuming the expression to be known for n and multiplying the expression with $(1+t^2)^{-1} = [(1+it)^{-1} + (1-it)^{-1}]/2$ and applying this identity to the resulting products of $(1+it)^{-1}$ and $(1-it)^{-1}$. This gives

$$I(0, n) = -2i \sum_{k=2, n} \frac{c(n, k)}{(k-1)} [1 + 2^{(k-1)/2} \sin((k-1)\pi/4)] + c(n, 1) \log \left(\frac{1+i}{1-i} \right) . \quad (4.30)$$

This boils down to the following expression for CUT_1

$$\begin{aligned}
CUT_1 &= D_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \left(\frac{b}{b_0}\right)^n I_n , \\
I_n &= \sum_{\pm} [A \mp iBc_{\pm}] I_n(\cos(c_{\pm})) , \\
I_n(c) &= \sum_{m=1}^{n-1} \sum_{k=0}^{n-m-1} \binom{n}{m} \binom{n-m-1}{k} (1-c)^{n-m-1-k} (1+c)^k c^m I(k, n-m-1) \\
&\quad + \frac{c^n}{1-c} \times \log \left[\frac{\sqrt{1-c} + \sqrt{1+c}}{\sqrt{1-c} - \sqrt{1+c}} \right] , \\
I(k, n) &= -\frac{2^{-n+1}}{n-1} + \frac{2k-1}{2(n-1)} \times I(k-1, n-1) , \\
I(0, n) &= -2i \sum_{k=2}^n \frac{c(n, k)}{(k-1)} [1 + 2^{(k-1)/2} \sin((k-1)\pi/4)] - c(n, 1) , \\
c(n, k) &= \sum_{l=0}^{n-k-1} c(n-1, k+l) 2^{-l-2} + c(n-1, n-1) 2^{-n+k-1} . \tag{4.31}
\end{aligned}$$

This expansion in powers of c_{\pm} fails to converge when their values are very large. This happens in the case of τ -pion production amplitude. In this case one typically has however the situation in which the conditions $A_1 \simeq iB\cos(\psi)/D$ holds true in excellent approximation and one can write

$$\begin{aligned}
CUT_1 &\simeq i \frac{D_1 B}{D} \times \sum_{n=0,1,\dots} \frac{(-1)^n}{n! 2^n} \left(\frac{b}{b_0}\right)^n I_n \times , \\
I_n &= \int_0^{\pi/2} \cos(\psi)^{n+1} d\psi = \sum_{k=0}^{n+1} \binom{n+1}{k} \frac{i^{n-2k} - 1}{n+1-2k} . \tag{4.32}
\end{aligned}$$

The denominator X_1 vanishes, when the conditions

$$\begin{aligned}
\cos(\theta) &= \frac{\beta}{v_{cm}} , \\
\sin(\phi) &= \cos(\psi) \tag{4.33}
\end{aligned}$$

hold. In forward direction the conditions express the vanishing of the z-component of the lepton velocity in velocity cm frame as one can realize by noticing that condition reduces to the condition $v = \beta/2$ in non-relativistic limit. This corresponds to the production of lepto-pion with momentum in scattering plane and with direction angle $\cos(\theta) = \beta/v_{cm}$.

CUT_1 diverges logarithmically for these values of kinematical variables at the limit $\phi \rightarrow 0$ as is easy to see by studying the behavior of the integral near as K approaches zero so that X_1 approaches zero at $\sin(\phi) = \cos(\Phi)$ and the integral over a small interval of length $\Delta\Psi$ around $\cos(\Psi) = \sin(\phi)$ gives a contribution proportional to $\log(A + B\Delta\Psi)/B$, $A = K[K - 2i\sin(\theta)\sin^2(\phi)]$ and $B = 2\sin(\theta)\cos(\phi)[\sin(\theta)\sin(\phi) - iK\cos(\phi)]$. Both A and B vanish at the limit $\phi \rightarrow 0$, $K \rightarrow 0$. The exponential damping reduces the magnitude of the singular contribution for large values of $\sin(\phi)$ as is clear from the first formula.

2. The contribution of the second cut

The expression for CUT_2 reads as

$$\begin{aligned}
CUT_2 &= D_2 \exp\left(-\frac{b}{b_2}\right) \times \int_0^{\pi/2} \exp\left(i\frac{b}{b_1} \cos(\psi)\right) A_2 d\psi , \\
D_2 &= -\frac{\sin(\frac{\phi}{2})}{u \sin(\theta)} , \\
b_1 &= \frac{\hbar \beta}{m \gamma_1} , \quad b_2 = \frac{\hbar}{mb \gamma_1} \frac{1}{\sin(\theta) \cos(\phi)} \\
A_2 &= \frac{A \cos(\psi) + B}{\cos^2(\psi) + 2iC \cos(\psi) + D} , \\
A &= \sin(\theta) \cos(\phi) u , \quad B = \frac{w}{v_{cm}} + \frac{v}{\beta} \sin^2(\theta) [\sin^2(\phi) - \cos^2(\phi)] , \\
C &= \frac{\beta w \cos(\phi)}{w v_{cm} \sin(\theta)} , \quad D = -\frac{1}{u^2} \left(\frac{\sin^2(\phi)}{\gamma^2} + \beta^2 (v^2 \sin^2(\theta) - \frac{2vw}{v_{cm}}) \cos^2(\phi) \right) \\
&\quad + \frac{w^2}{v_{cm}^2 u^2 \sin^2(\theta)} + 2i \frac{\beta v}{u} \sin(\theta) \cos(\phi) , \\
u &= 1 - \beta v \cos(\theta) , \quad w = 1 - \frac{v_{cm}}{\beta} \cos(\theta) .
\end{aligned} \tag{4.34}$$

$$(4.35)$$

The denominator X_2 has no poles and the contribution of the second cut is therefore always finite.

1. The factor $\exp(-b/b_2)$ gives an exponential reduction and the contribution of CUT_2 is large only when the criterion

$$b < \frac{\hbar}{m} \times \frac{1}{v \gamma_1 \sin(\theta) \cos(\phi)}$$

for the impact parameter b is satisfied. Large values of \hbar increase the range of allowed impact parameters since the Compton length of lepto-pion increases.

2. At the limit when the exponent becomes very large the variation of the phase factor implies destructive interference and one can perform stationary phase approximation around $\psi = \pi/2$. This gives

$$\begin{aligned}
CUT_2 &\simeq \sqrt{\frac{2\pi b_1}{b}} \times D_2 \times \exp\left(\frac{b}{b_2}\right) A_2(\psi = 0) , \\
D_2 &= -\frac{\sin(\frac{\phi}{2})}{u \sin(\theta)} , \quad A_2 = \frac{A}{D} .
\end{aligned} \tag{4.36}$$

3. As for CUT_1 , the integral over ψ can be expressed as a finite sum of integrals of rational functions, when the value of $(b/b_1) \cos(\psi)$ is so small that $\exp(i(b/b_1) \cos(\psi))$ can be approximated by a Taylor polynomial. More generally, one obtains the expansion

$$\begin{aligned}
CUT_2 &= D_2 \exp\left(-\frac{b}{b_2}\right) \times \sum_{n=0}^{\infty} \frac{1}{n!} i^n \left(\frac{b}{b_1}\right)^n I_n(A, B, C, D) , \\
I_n(A, B, C, D) &= \int_0^{\pi/2} \cos(\psi)^n \frac{A + iB \cos(\psi)}{\cos^2(\psi) + C \cos(\psi) + D} .
\end{aligned} \tag{4.37}$$

The integrand of $I_n(A, B, C, D)$ is same rational function as in the case of CUT_1 but the parameters A, B, C, D given in the expression for CUT_2 are different functions of the kinematical variables. The functions appearing in the expression for integrals $I_n(c)$ correspond to the roots of the denominator of A_2 and are given by $c_{\pm} = -iC \pm \sqrt{-C^2 - D}$, where C and D are the function appearing in the general expression for CUT_2 in Eq. 4.35.




Figure 3: Evaluation of k_y -integral using residue calculus.




Figure 4: Evaluation of k_x -integral using residue calculus.

4.2 Production Amplitude In Quantum Model

The previous expressions for CUT_1 and CUT_2 as such give the production amplitude for given b in the classical model and the cross section can be calculated by integrating over the values of b . The finite Taylor expansion of the amplitude in powers of b allows explicit formulas when impact parameter cutoff is assumed.

4.2.1 General expression of the production amplitude

In quantum model the production amplitude can be reduced to simpler form by using the defining integral representation of Bessel functions

$$\begin{aligned}
 f_B &= i \int F(b) J_0(\Delta kb) (CUT_1 + CUT_2) b db , \\
 F &= 1 \text{ for } \exp(i(S)) \text{ option} , \\
 F(b \geq b_{cr}) &= \int dz \frac{1}{\sqrt{z^2 + b^2}} = 2 \ln\left(\frac{\sqrt{a^2 - b^2} + a}{b}\right) \text{ for } \exp(i(S)) - 1 \text{ option} , \\
 \Delta k &= 2k \sin\left(\frac{\alpha}{2}\right) , \quad k = M_R \beta .
 \end{aligned} \tag{4.38}$$

Note that F is a rather slowly varying function of b and in good approximation can be replaced by its average value $A(b, p)$, which has been already explicitly calculated as power series in b . α_{em} corresponds to the value of α_{em} for the standard value of Planck constant.

4.2.2 The limit $\Delta k = 0$

The integral of the contribution of CUT_1 over the impact parameter b involves integrals of the form

$$\begin{aligned}
 J_{1,n} &= b_0^2 \int J_0(\Delta kb) F(b) x^{n+1} dx , \\
 x &= \frac{b}{b_0} .
 \end{aligned} \tag{4.39}$$

Here a is the upper impact parameter cutoff. For CUT_2 one has integrals of the form

$$\begin{aligned}
 J_{2,n} &= b_1^2 \left(\frac{b_2}{b_1}\right)^{n+2} \int J_0(\Delta kb) F(b) \exp(-x) x^{n+1} dx , \\
 x &= \frac{b}{b_2} .
 \end{aligned} \tag{4.40}$$

Using the following approximations it is possible to estimate the integrals analytically.

1. The logarithmic term is slowly varying function and can be replaced with its average value

$$F(b) \rightarrow \langle F(b) \rangle \equiv F . \tag{4.41}$$

2. Δk is fixed once the value of the impact parameter is known. At the limit $\Delta k = 0$ making sense for very high energy collisions one can put the value of Bessel function to $J_0(0) = 1$. Hence it is advantageous to calculate the integrals of $\int CUT_i b db$.

Consider first the integral $\int CUT_1 b db$. If exponential series converges rapidly one can use Taylor polynomial and calculate the integrals explicitly. When this is not the case one can calculate integral approximately and the total integral is sum over two contributions:

$$\int CUT_1 b db = I_a + I_b . \tag{4.42}$$

1. The region in which Taylor expansion converges rapidly gives rise integrals

$$\begin{aligned} I_{1,n} &\simeq b_0^2 \int x^{n+1} dx = b_0^2 \frac{1}{n+2} \left[\left(\frac{b_{max}}{b_0} \right)^{n+2} - \left(\frac{b_{cr}}{b_0} \right)^{n+2} \right] \simeq b_0^2 \frac{1}{n+2} \left(\frac{b_{max}}{b_0} \right)^{n+2} , \\ I_{2,n} &\simeq b_1^2 \left(\frac{b_2}{b_1} \right)^{n+2} \int exp(-x) x^{n+1} dx = b_1^2 \left(\frac{b_2}{b_1} \right)^{n+2} (n+1)! . \end{aligned} \quad (4.43)$$

2. For the perturbative part of CUT_1 one obtains the expression

$$\begin{aligned} I_a &= \int_0^{b_{max}} CUT_1 b db = D_1 \times b_0^2 \times \sum_{n=0}^{\infty} \frac{1}{n!(n+2)} \left(\frac{b_{max}}{b_0} \right)^{n+2} I_n(A, B, C, D) , \\ D_1 &= -\frac{1}{2} \frac{\sin(\phi)}{\sin(\theta)} , \quad b_0 = \frac{\hbar\beta\gamma}{m\gamma_1} . \end{aligned} \quad (4.44)$$

There b_{max} is the largest value of b for which the series converges sufficiently rapidly.

3. The convergence of the exponential series is poor for large values of b/b_0 , that is for $b > b_m$. In this case one can use the approximation in which the multiplier of exponent function in the integrand is replaced with its value at $\psi = \pi/2$ so that amplitude becomes proportional to b_0/b . In this case the integral over b gives a factor proportional to ab_0 , where a is the impact parameter cutoff.

$$\begin{aligned} I_b &\equiv \int_{b_m}^a CUT_1 b db \simeq b_0(a - b_m) D_1 \times A_1(\psi = \pi/2) \\ &= \frac{\beta\gamma}{\gamma_1} \times \frac{\hbar}{m} \times \frac{\sin^2(\theta) \cos(\phi) \sin(\phi)}{\sin^2(\theta) \sin^2(\phi) + K^2} , \\ D_1 &= -\frac{1}{2} \frac{\sin(\phi)}{\sin(\theta)} , \quad A_1(\psi = \pi/2) = \frac{A}{D} . \end{aligned} \quad (4.45)$$

4. As already explained, the expansion based on partial fractions does not converge, when the roots c_{\pm} have very large values. This indeed occurs in the case of τ -pion production cross section. In this case one has $A_1 \simeq iB \cos(\psi)/D$ in excellent approximation and one can calculate CUT_1 in much easier manner. Using the formula of Eq. 4.32 for CUT_1 , one obtains

$$\begin{aligned} \int CUT_1 b db &\simeq b_0^2 \frac{D_1 B}{D} \times \sum_{n=0,1,\dots} \frac{(-1)^n}{n!(n+2)2^n} \times \sum_{k=0}^{n+1} \binom{n+1}{k} c_{n,k} \times \left(\frac{b_{max}}{b_0} \right)^n , \\ c_{n,k} &= \frac{i^{n+1-2k} - 1}{n+1-2k} \text{ for } n \neq 2k-1 , \quad c_{n,k} = \frac{i\pi}{2} \text{ for } n = 2k-1 , \end{aligned} \quad (4.46)$$

Note that for $n = 2k + 1 = k$ the coefficient diverges formally and actual

Highly analogous treatment applies to the integral of CUT_2 .

1. For the perturbative contribution to $\int CUT_2 b db$ one obtains

$$\begin{aligned}
I_a &= \int_0^{b_{1,max}} CUT_2 b db = b_1^2 D_2 \sum_{n=0}^{\infty} (n+1) i^n I_n(A, B, C, D) \times \left(\frac{b_2}{b_1}\right)^{n+2} , \\
D_2 &= -\frac{\sin(\frac{\phi}{2})}{u \sin(\theta)} , \\
b_1 &= \frac{\hbar \beta}{m \gamma_1} , \quad b_2 = \frac{\hbar}{m \gamma_1} \frac{1}{\sin(\theta) \cos(\phi)} .
\end{aligned} \tag{4.47}$$

2. Taylor series converges slowly for

$$\frac{b_1}{b_2} = \frac{\sin(\theta) \cos(\phi)}{\beta} \rightarrow 0 .$$

In this case one can replace $\exp(-b/b_2)$ with unity or expand it as Taylor series taking only few terms. This gives the expression for the integral which is of the same general form as in the case of CUT_1

$$I_a = \int_0^{b_{max}} CUT_2 b db = b_1^2 D_2 \sum_{n=0}^{\infty} \frac{i^n}{n!(n+2)} I_n(A, B, C, D) \left(\frac{b_{max}}{b_1}\right)^{n+1} . \tag{4.48}$$

3. Also when b/b_1 becomes very large, one must apply stationary phase approximation to calculate the contribution of CUT_2 which gives a result proportional to $\sqrt{b_1/b}$. Assume that $b_m \gg b_1$ is the value of impact parameter above which stationary phase approximation is good. This gives for the non-perturbative contribution to the production amplitude the expression

$$\begin{aligned}
I_b &= \int_{b_m}^a CUT_2 b db = k \sqrt{\frac{2\pi b_1}{b_2}} b_2^2 \times D_2 \times A_2(\psi = \pi/2) , \\
k &= \int_{x_1}^{x_2} \exp(-x) x^{1/2} dx = 2 \int_{\sqrt{x_1}}^{\sqrt{x_2}} \exp(-u^2) u^2 du , \\
x_1 &= \frac{b_m}{b_2} , \quad x_2 = \frac{a}{b_2} .
\end{aligned} \tag{4.49}$$

In good approximation one can take $x_2 = \infty$. $x_1 = 0$ gives the upper bound $k \leq \sqrt{\pi}$ for the integral.

Some remarks relating to the numerics are in order.

1. The contributions of both CUT_1 and CUT_2 are proportional to $1/\sin(\theta)$ in the forward direction. The denominators of A_i however behave like $1/\sin^2(\theta)$ at this limit so that the amplitude behaves as $\sin(\theta)$ at this limit and the amplitude approaches to zero like $\sin(\theta)$. Therefore the singularity is only apparent but must be taken into account in the calculation since one has $c_{\pm} \rightarrow i\infty$ at this limit for CUT_2 and for CUT_1 the roots approach to $c_+ = c_- = i\infty$. One must pose a cutoff θ_{min} below which the contribution of CUT_1 and CUT_2 are calculated directly using approximate the expressions for $D_i A_i$.

$$\begin{aligned}
D_1 A_1 &\rightarrow -\frac{i}{K} \cos(\psi) \times \sin(\theta) \rightarrow 0 \\
D_2 A_2 &\rightarrow -\frac{uw_{cm}}{w} \times \sin(\theta) \rightarrow 0 .
\end{aligned} \tag{4.50}$$

In good approximation both contributions vanish since also $\sin^2(\theta)$ factor from the phase space integration reduces the contribution.

2. A second numerical problem is posed by the possible vanishing of

$$K = \beta\gamma\left(1 - \frac{v_c m}{\beta} \cos(\theta)\right) .$$

In this case the roots $c_{\pm} = \pm \sin(\phi)$ are real and c_+ gives rise to a pole in the integrand.

The singularity to the amplitude comes from the logarithmic contributions in the Taylor series expansion of the amplitude. The sum of the singular contributions coming from c_+ and c_- are of form

$$\frac{c_n}{2} (\sqrt{1 - \sin(\phi)} + \sqrt{1 + \sin(\phi)}) \log\left(\frac{1+u}{1-u}\right) , \quad u = \sqrt{\frac{1 + \sin(\phi)}{1 - \sin(\phi)}} .$$

Here c_n characterizes the $1/(\cos(\psi) - c_{\pm})$ term of associated with the $\cos(\psi)^n$ term in the Taylor expansion. Logarithm becomes singular for the two terms in the sum at the limit $\phi \rightarrow 0$. The sum however behaves as

$$\frac{c_n}{2} \sin(\phi) \log\left(\frac{\sin(\phi)}{2}\right) .$$

so that the net result vanishes at the limit $\phi \rightarrow 0$. It is essential that the logarithmic singularities corresponding to the roots c_+ and c_- cancel each other and this must be taken into account in numerics. There is also apparent singularity at $\phi = \pi/2$ canceled by $\cos(\phi)$ factor in D_1 . The simplest manner to get rid of the problem is to exclude small intervals $[0, \epsilon]$ and $[\pi/2 - \epsilon, \pi/2]$ from the phase space volume.

4.2.3 Improved approximation to the production cross section

The approximation $J_0(\Delta k_T(b)b) = 1$ and $F(b) = F = \text{constant}$ allows to perform the integrations over impact parameter explicitly (for $\exp(iS)$ option $F = 1$ holds true identically in the lowest order approximation). An improved approximation is obtained by dividing the range of impact parameters to pieces and performing the integrals over the impact parameter ranges exactly using the average values of these functions. This requires only a straightforward generalization of the formulas derived above involving integrals of the functions x^n and $\exp(-x)x^n$ over finite range. Obviously this is still numerically well-controlled procedure.

4.3 Evaluation Of The Singular Parts Of The Amplitudes

The singular parts of the amplitudes $CUT_{1,sing}$ and $B_{1,sing}$ are rational functions of $\cos(\psi)$ and the integrals over ψ can be evaluated exactly.

In the classical model the expression for $U_{1,sing}$ appearing as integrand in the expression of $CUT_{1,sing}$ reads as

$$\begin{aligned} A_{1,sing} &= -\frac{1}{2\sqrt{K^2 + \sin^2(\theta)}} (\sin(\theta)\cos(\phi)A_a + iKA_b) , \\ A_a &= I_1(\beta, \pi/2) = \int_0^{\pi/2} d\psi f_1 , \\ A_b &= I_2(\beta, \pi/2) = \int_0^{\pi/2} d\psi f_2 , \\ f_1 &= \frac{1}{(\cos(\psi) - c_1)(\cos(\psi) - c_2)} , \\ f_2 &= \cos(\psi)f_1 , \\ c_1 &= \frac{-iK\cos(\phi) + \sin(\phi)\sqrt{K^2 + \sin^2(\theta)}}{\sin(\theta)} , \\ c_2 &= -\bar{c}_1 . \end{aligned} \tag{4.51}$$

Here c_i are the roots of the polynomial X_1 appearing in the denominator of the integrand.

In quantum model the approximate expression for the singular contribution to the production amplitude can be written as

$$\begin{aligned}
B_{1,sing} &\simeq k_1 \frac{\sin(\theta)\sin(\phi)}{2\sqrt{K^2 + \sin^2(\theta)}} \sum_n \langle F \rangle_n (I(x(n+1)) - I(x(n))) , \\
I(x) &= \exp\left(-\frac{\sin(\phi)x}{\sin(\phi_0)}\right) (\sin(\theta)\cos(\phi)A_a(\Delta ka, x) + iKA_b(\Delta ka, x)) , \\
k_1 &= 2\pi^2 M_R Z_1 Z_2 \alpha_{em} \frac{\sqrt{2}}{\sqrt{\Delta k \pi}} \sin(\phi_0) .
\end{aligned} \tag{4.52}$$

The expressions for the amplitudes $A_a(k, x)$ and $A_b(k, x)$ read as

$$\begin{aligned}
A_a(k, x) &= \cos(kx)I_3(k, 0, \pi/2) + i\sin(\phi_0)k\sin(kx)I_5(k, 0, \pi/2) , \\
A_b(k, x) &= \cos(kx)I_4(k, 0, \pi/2) + i\sin(\phi_0)k\sin(kx)I_3(k, 0, \pi/2) , \\
I_i(k, \alpha, \beta) &= \int_\alpha^\beta f_i(k)d\psi , \\
f_3(k) &= \frac{\cos(\psi)}{(\cos^2(\psi) + \sin^2(\phi_0)k^2)} f_1(k) , \\
f_4(k) &= \cos(\psi)f_3(k) , \\
f_5(k) &= \frac{1}{(\cos^2(\psi) + \sin^2(\phi_0)k^2)} f_1(k) .
\end{aligned} \tag{4.53}$$

The expressions for the integrals I_i as functions of the endpoints α and β can be written as

$$\begin{aligned}
I_1(k, \alpha, \beta) &= I_0(c_1, \alpha, \beta) - I_0(c_2, \alpha, \beta) , \\
I_2(\alpha, \beta) &= c_1 I_0(c_1, \alpha, \beta) - c_2 I_0(c_2, \alpha, \beta) , \\
I_3 &= C_{34} \sum_{i=1,2,j=3,4} \frac{1}{(c_i - c_j)} (c_i I_0(c_i, \alpha, \beta) - c_j I_0(c_j, \alpha, \beta)) , \\
I_4 &= C_{34} \sum_{i=1,2,j=3,4} \frac{1}{(c_i - c_j)} ((c_i - c_j)(\beta - \alpha) - c_i^2 I_0(c_i, \alpha, \beta) + c_j^2 I_0(c_j, \alpha, \beta)) , \\
I_5 &= C_{34} \sum_{i=1,2,j=3,4} \frac{1}{(c_i - c_j)} (I_0(c_i, \alpha, \beta) - I_0(c_j, \alpha, \beta)) , \\
C_{34} &= \frac{1}{c_3 - c_4} = \frac{1}{2ikas(\phi_0)} .
\end{aligned} \tag{4.54}$$

The parameters c_1 and c_2 are the zeros of X_1 as function of $\cos(\psi)$ and c_3 and c_4 the zeros of the function $\cos^2(\psi) + k^2 a^2 \sin^2(\phi_0)$:

$$\begin{aligned}
c_1 &= \frac{-iK\cos(\phi) + \sin(\phi)\sqrt{K^2 + \sin^2(\theta)}}{\sin(\theta)} , \\
c_2 &= \frac{-iK\cos(\phi) - \sin(\phi)\sqrt{K^2 + \sin^2(\theta)}}{\sin(\theta)} , \\
c_3 &= ikas(\phi_0) , \\
c_4 &= -ikas(\phi_0) .
\end{aligned} \tag{4.55}$$

The basic integral $I_0(c, \alpha, \beta)$ appearing in the formulas is given by

$$\begin{aligned}
 I_0(c, \alpha, \beta) &= \int_{\alpha}^{\beta} d\psi \frac{1}{(\cos(\psi) - c)} , \\
 &= \frac{1}{\sqrt{1-c^2}} (f(\alpha) - f(\beta)) , \\
 f(x) &= \ln\left(\frac{(1 + \tan(x/2)t_0)}{(1 - \tan(x/2)t_0)}\right) , \\
 t_0 &= \sqrt{\frac{1-c}{1+c}} .
 \end{aligned} \tag{4.56}$$

From the expression of I_0 one discovers that scattering amplitude has logarithmic singularity, when the condition $\tan(\alpha/2) = 1/t_0$ or $\tan(\beta/2) = 1/t_0$ is satisfied and appears, when c_1 and c_2 are real. This happens at the cone $K = 0$ ($\theta = \theta_0$), when the condition

$$\begin{aligned}
 \sqrt{\frac{(1 - \sin(\phi))}{(1 + \sin(\phi))}} &= \tan(x/2) , \\
 x &= \alpha \text{ or } \beta .
 \end{aligned} \tag{4.57}$$

holds true. The condition is satisfied for $\phi \simeq x/2$. $x = 0$ is the only interesting case and gives singularity at $\phi = 0$. In the classical case this gives logarithmic singularity in production amplitude for all scattering angles.

REFERENCES

Theoretical Physics

- [B1] Zuber J-B Itzykson C. *Quantum Field Theory*. Mc Graw-Hill, New York, 1980.
- [B2] Shajesh KV. Eikonal approximation. Available at: <http://www.nhn.ou.edu/~shajesh/eikonal/sp.pdf>, 2008.
- [B3] Okun LB. *Leptons and Quarks*. North-Holland, Amsterdam, 1982.

Particle and Nuclear Physics

- [C1] Pamela is coming out. Available at: <http://resonaances.blogspot.com/2008/10/pamelas-coming-out.html>.
- [C2] Particle Data Tables. Available at: http://pdg.lbl.gov/2010/listings/contents_listings.html.
- [C3] Who ordered that?! An X-traordinary particle? Available at: <http://www.quantumdiaries.org/2011/10/10/who-ordered-that-an-x-traordinary-particle/>.
- [C4] Chodos A. 17(4):211–223, 1987.
- [C5] Strumia A. Dark Matter Interpretations of the Cosmic-Ray e Excesses. *Prog Theor Phys Supplement*. Available at: <http://ptp.ipap.jp/link?PTPS/180/128/>, (180):128–134, 2009.
- [C6] Weiner N Arkani-Hamed N. LHC Signals for a SuperUnified Theory of Dark Matter. Available at: <http://arxiv.org/abs/0810.0714>, 2008.
- [C7] Sarkar S Barger V, Phillips RJ. *Phys Lett B*, 352:365–371, 1995.

- [C8] Cofield C. Antimatter from lightning flashes the Fermi space telescope. *Symmetry Breaking (joint Fermilab/SLAC publication)*. Available at: <http://tinyurl.com/p2z3n9p>, (5), October 2009.
- [C9] Weniger C. A Tentative Gamma -Ray Line from Dark Matter Annihilation at the Fermi Large Area Telescope. Available at: <http://arxiv.org/abs/1204.2797>, 2012.
- [C10] CDF Collaboration. Study of multi-muon events produced in p-pbar collisions at $\sqrt{s}=1.96$ TeV. Available at: http://arxiv.org/PS_cache/arxiv/pdf/0810/0810.0714v1.pdf, 2008.
- [C11] CDMS collaboration. Results from the Final Exposure of the CDMS II Experiment. Available at: <http://arxiv.org/abs/0912.3592>, 2009.
- [C12] CoGENT collaboration. Results from a Search for Light-Mass Dark Matter with a P-type Point Contact Germanium Detector. Available at: <http://arxiv.org/abs/1002.4703>, 2010.
- [C13] DAMA Collaboration. New results from DAMA/LIBRA. Available at: <http://arxiv.org/abs/1002.1028>, 2010.
- [C14] DAMA collaboration. Results from DAMA/LIBRA at Gran Sasso. *Found Phys* ..Available at: <http://people.roma2.infn.it/~dama/web/publ10.html>, 40:900, 2010.
- [C15] PAMELA Collaboration. Observation of an anomalous positron abundance in the cosmic radiation. Available at: http://arxiv.org/PS_cache/arxiv/pdf/0810/0810.4995v1.pdf, 2008.
- [C16] Xenon100 collaboration. Dark Matter Results from 100 Live Days of XENON100 Data. Available at: <http://arxiv.org/abs/1104.2549>, 2011.
- [C17] Toldra R Escribano R, Masso E. *Phys Lett B*, 356, 1995.
- [C18] Abeysekera AU et al. Extended gamma-ray sources around pulsars constrain the origin of the positron flux at Earth. *Science*. <http://tinyurl.com/yach5g83>, 358(6365):911–914, 2017.
- [C19] Akesson T et al. *Phys Lett B*, 463:36, 1987.
- [C20] Armbruster Collaboration: KARMEN B et al. *Phys Lett B*, 348, 1995.
- [C21] Chliapnikov PV et al. *Phys Lett B*, 141, 1984.
- [C22] Churazov E et al. Positron annihilation spectrum from the Galactic Center region observed by SPI/INTEGRAL. *Mon Not Roy Astron Soc*, 357, 2005.
- [C23] Clemente M et al. *Phys Rev*, 137, 1984.
- [C24] Cowan T et al. *Phys Rev*, 54:56, 1985.
- [C25] Daniels T et al. $j/\psi, \psi' \rightarrow \mu^+\mu^-$ and $b \rightarrow j/\psi, \psi'$ cross-sections. In *DPP Conf.*, pages 552–555, 1994.
- [C26] Dantzman K et al. *Phys Rev*, 62, 1989.
- [C27] Giromini P et al. Phenomenological interpretation of the multi-muon events reported by the CDF collaboration. Available at: <http://arxiv.org/abs/0810.5730>, 2008.
- [C28] Goshaw AT et al. *Phys Rev*, 43, 1979.
- [C29] Judge S et al. *Phys Rev*, 65(8), 1990.
- [C30] Koenig W et al. *Z Physik A*, 3288, 1987.
- [C31] Salapura P et al. *Phys Lett B*, 245:153, 1990.

- [C32] Schweppe J et al. *Phys Rev*, 51, 1983.
- [C33] Tsertos H et al. *Phys Lett*, 273:326, 1985.
- [C34] Weidenspointner G et al. The sky distribution of positronium annihilation continuum emission measured with SPI/INTEGRAL. *Astron Astrophys* . Available at: <http://arxiv.org/abs/astro-ph/0601673>, 450, 2006.
- [C35] Westbrook CI et al. *Phys Rev*, 58:1328, 1987.
- [C36] Yue AT et al. Improved determination of the neutron lifetime. *PRL*. Available at:<http://tinyurl.com/h88n57j>, 111(222501), 2013.
- [C37] Chang J et al (ATIC). In *29th International Cosmic Ray Conferences (ICRC 2005), Pune, India, 31 August 03 - 10 2005*, 2005.
- [C38] Valencia G He XG, Tandean J. Has HyperCP Observed a Light Higgs Boson? *Phys Rev D*. Available at: <http://arxiv.org/abs/hep-ph/0610274>, 74, 2007.
- [C39] Valencia G He XG, Tandean J. Light Higgs Production in Hyperon Decay. *Phys Rev* . Available at: <http://arxiv.org/abs/hep-ph/0610362>, 98, 2007.
- [C40] Goodenough L Hooper D. Dark Matter Annihilation in The Galactic Center As Seen by the Fermi Gamma Ray Space Telescope. Available at: <http://arxiv.org/pdf/1010.2752v1>, 2010.
- [C41] Zeller M Kraus L. *Phys Rev D*, 34, 1986.
- [C42] McKee M. Earth creates powerful gamma-ray flashes. Available at: <http://www.newscientist.com/channel/space/dn7025>, February 2005.
- [C43] Jones N. Lightning strikes release powerful X-ray bursts. *New Scientist* . Available at: <http://tinyurl.com/zqc7r7z>, 2381, February 2003.
- [C44] Voit P. A new long-lived particle by CDF experiment. Available at: <http://www.math.columbia.edu/~voit/wordpress/?p=1045>, 2008.
- [C45] Courtland R. Hints of lightweight dark matter particle found in space. Available at: <http://tinyurl.com/3y3mj9c>, 2010.
- [C46] Foot R. A CoGeNT confirmation of the DAMA signal. Available at: <http://arxiv.org/abs/1004.1424>, 2010.
- [C47] Barshay S. *Mod. Phys Lett A*, 7(20):1843, 1992.
- [C48] Dorigo T. Saving a good text from a few mistakes. Available at: <http://dorigo.wordpress.com/2008/11/21/saving-a-good-text-from-a-few-mistakes/>, 2008.
- [C49] Dorigo T. Some notes on the multi-muon analysis - part I. Available at: <http://dorigo.wordpress.com/2008/11/08/some-notes-on-the-multi-muon-analysis-part-i/>, 2008.
- [C50] Tomasi-Gustafsson E Tatischeff B. Search for low-mass exotic mesonic structures: II. Attempts to understand the experimental results. *Part Nucl Lett*, 5(5):709–713, 2008.

Cosmology and Astro-Physics

- [E1] Fermi 130 GeV gamma-ray excess and dark matter annihilation in sub-haloes and in the Galactic centre. Available at: <http://arxiv.org/pdf/1205.1045.pdf>, 2012.
- [E2] Nottale L Da Rocha D. Gravitational Structure Formation in Scale Relativity. Available at: <http://arxiv.org/abs/astro-ph/0310036>, 2003.
- [E3] Boehm C et al. MeV Dark Matter: Has It Been Detected? *New Scientist*. Available at: <http://arxiv.org/abs/astro-ph/0309686>, October 2003.

Neuroscience and Consciousness

- [J1] Blackman CF. *Effect of Electrical and Magnetic Fields on the Nervous System*, pages 331–355. Plenum, New York, 1994.
- [J2] Cowan JD. Spontaneous symmetry breaking in large-scale nervous activity. *Int J Quant Chem*, 22:1059–1082, 1982.

Books related to TGD

- [K1] Pitkänen M. The Status of Lepto-pion Hypothesis. In *TGD and p-Adic Numbers*. Online book. Available at: http://tgdtheory.fi/public_html/padtgd/padtgd.html#leptc, 2000.
- [K2] Pitkänen M. About Strange Effects Related to Rotating Magnetic Systems . In *TGD and Fringe Physics*. Online book. Available at: http://tgdtheory.fi/public_html/freenergy/freenergy.html#Faraday, 2006.
- [K3] Pitkänen M. Cosmic Strings. In *Physics in Many-Sheeted Space-Time*. Online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#cstrings, 2006.
- [K4] Pitkänen M. Dark Forces and Living Matter. In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#darkforces, 2006.
- [K5] Pitkänen M. Dark Nuclear Physics and Condensed Matter. In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#exonuclear, 2006.
- [K6] Pitkänen M. Does TGD Predict the Spectrum of Planck Constants? In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#Planck, 2006.
- [K7] Pitkänen M. General Ideas about Many-Sheeted Space-Time: Part I. In *Physics in Many-Sheeted Space-Time*. Online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#topcond, 2006.
- [K8] Pitkänen M. Magnetic Sensory Canvas Hypothesis. In *TGD and EEG*. Online book. Available at: http://tgdtheory.fi/public_html/tgdeeg/tgdeeg.html#mec, 2006.
- [K9] Pitkänen M. Massless states and particle massivation. In *p-Adic Physics*. Online book. Available at: http://tgdtheory.fi/public_html/padphys/padphys.html#mless, 2006.
- [K10] Pitkänen M. New Particle Physics Predicted by TGD: Part I. In *p-Adic Physics*. Online book. Available at: http://tgdtheory.fi/public_html/padphys/padphys.html#mass4, 2006.
- [K11] Pitkänen M. New Particle Physics Predicted by TGD: Part II. In *p-Adic Physics*. Online book. Available at: http://tgdtheory.fi/public_html/padphys/padphys.html#mass5, 2006.
- [K12] Pitkänen M. Nuclear String Hypothesis. In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#nuclstring, 2006.
- [K13] Pitkänen M. p-Adic Particle Massivation: Hadron Masses. In *p-Adic Length Scale Hypothesis and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/padphys/padphys.html#mass3, 2006.
- [K14] Pitkänen M. TGD and Nuclear Physics. In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#padnucl, 2006.

- [K15] Pitkänen M. TGD as a Generalized Number Theory: Infinite Primes. In *TGD as a Generalized Number Theory*. Online book. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#visionc, 2006.
- [K16] Pitkänen M. The Recent Status of Lepto-hadron Hypothesis. In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#leptc, 2006.
- [K17] Pitkänen M. Three new physics realizations of the genetic code and the role of dark matter in bio-systems. In *Genes and Memes*. Online book. Available at: http://tgdtheory.fi/public_html/genememe/genememe.html#dnatqccodes, 2006.
- [K18] Pitkänen M. Summary of TGD Inspired Ideas about Free Energy. In *TGD and Fringe Physics*. Online book. Available at: http://tgdtheory.fi/public_html/freenergy/freenergy.html#freerg, 2013.
- [K19] Pitkänen M. Criticality and dark matter. In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#qcritdark, 2014.
- [K20] Pitkänen M. Quantum gravity, dark matter, and prebiotic evolution. In *Genes and Memes*. Online book. Available at: http://tgdtheory.fi/public_html/genememe/genememe.html#hgrprebio, 2014.
- [K21] Pitkänen M. Unified Number Theoretical Vision. In *TGD as a Generalized Number Theory*. Online book. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#numbervision, 2014.

Articles about TGD

- [L1] Pitkänen M. Further Progress in Nuclear String Hypothesis. Available at: <http://tgdtheory.fi/articles/nuc1string.pdf>, 2007.
- [L2] Pitkänen M. CMAP representations about TGD, and TGD inspired theory of consciousness and quantum biology. Available at: <http://www.tgdtheory.fi/tgdglossary.pdf>, 2014.
- [L3] Pitkänen M. Cold Fusion Again . Available at: http://tgdtheory.fi/public_html/articles/cfagain.pdf, 2015.
- [L4] Pitkänen M. About Physical Representations of Genetic Code in Terms of Dark Nuclear Strings. Available at: http://tgdtheory.fi/public_html/articles/genecodemodels.pdf, 2016.