

Recent View about Kähler Geometry and Spin Structure of "World of Classical Worlds"

M. Pitkänen,

April 1, 2017

Email: matpitka@luukku.com.

http://tgdtheory.com/public_html/.

Recent postal address: Karkinkatu 3 I 3, 00360, Karkkila, Finland.

Contents

1	Introduction	4
2	WCW As A Union Of Homogenous Or Symmetric Spaces	7
2.1	Basic Vision	7
2.2	Equivalence Principle And WCW	9
2.3	Ep At Quantum And Classical Level	9
2.4	Criticism Of The Earlier Construction	11
2.5	Is WCW Homogenous Or Symmetric Space?	12
2.6	Symplectic And Kac-Moody Algebras As Basic Building Bricks	13
3	Updated View About Kähler Geometry Of WCW	13
3.1	Kähler Function, Kähler Action, And Connection With String Models	14
3.2	Realization Of Super-Conformal Symmetries	15
3.3	Interior Dynamics For Fermions, The Role Of Vacuum Extremals, And Dark Matter	16
3.4	Classical Number Fields And Associativity And Commutativity As Fundamental Law Of Physics	18

3.4.1	<i>How to achieve associativity in the fermionic sector?</i>	18
3.4.2	<i>Is super-symmetrized Kähler-Dirac action enough?</i>	20
3.4.3	<i>Are 4-D spinor modes consistent with associativity?</i>	22
4	About some unclear issues of TGD	24
4.1	Adelic vision and symmetries	25
4.2	Quantum-classical correspondence for fermions	25
4.3	Strong form of holography for fermions	26
4.4	The relationship between spinors in space-time interior and at boundaries between Euclidian and Minkoskian regions	26
4.5	About second quantization of the induced spinor fields	27
4.6	Is statistical entanglement “real” entanglement?	28
5	About The Notion Of Four-Momentum In TGD Framework	29
5.1	Scale Dependent Notion Of Four-Momentum In Zero Energy Ontology	30
5.2	Are The Classical And Quantal Four-Momenta Identical?	30
5.3	What Equivalence Principle (EP) Means In Quantum TGD?	31
5.4	TGD-GRT Correspondence And Equivalence Principle	33
5.5	How Translations Are Represented At The Level Of WCW ?	34
5.5.1	Discrete Lorentz transformations and time translations act in the space of CDs: inertial four-momentum	34
5.5.2	The action of translations at space-time sheets	35
5.6	Yangian And Four-Momentum	36
5.6.1	Yangian symmetry	36
5.6.2	How to generalize Yangian symmetry in TGD framework?	37
5.6.3	Could Yangian symmetry provide a new view about conserved quantum numbers?	38
6	Generalization Of Ads/CFT Duality To TGD Framework	39
6.1	Does The Exponent Of Chern-Simons Action Reduce To The Exponent Of The Area Of Minimal Surfaces?	40
6.2	Does Kähler Action Reduce To The Sum Of Areas Of Minimal Surfaces In Effective Metric?	40
6.3	Surface Area As Geometric Representation Of Entanglement Entropy?	43
6.3.1	Generalization of AdS/CFT to TGD context	43
6.3.2	Second interpretation	44
6.3.3	With what kind of systems 3-surfaces can entangle?	45
6.3.4	Minimal surface property is not favored in TGD framework	46
6.4	Related Ideas	47
6.4.1	p-Adic variant of Bekenstein-Hawking law	47
6.4.2	What is the space-time correlate for negentropic entanglement?	48
6.5	The Importance Of Being Light-Like	48
6.5.1	The singular objects associated with the induced metric	48
6.5.2	The singular objects associated with the effective metric	49

7	Could One Define Dynamical Homotopy Groups In WCW?	50
7.1	About Cobordism As A Concept	51
7.2	Prastaro's Generalization Of Cobordism Concept To The Level Of Partial Differential Equations	52
7.3	Why Prastaro's Idea Resonates So Strongly With TGD	53
7.4	What Preferred Extremals Are?	54
7.5	Could Dynamical Homotopy/Homology Groups Characterize WCW Topology?	57
7.6	Appendix: About Field Equations Of TGD In Jet Bundle Formulation	60
7.6.1	Field equations as conservation laws, Frobenius integrability conditions, and a connection with quaternion analyticity . . .	61
7.6.2	Jet bundle formalism	65

Abstract

The construction of Kähler geometry of WCW (“world of classical worlds”) is fundamental to TGD program. I ended up with the idea about physics as WCW geometry around 1985 and made a breakthrough around 1990, when I realized that Kähler function for WCW could correspond to Kähler action for its preferred extremals defining the analogs of Bohr orbits so that classical theory with Bohr rules would become an exact part of quantum theory and path integral would be replaced with genuine integral over WCW. The motivating construction was that for loop spaces leading to a unique Kähler geometry. The geometry for the space of 3-D objects is even more complex than that for loops and the vision still is that the geometry of WCW is unique from the mere existence of Riemann connection.

This chapter represents the updated version of the construction providing a solution to the problems of the previous construction. The basic formulas remain as such but the expressions for WCW super-Hamiltonians defining WCW Hamiltonians (and matrix elements of WCW metric) as their anticommutator are replaced with those following from the dynamics of the Kähler-Dirac action.

1 Introduction

The construction of Kähler geometry of WCW (“world of classical worlds”) is fundamental to TGD program. I ended up with the idea about physics as WCW geometry around 1985 and made a breakthrough around 1990, when I realized that Kähler function for WCW could correspond to Kähler action for its preferred extremals defining the analogs of Bohr orbits so that classical theory with Bohr rules would become an exact part of quantum theory and path integral would be replaced with genuine integral over WCW. The motivating construction was that for loop spaces leading to a unique Kähler geometry [A8]. The geometry for the space of 3-D objects is even more complex than that for loops and the vision still is that the geometry of WCW is unique from the mere existence of Riemann connection.

The basic idea is that WCW is union of symmetric spaces G/H labelled by zero modes which do not contribute to the WCW metric. There have been many open questions and it seems the details of the earlier approach [?] must be modified at the level of detailed identifications and interpretations.

1. A longstanding question has been whether one could assign Equivalence Principle (EP) with the coset representation formed by the super-Virasoro representation assigned to G and H in such a manner that the four-momenta associated with the representations and identified as inertial and gravitational four-momenta would be identical. This does not seem to be the case. The recent view will be that EP reduces to the view that the classical four-momentum associated with Kähler action is equivalent with that assignable to Kähler-Dirac action supersymmetrically related to Kähler action: quantum classical correspondence (QCC) would be in question. Also strong form of general coordinate invariance implying strong form of holography in turn implying

that the super-symplectic representations assignable to space-like and light-like 3-surfaces are equivalent could imply EP with gravitational and inertial four-momenta assigned to these two representations.

At classical level EP follows from the interpretation of GRT space-time as effective space-time obtained by replacing many-sheeted space-time with Minkowski space with effective metric determined as a sum of Minkowski metric and sum over the deviations of the induced metrics of space-time sheets from Minkowski metric. Poincare invariance suggests strongly classical EP for the GRT limit in long length scales at least.

2. The detailed identification of groups G and H and corresponding algebras has been a longstanding problem. Symplectic algebra associated with $\delta M_{\pm}^4 \times CP^2$ (δM_{\pm}^4 is light-cone boundary - or more precisely, with the boundary of causal diamond (CD) defined as Cartesian product of CP_2 with intersection of future and past direct light cones of M^4 has Kac-Moody type structure with light-like radial coordinate replacing complex coordinate z . Virasoro algebra would correspond to radial diffeomorphisms. I have also introduced Kac-Moody algebra assigned to the isometries and localized with respect to internal coordinates of the light-like 3-surfaces at which the signature of the induced metric changes from Minkowskian to Euclidian and which serve as natural correlates for elementary particles (in very general sense!). This kind of localization by force could be however argued to be rather ad hoc as opposed to the inherent localization of the symplectic algebra containing the symplectic algebra of isometries as sub-algebra. It turns out that one obtains direct sum of representations of symplectic algebra and Kac-Moody algebra of isometries naturally as required by the success of p-adic mass calculations.
3. The dynamics of Kähler action is not visible in the earlier construction. The construction also expressed WCW Hamiltonians as 2-D integrals over partonic 2-surfaces. Although strong form of general coordinate invariance (GCI) implies strong form of holography meaning that partonic 2-surfaces and their 4-D tangent space data should code for quantum physics, this kind of outcome seems too strong. The progress in the understanding of the solutions of Kähler-Dirac equation led however to the conclusion that spinor modes other than right-handed neutrino are localized at string world sheets with strings connecting different partonic 2-surfaces. This leads to a modification of earlier construction in which WCW super-Hamiltonians are essentially integrals with integrand identified as a Noether super current for the deformations in G . Each spinor mode gives rise to super current and the modes of right-handed neutrino and other fermions differ in an essential manner. Right-handed neutrino would correspond to symplectic algebra and other modes to the Kac-Moody algebra and one obtains the crucial 5 tensor factors of Super Virasoro required by p-adic mass calculations.

The matrix elements of WCW metric between Killing vectors are expressible as anti-commutators of super-Hamiltonians identifiable as contractions of WCW gamma matrices with these vectors and give Poisson brackets of corresponding Hamiltonians. The anti-commutation relates of induced spinor fields are

dictated by this condition. Everything is 3-dimensional although one expects that symplectic transformations localized within interior of X^3 act as gauge symmetries so that in this sense effective 2-dimensionality is achieved. The components of WCW metric are labelled by standard model quantum numbers so that the connection with physics is extremely intimate.

4. An open question in the earlier visions was whether finite measurement resolution is realized as discretization at the level of fundamental dynamics. This would mean that only certain string world sheets from the slicing by string world sheets and partonic 2-surfaces are possible. The requirement that anti-commutations are consistent suggests that string world sheets correspond to surfaces for which Kähler magnetic field is constant along string in well defined sense ($J_{\mu\nu}\epsilon^{\mu\nu}g^{1/2}$ remains constant along string). It however turns that by a suitable choice of coordinates of 3-surface one can guarantee that this quantity is constant so that no additional constraint results.
5. Quantum criticality is one of the basic notions of quantum TGD and its relationship to coset construction has remained unclear. In this chapter the concrete realization of criticality in terms of symmetry breaking hierarchy of Super Virasoro algebra acting on symplectic and Kac-Moody algebras. Also a connection with finite measurement resolution - second key notion of TGD - emerges naturally.

The appendix of the book gives a summary about basic concepts of TGD with illustrations. There are concept maps about topics related to the contents of the chapter prepared using CMAP realized as html files. Links to all CMAP files can be found at <http://tgdtheory.fi/cmaphtml.html> [?]. Pdf representation of same files serving as a kind of glossary can be found at <http://tgdtheory.fi/tgdglossary.pdf> [?]. The topics relevant to this chapter are given by the following list.

- Hierarchy of Planck constants [?]
- Hyperfinite factors and TGD [?]
- Structure of WCW [?]
- TGD as infinite-dimensional geometry [?]
- WCW gamma matrices [?]
- WCW spinor fields [?]
- Weak form of electric-magnetic duality [?]
- Zero Energy Ontology (ZEO) [?]
- 4-D spin glass degeneracy [?]
- Equivalence Principle [?]
- Holography [?]

- Quantum Classical Correspondence [?]
- Quantum criticality [?]
- Symmetries of WCW [?]
- TGD as ATQFT [?]
- Vacuum functional in TGD [?]
- KD equation [?]
- Kaehler-Dirac action [?]

2 WCW As A Union Of Homogenous Or Symmetric Spaces

The physical interpretation and detailed mathematical understanding of superconformal symmetries has developed rather slowly and has involved several side tracks. In the following I try to summarize the basic picture with minimal amount of formulas with the understanding that the statement “Noether charge associated with geometrically realized Kac-Moody symmetry” is enough for the reader to write down the needed formula explicitly. Formula oriented reader might deny the value of the approach giving weight to principles. My personal experience is that piles of formulas too often hide the lack of real understanding.

In the following the vision about WCW as union of coset spaces is discussed in more detail.

2.1 Basic Vision

The basic view about coset space construction for WCW has not changed.

1. The idea about WCW as a union of coset spaces G/H labelled by zero modes is extremely attractive. The structure of homogenous space [A1] (http://en.wikipedia.org/wiki/Homogenous_space) means at Lie algebra level the decomposition $g = h \oplus t$ to sub-Lie-algebra h and its complement t such that $[h, t] \subset t$ holds true. Homogeneous spaces have G as its isometries. For symmetric space the additional condition $[t, t] \subset h$ holds true and implies the existence of involution changing at the Lie algebra level the sign of elements of t and leaving the elements of h invariant. The assumption about the structure of symmetric space [A4] (http://en.wikipedia.org/wiki/Symmetric_space) implying covariantly constant curvature tensor is attractive in infinite-dimensional case since it gives hopes about calculability.

An important source of intuition is the analogy with the construction of CP_2 , which is symmetric space. A particular choice of h corresponds to Lie-algebra elements realized as Killing vector fields which vanish at particular point of WCW and thus leave 3-surface invariant. A preferred choice for this point is

as maximum or minimum of Kähler function. For this 3-surface the Hamiltonians of h should be stationary. If symmetric space property holds true then commutators of $[t, t]$ also vanish at the minimum/maximum. Note that Euclidian signature for the metric of WCW requires that Kähler function can have only maximum or minimum for given zero modes.

2. The basic objection against TGD is that one cannot use the powerful canonical quantization using the phase space associated with configuration space - now WCW . The reason is the extreme non-linearity of the Kähler action and its huge vacuum degeneracy, which do not allow the construction of Hamiltonian formalism. Symplectic and Kähler structure must be realized at the level of WCW . In particular, Hamiltonians must be represented in completely new manner. The key idea is to construct WCW Hamiltonians as anti-commutators of super-Hamiltonians defining the contractions of WCW gamma matrices with corresponding Killing vector fields and therefore defining the matrix elements of WCW metric in the tangent vector basis defined by Killing vector fields. Super-symmetry therefore gives hopes about constructing quantum theory in which only induced spinor fields are second quantized and imbedding space coordinates are treated purely classically.
3. It is important to understand the difference between symmetries and isometries assigned to the Kähler function. Symmetries of Kähler function do not affect it. The symmetries of Kähler action are also symmetries of Kähler action because Kähler function is Kähler action for a preferred extremal (here there have been a lot of confusion). Isometries leave invariant only the quadratic form defined by Kähler metric $g_{M\bar{N}} = \partial_M \partial_{\bar{L}} K$ but not Kähler function in general. For G/H decomposition G represents isometries and H both isometries and symmetries of Kähler function.

CP_2 is familiar example: $SU(3)$ represents isometries and $U(2)$ leaves also Kähler function invariant since it depends on the $U(2)$ invariant radial coordinate r of CP_2 . The origin $r = 0$ is left invariant by $U(2)$ but for $r > 0$ $U(2)$ performs a rotation at $r = \text{constant}$ 3-sphere. This simple picture helps to understand what happens at the level of WCW .

How to then distinguish between symmetries and isometries? A natural guess is that one obtains also for the isometries Noether charges but the vanishing of boundary terms at spatial infinity crucial in the argument leading to Noether theorem as $\Delta S = \Delta Q = 0$ does not hold true anymore and one obtains charges which are not conserved anymore. The symmetry breaking contributions would now come from effective boundaries defined by wormhole throats at which the induce metric changes its signature from Minkowskian to Euclidian. A more delicate situation is in which first order contribution to ΔS vanishes and therefore also ΔQ and the contribution to ΔS comes from second variation allowing also to define Noether charge which is not conserved.

4. The simple picture about CP_2 as symmetric space helps to understand the general vision if one assumes that WCW has the structure of symmetric space. The decomposition $g = h + t$ corresponds to decomposition of symplectic

deformations to those which vanish at 3-surface (h) and those which do not (t).

For the symmetric space option, the Poisson brackets for super generators associated with t give Hamiltonians of h identifiable as the matrix elements of WCW metric. They would not vanish although they are stationary at 3-surface meaning that Riemann connection vanishes at 3-surface. The Hamiltonians which vanish at 3-surface X^3 would correspond to t and the Hamiltonians for which Killing vectors vanish and which therefore are stationary at X^3 would correspond to h . Outside X^3 the situation would of course be different. The metric would be obtained by parallel translating the metric from the preferred point of WCW to elsewhere and symplectic transformations would make this parallel translation.

For the homogenous space option the Poisson brackets for super generators of t would still give Hamiltonians identifiable as matrix elements of WCW metric but now they would be necessary those of h . In particular, the Hamiltonians of t do not in general vanish at X^3 .

2.2 Equivalence Principle And WCW

2.3 Ep At Quantum And Classical Level

Quite recently I returned to an old question concerning the meaning of Equivalence Principle (EP) in TGD framework.

Heretic would of course ask whether the question about whether EP is true or not is a pseudo problem due to uncritical assumption there really are two different four-momenta which must be identified. If even the identification of these two different momenta is difficult, the pondering of this kind of problem might be waste of time.

At operational level EP means that the scattering amplitudes mediated by graviton exchange are proportional to the product of four-momenta of particles and that the proportionality constant does not depend on any other parameters characterizing particle (except spin). The are excellent reasons to expect that the stringy picture for interactions predicts this.

1. The old idea is that EP reduces to the coset construction for Super Virasoro algebra using the algebras associated with G and H . The four-momenta assignable to these algebras would be identical from the condition that the differences of the generators annihilate physical states and identifiable as inertial and gravitational momenta. The objection is that for the preferred 3-surface H by definition acts trivially so that time-like translations leading out from the boundary of CD cannot be contained by H unlike G . Hence four-momentum is not associated with the Super-Virasoro representations assignable to H and the idea about assigning EP to coset representations does not look promising.
2. Another possibility is that EP corresponds to quantum classical correspondence (QCC) stating that the classical momentum assignable to Kähler action is identical with gravitational momentum assignable to Super Virasoro representations. This forced to reconsider the questions about the precise iden-

tification of the Kac-Moody algebra and about how to obtain the magic five tensor factors required by p-adic mass calculations [K13].

A more precise formulation for EP as QCC comes from the observation that one indeed obtains two four-momenta in TGD approach. The classical four-momentum assignable to the Kähler action and that assignable to the Kähler-Dirac action. This four-momentum is an operator and QCC would state that given eigenvalue of this operator must be equal to the value of classical four-momentum for the space-time surfaces assignable to the zero energy state in question. In this form EP would be highly non-trivial. It would be justified by the Abelian character of four-momentum so that all momentum components are well-defined also quantum mechanically. One can also consider the splitting of four-momentum to longitudinal and transversal parts as done in the parton model for hadrons: this kind of splitting would be very natural at the boundary of CD. The objection is that this correspondence is nothing more than QCC.

3. A further possibility is that duality of light-like 3-surfaces and space-like 3-surfaces holds true. This is the case if the action of symplectic algebra can be defined at light-like 3-surfaces or even for the entire space-time surfaces. This could be achieved by parallel translation of light-cone boundary providing slicing of CD. The four-momenta associated with the two representations of super-symplectic algebra would be naturally identical and the interpretation would be in terms of EP.

One should also understand how General Relativity and EP emerge at classical level. The understanding comes from the realization that GRT is only an effective theory obtained by endowing M^4 with effective metric.

1. The replacement of superposition of fields with superposition of their effects means replacing superposition of fields with the set-theoretic union of space-time surfaces. Particle experiences sum of the effects caused by the classical fields at the space-time sheets.
2. This is true also for the classical gravitational field defined by the deviation from flat Minkowski metric in standard M^4 coordinates for the space-time sheets. One can define effective metric as sum of M^4 metric and deviations. This effective metric would correspond to that of General Relativity. This resolves long standing issues relating to the interpretation of TGD.
3. Einstein's equations could hold true for the effective metric. They are motivated by the underlying Poincare invariance which cannot be realized as global conservation laws for the effective metric. The conjecture vanishing of divergence of Kähler energy momentum tensor can be seen as the microscopic justification for the claim that Einstein's equations hold true for the effective space-time.
4. The breaking of Poincare invariance could have interpretation as effective breaking in zero energy ontology (ZEO), in which various conserved charges are length dependent and defined separately for each causal diamond (CD).

One can of course consider the possibility that Einstein's equations generalize for preferred extremals of Kähler action. This would actually represent at space-time level the notion of QCC rather than realize QCC interpreted as EP. The condition that the energy momentum tensor for Kähler action has vanishing covariant divergence would be satisfied in GRT if Einstein's equations with cosmological term hold true. This is the case also now but one can consider also more general solutions in which one has two cosmological constants which are not genuine constants anymore [K18].

An interesting question is whether inertial-gravitational duality generalizes to the case of color gauge charges so that color gauge fluxes would correspond to "gravitational" color charges and the charges defined by the conserved currents associated with color isometries would define "inertial" color charges. Since the induced color fields are proportional to color Hamiltonians multiplied by Kähler form they vanish identically for vacuum extremals in accordance with "gravitational" color confinement.

2.4 Criticism Of The Earlier Construction

The earlier detailed realization of super-Hamiltonians and Hamiltonians can be criticized.

1. Even after these more than twenty years it looks strange that the Hamiltonians should reduce to flux integrals over partonic 2-surfaces. The interpretation has been in terms of effective 2-dimensionality suggested strongly by strong form of general coordinate invariance stating that the descriptions based on light-like orbits of partonic 2-surfaces and space-like three surfaces at the ends of causal diamonds are dual so that only partonic 2-surfaces and 4-D tangent space data at them would matter. Strong form of holography implies effective 2-dimensionality but this should correspond gauge character for the action of symplectic generators in the interior the space-like 3-surfaces at the ends of CDs, which is something much milder.

One expects that the strings connecting partonic 2-surfaces could bring something new to the earlier simplistic picture. The guess is that imbedding space Hamiltonian assignable to a point of partonic 2-surface should be replaced with that defined as integral over string attached to the point. Therefore the earlier picture would suffer no modification at the level of general formulas.

2. The fact that the dynamics of Kähler action and Kähler-Dirac action are not directly involved with the earlier construction raises suspicions. I have proposed that Kähler function could allow identification as Dirac determinant [K15] but one would expect more intimate connection. Here the natural question is whether super-Hamiltonians for the Kähler-Dirac action could correspond to Kähler charges constructible using Noether's theorem for corresponding deformations of the space-time surface and would also be identifiable as WCW gamma matrices.

2.5 Is WCW Homogenous Or Symmetric Space?

A key question is whether WCW can be symmetric space [A4] (http://en.wikipedia.org/wiki/Riemannian_symmetric_space) or whether only homogenous structure is needed. The lack of covariant constancy of curvature tensor might produce problems in infinite-dimensional context.

The algebraic conditions for symmetric space are $g = h + t$, $[h, t] \subset t$, $[t, t] \subset h$. The latter condition is the difficult one.

1. δCD Hamiltonians should induce diffeomorphisms of X^3 indeed leaving it invariant. The symplectic vector fields would be parallel to X^3 . A stronger condition is that they induce symplectic transformations for which all points of X^3 remain invariant. Now symplectic vector fields vanish at preferred 3-surface (note that the symplectic transformations are r_M local symplectic transformations of $S^2 \times CP_2$).
2. For Kac-Moody algebra inclusion $H \subset G$ for the finite-dimensional Lie-algebra induces the structure of symmetric space. If entire algebra is involved this does not look physically very attractive idea unless one believes on symmetry breaking for both $SU(3)$, $U(2)_{ew}$, and $SO(3)$ and E_2 (here complex conjugation corresponds to the involution). If one assumes only Kac-Moody algebra as critical symmetries, the number of tensor factors is 4 instead of five, and it is not clear whether one can obtain consistency with p-adic mass calculations.

Examples of 3-surfaces remaining invariant under $U(2)$ are 3-spheres of CP_2 . They could correspond to intersections of deformations of CP_2 type vacuum extremals with the boundary of CD. Also geodesic spheres S^2 of CP_2 are invariant under $U(2)$ subgroup and would relate naturally to cosmic strings. The corresponding 3-surface would be $L \times S^2$, where L is a piece of light-like radial geodesic.

3. In the case of symplectic algebra one can construct the imbedding space Hamiltonians inducing WCW Hamiltonians as products of elements of the isometry algebra of $S^2 \times CP_2$ for with parity under involution is well-defined. This would give a decomposition of the symplectic algebra satisfying the symmetric space property at the level imbedding space. This decomposition does not however look natural at the level of WCW since the only single point of CP_2 and light-like geodesic of δM_+^4 can be fixed by $SO(2) \times U(2)$ so that the 3-surfaces would reduce to pieces of light rays.
4. A more promising involution is the inversion $r_M \rightarrow 1/r_M$ of the radial coordinate mapping the radial conformal weights to their negatives. This corresponds to the inversion in Super Virasoro algebra. t would correspond to functions which are odd functions of $u \equiv \log(r_M/r_0)$ and h to even function of u . Stationary 3-surfaces would correspond to $u = 1$ surfaces for which $\log(u) = 0$ holds true. This would assign criticality with Virasoro algebra as one expects on general grounds.

$r_M = \text{constant}$ surface would most naturally correspond to a maximum of Kähler function which could indeed be highly symmetric. The elements with

even u -parity should define Hamiltonians, which are stationary at the maximum of Kähler function. For other 3-surfaces obtained by (r_M -local) symplectic transformations the situation is different: now H is replaced with its symplectic conjugate hHg^{-1} of H is acceptable and if the conjecture is true one would obtain 3-surfaces assignable to perturbation theory around given maximum as symplectic conjugates of the maximum. The condition that H leaves X^3 invariant in point-wise manner is certainly too strong and imply that the 3-surface has single point as CP_2 projection.

5. One can also consider the possibility that critical deformations correspond to h and non-critical ones to t for the preferred 3-surface. Criticality for given h would hold only for a preferred 3-surface so that this picture would be very similar that above. Symplectic conjugates of h would define criticality for other 3-surfaces. WCW would decompose to a union corresponding to different criticalities perhaps assignable to the hierarchy of sub-algebras of conformal algebra labelled by integer whose multiples give the allowed conformal weights. Hierarchy of breakings of conformal symmetries would characterize this hierarchy of sectors of WCW .

For sub-algebras of the conformal algebras (Kac-Moody and symplectic algebra) the condition $[t, t] \subset h$ cannot hold true so that one would obtain only the structure of homogenous space.

2.6 Symplectic And Kac-Moody Algebras As Basic Building Bricks

3 Updated View About Kähler Geometry Of WCW

During last years the understanding of the mathematical aspects of TGD and of its connection with the experimental world has developed rapidly.

TGD differs in several respects from quantum field theories and string models. The basic mathematical difference is that the mathematically poorly defined notion of path integral is replaced with the mathematically well-defined notion of functional integral defined by the Kähler function defining Kähler metric for WCW (“world of classical worlds”). Apart from quantum jump, quantum TGD is essentially theory of classical WCW spinor fields with WCW spinors represented as fermionic Fock states. One can say that Einstein’s geometrization of physics program is generalized to the level of quantum theory.

It has been clear from the beginning that the gigantic super-conformal symmetries generalizing ordinary super-conformal symmetries are crucial for the existence of WCW Kähler metric. The detailed identification of Kähler function and WCW Kähler metric has however turned out to be a difficult problem. It is now clear that WCW geometry can be understood in terms of the analog of AdS/CFT duality between fermionic and space-time degrees of freedom (or between Minkowskian and Euclidian space-time regions) allowing to express Kähler metric either in terms of Kähler function or in terms of anti-commutators of WCW gamma matrices identifiable as super-conformal Noether super-charges for the symplectic algebra assignable

to $\delta M_{\pm}^4 \times CP_2$. The string model type description of gravitation emerges and also the TGD based view about dark matter becomes more precise. String tension is however dynamical rather than pre-given and the hierarchy of Planck constants is necessary in order to understand the formation of gravitationally bound states. Also the proposal that sparticles correspond to dark matter becomes much stronger: sparticles actually are dark variants of particles.

A crucial element of the construction is the assumption that super-symplectic and other super-conformal symmetries having the same structure as 2-D super-conformal groups can be seen as broken gauge symmetries such that sub-algebra with conformal weights coming as n -ples of those for full algebra act as gauge symmetries. In particular, the Noether charges of this algebra vanish for preferred extremals- this would realize the strong form of holography implied by strong form of General Coordinate Invariance. This gives rise to an infinite number of hierarchies of conformal gauge symmetry breakings with levels labelled by integers $n(i)$ such that $n(i)$ divides $n(i + 1)$ interpreted as hierarchies of dark matter with levels labelled by the value of Planck constant $h_{eff} = n \times h$. These hierarchies define also hierarchies of quantum criticalities, and are proposed to give rise to inclusion hierarchies of hyperfinite factors of II_1 having interpretation in terms of finite cognitive resolution with inclusions being characterized by the integers $n(+1)/n(i)$.

These hierarchies are fundamental for the understanding of living matter. Living matter is fighting in order to stay at criticality and uses metabolic energy and homeostasis to achieve this. In the biological death of the system (self) a phase transition increasing h_{eff} finally takes place. The sub-selves of self experienced by self as mental images however die and are reborn at opposite boundary of the corresponding causal diamond (CD) and they genuinely evolve so that self can be said to become wiser even without dying! The purpose of this fighting against criticality would thus allow a possibility for sub-selves to evolve via subsequent re-incarnations. One interesting prediction is the possibility of time reversed mental images. The challenge is to understand what they do mean at the level of conscious experience.

3.1 Kähler Function, Kähler Action, And Connection With String Models

The definition of Kähler function in terms of Kähler action is possible because space-time regions can have also Euclidian signature of induced metric. Euclidian regions with 4-D CP_2 projection - wormhole contacts - are identified as lines of generalized Feynman diagrams - space-time correlates for basic building bricks of elementary particles. Kähler action from Minkowskian regions is imaginary and gives to the functional integrand a phase factor crucial for quantum field theoretic interpretation. The basic challenges are the precise specification of Kähler function of “world of classical worlds” (WCW) and Kähler metric.

There are two approaches concerning the definition of Kähler metric: the conjecture analogous to AdS/CFT duality is that these approaches are mathematically equivalent.

1. The Kähler function defining Kähler metric can be identified as Kähler action

for space-time regions with Euclidian signature for a preferred extremal containing 3-surface as the ends of the space-time surfaces inside causal diamond (CD). Minkowskian space-time regions give to Kähler action an imaginary contribution interpreted as the counterpart of quantum field theoretic action. The exponent of Kähler function gives rise to a mathematically well-defined functional integral in WCW. WCW metric is dictated by the Euclidian regions of space-time with 4-D CP_2 projection.

The basic question concerns the attribute "preferred". Physically the preferred extremal is analogous to Bohr orbit. What is the mathematical meaning of preferred extremal of Kähler action? The latest step of progress is the realization that the vanishing of generalized conformal charges for the ends of the space-time surface fixes the preferred extremals to high extent and is nothing but classical counterpart for generalized Virasoro and Kac-Moody conditions.

2. Fermions are also needed. The well-definedness of electromagnetic charge led to the hypothesis that spinors are restricted at string world sheets. One could also consider associativity as basic constraint to fermionic dynamics combined with the requirement that octonionic representation for gamma matrices is equivalent with the ordinary one. The conjecture is that this leads to the same outcome. This point is highly non-trivial and will be discussed below separately.
3. Second manner to define Kähler metric is as anticommutators of WCW gamma matrices identified as super-symplectic Noether charges for the Dirac action for induced spinors with string tension proportional to the inverse of Newton's constant. These charges are associated with the 1-D space-like ends of string world sheets connecting the wormhole throats. WCW metric contains contributions from the spinor modes associated with various string world sheets connecting the partonic 2-surfaces associated with the 3-surface.

It is clear that the information carried by WCW metric about 3-surface is rather limited and that the larger the number of string world sheets, the larger the information. This conforms with strong form of holography and the notion of measurement resolution as a property of quantum state. Duality clearly means that Kähler function is determined either by space-time dynamics inside Euclidian wormhole contacts or by the dynamics of fermionic strings in Minkowskian regions outside wormhole contacts. This duality brings strongly in mind AdS/CFT duality. One could also speak about fermionic emergence since Kähler function is dictated by the Kähler metric part from a real part of gradient of holomorphic function: a possible identification of the exponent of Kähler function is as Dirac determinant.

3.2 Realization Of Super-Conformal Symmetries

The detailed realization of various super-conformal symmetries has been also a long standing problem.

1. Super-conformal symmetry requires that Dirac action for string world sheets is accompanied by string world sheet area as part of bosonic action. String

world sheets are implied and can be present only in Minkowskian regions if one demands that octonionic and ordinary representations of induced spinor structure are equivalent (this requires vanishing of induced spinor curvature to achieve associativity in turn implying that CP_2 projection is 1-D). Note that 1-dimensionality of CP_2 projection is symplectically invariant property. Kähler action is not invariant under symplectic transformations. This is necessary for having non-trivial Kähler metric. Whether WCW really possesses super-symplectic isometries remains an open problem.

2. Super-conformal symmetry also demands that Kähler action is accompanied by what I call Kähler-Dirac action with gamma matrices defined by the contractions of the canonical momentum currents with imbedding space-gamma matrices. Both the well-definedness of em charge and equivalence of octonionic spinor dynamics with ordinary one require the restriction of spinor modes to string world sheets with light-like boundaries at wormhole throats. K-D action with the localization of induced spinors at string world sheets is certainly the minimal option to consider.
3. Strong form of holography implied by strong form of general coordinate invariance strongly suggests that super-conformal symmetry is broken gauge invariance in the sense that the classical super-conformal charges for a sub-algebra of the symplectic algebra with conformal weights vanishing modulo some integer n vanish. The proposal is that n corresponds to the effective Planck constant as $h_{eff}/h = n$. The standard conformal symmetries for spinors modes at string world sheets is always unbroken gauge symmetry.

3.3 Interior Dynamics For Fermions, The Role Of Vacuum Extremals, And Dark Matter

The key role of CP_2 -type and M^4 -type vacuum extremals has been rather obvious from the beginning but the detailed understanding has been lacking. Both kinds of extremals are invariant under symplectic transformations of $\delta M^4 \times CP_2$, which inspires the idea that they give rise to isometries of WCW. The deformations CP_2 -type extremals correspond to lines of generalized Feynman diagrams. M^4 type vacuum extremals in turn are excellent candidates for the building bricks of many-sheeted space-time giving rise to GRT space-time as approximation. For M^4 type vacuum extremals CP_2 projection is (at most 2-D) Lagrangian manifold so that the induced Kähler form vanishes and the action is fourth-order in small deformations. This implies the breakdown of the path integral approach and of canonical quantization, which led to the notion of WCW.

If the action in Minkowskian regions contains also string area, the situation changes dramatically since strings dominate the dynamics in excellent approximation and string theory should give an excellent description of the situation: this of course conforms with the dominance of gravitation.

String tension would be proportional to $1/\hbar G$ and this raises a grave classical counter argument. In string model massless particles are regarded as strings, which have contracted to a point in excellent approximation and cannot have length longer

than Planck length. How this can be consistent with the formation of gravitationally bound states is however not understood since the required non-perturbative formulation of string model required by the large valued of the coupling parameter GMm is not known.

In TGD framework strings would connect even objects with macroscopic distance and would obviously serve as correlates for the formation of bound states in quantum level description. The classical energy of string connecting say the two wormhole contacts defining elementary particle is gigantic for the ordinary value of \hbar so that something goes wrong.

I have however proposed [K9, K8, K20] that gravitons - at least those mediating interaction between dark matter have large value of Planck constant. I talk about gravitational Planck constant and one has $\hbar_{eff} = \hbar_{gr} = GMm/v_0$, where $v_0/c < 1$ (v_0 has dimensions of velocity). This makes possible perturbative approach to quantum gravity in the case of bound states having mass larger than Planck mass so that the parameter GMm analogous to coupling constant is very large. The velocity parameter v_0/c becomes the dimensionless coupling parameter. This reduces the string tension so that for string world sheets connecting macroscopic objects one would have $T \propto v_0/G^2Mm$. For $v_0 = GMm/\hbar$, which remains below unity for Mm/m_{Pl}^2 one would have $\hbar_{gr}/\hbar = 1$. Hence action remains small and its imaginary exponent does not fluctuate wildly to make the bound state forming part of gravitational interaction short ranged. This is expected to hold true for ordinary matter in elementary particle scales. The objects with size scale of large neutron ($100 \mu\text{m}$ in the density of water) - probably not an accident - would have mass above Planck mass so that dark gravitons and also life would emerge as massive enough gravitational bound states are formed. $\hbar_{gr} = \hbar_{eff}$ hypothesis is indeed central in TGD based view about living matter.

If one assumes that for non-standard values of Planck constant only n -multiples of super-conformal algebra in interior annihilate the physical states, interior conformal gauge degrees of freedom become partly dynamical. The identification of dark matter as macroscopic quantum phases labeled by $\hbar_{eff}/\hbar = n$ conforms with this.

The emergence of dark matter corresponds to the emergence of interior dynamics via breaking of super-conformal symmetry. The induced spinor fields in the interior of flux tubes obeying Kähler Dirac action should be highly relevant for the understanding of dark matter. The assumption that dark particles have essentially same masses as ordinary particles suggests that dark fermions correspond to induced spinor fields at both string world sheets and in the space-time interior: the spinor fields in the interior would be responsible for the long range correlations characterizing $\hbar_{eff}/\hbar = n$. Magnetic flux tubes carrying dark matter are key entities in TGD inspired quantum biology. Massless extremals represent second class of M^4 type non-vacuum extremals.

This view forces once again to ask whether space-time SUSY is present in TGD and how it is realized. With a motivation coming from the observation that the mass scales of particles and sparticles most naturally have the same p-adic mass scale as particles in TGD Universe I have proposed that sparticles might be dark in TGD sense. The above argument leads to ask whether the dark variants of particles correspond to states in which one has ordinary fermion at string world sheet and 4-D fermion in the space-time interior so that dark matter in TGD sense would almost

by definition correspond to sparticles!

3.4 Classical Number Fields And Associativity And Commutativity As Fundamental Law Of Physics

The dimensions of classical number fields appear as dimensions of basic objects in quantum TGD. Imbedding space has dimension 8, space-time has dimension 4, light-like 3-surfaces are orbits of 2-D partonic surfaces. If conformal QFT applies to 2-surfaces (this is questionable), one-dimensional structures would be the basic objects. The lowest level would correspond to discrete sets of points identifiable as intersections of real and p-adic space-time sheets. This suggests that besides p-adic number fields also classical number fields (reals, complex numbers, quaternions, octonions [A9]) are involved [K11] and the notion of geometry generalizes considerably. In the recent view about quantum TGD the dimensional hierarchy defined by classical number field indeed plays a key role. $H = M^4 \times CP_2$ has a number theoretic interpretation and standard model symmetries can be understood number theoretically as symmetries of hyper-quaternionic planes of hyper-octonionic space.

The associativity condition $A(BC) = (AB)C$ suggests itself as a fundamental physical law of both classical and quantum physics. Commutativity can be considered as an additional condition. In conformal field theories associativity condition indeed fixes the n-point functions of the theory. At the level of classical TGD space-time surfaces could be identified as maximal associative (hyper-quaternionic) sub-manifolds of the imbedding space whose points contain a preferred hyper-complex plane M^2 in their tangent space and the hierarchy finite fields-rationals-reals-complex numbers-quaternions-octonions could have direct quantum physical counterpart [K11]. This leads to the notion of number theoretic compactification analogous to the dualities of M-theory: one can interpret space-time surfaces either as hyper-quaternionic 4-surfaces of M^8 or as 4-surfaces in $M^4 \times CP_2$. As a matter fact, commutativity in number theoretic sense is a further natural condition and leads to the notion of number theoretic braid naturally as also to direct connection with super string models.

At the level of Kähler-Dirac action the identification of space-time surface as a hyper-quaternionic sub-manifold of H means that the modified gamma matrices of the space-time surface defined in terms of canonical momentum currents of Kähler action using octonionic representation for the gamma matrices of H span a hyper-quaternionic sub-space of hyper-octonions at each point of space-time surface (hyper-octonions are the subspace of complexified octonions for which imaginary units are octonionic imaginary units multiplied by commuting imaginary unit). Hyper-octonionic representation leads to a proposal for how to extend twistor program to TGD framework [K15, K12].

3.4.1 *How to achieve associativity in the fermionic sector?*

In the fermionic sector an additional complication emerges. The associativity of the tangent- or normal space of the space-time surface need not be enough to guarantee the associativity at the level of Kähler-Dirac or Dirac equation. The reason is the presence of spinor connection. A possible cure could be the vanishing of the

components of spinor connection for two conjugates of quaternionic coordinates combined with holomorphy of the modes.

1. The induced spinor connection involves sigma matrices in CP_2 degrees of freedom, which for the octonionic representation of gamma matrices are proportional to octonion units in Minkowski degrees of freedom. This corresponds to a reduction of tangent space group $SO(1,7)$ to G_2 . Therefore octonionic Dirac equation identifying Dirac spinors as complexified octonions can lead to non-associativity even when space-time surface is associative or co-associative.
2. The simplest manner to overcome these problems is to assume that spinors are localized at 2-D string world sheets with 1-D CP_2 projection and thus possible only in Minkowskian regions. Induced gauge fields would vanish. String world sheets would be minimal surfaces in $M^4 \times D^1 \subset M^4 \times CP_2$ and the theory would simplify enormously. String area would give rise to an additional term in the action assigned to the Minkowskian space-time regions and for vacuum extremals one would have only strings in the first approximation, which conforms with the success of string models and with the intuitive view that vacuum extremals of Kähler action are basic building bricks of many-sheeted space-time. Note that string world sheets would be also symplectic covariants.

Without further conditions gauge potentials would be non-vanishing but one can hope that one can gauge transform them away in associative manner. If not, one can also consider the possibility that CP_2 projection is geodesic circle S^1 : symplectic invariance is considerably reduces for this option since symplectic transformations must reduce to rotations in S^1 .

3. The first heavy objection is that action would contain Newton's constant G as a fundamental dynamical parameter: this is a standard recipe for building a non-renormalizable theory. The very idea of TGD indeed is that there is only single dimensionless parameter analogous to critical temperature. One can of course argue that the dimensionless parameter is $\hbar G/R^2$, " R CP_2 "radius".

Second heavy objection is that the Euclidian variant of string action exponentially damps out all string world sheets with area larger than $\hbar G$. Note also that the classical energy of Minkowskian string would be gigantic unless the length of string is of order Planck length. For Minkowskian signature the exponent is oscillatory and one can argue that wild oscillations have the same effect.

The hierarchy of Planck constants would allow the replacement $\hbar \rightarrow \hbar_{eff}$ but this is not enough. The area of typical string world sheet would scale as \hbar_{eff} and the size of CD and gravitational Compton lengths of gravitationally bound objects would scale as $\sqrt{\hbar_{eff}}$ rather than $\hbar_{eff} = GMm/v_0$, which one wants. The only way out of problem is to assume $T \propto (\hbar/\hbar_{eff})^2 \times (1/\hbar_{bar}G)$. This is however un-natural for genuine area action. Hence it seems that the visit of the basic assumption of superstring theory to TGD remains very short.

3.4.2 *Is super-symmetrized Kähler-Dirac action enough?*

Could one do without string area in the action and use only K-D action, which is in any case forced by the super-conformal symmetry? This option I have indeed considered hitherto. K-D Dirac equation indeed tends to reduce to a lower-dimensional one: for massless extremals the K-D operator is effectively 1-dimensional. For cosmic strings this reduction does not however take place. In any case, this leads to ask whether in some cases the solutions of Kähler-Dirac equation are localized at lower-dimensional surfaces of space-time surface.

1. The proposal has indeed been that string world sheets carry vanishing W and possibly even Z fields: in this manner the electromagnetic charge of spinor mode could be well-defined. The vanishing conditions force in the generic case 2-dimensionality.

Besides this the canonical momentum currents for Kähler action defining 4 imbedding space vector fields must define an integrable distribution of two planes to give string world sheet. The four canonical momentum currents $\Pi_k \alpha = \partial L_K / \partial_{\partial_\alpha h^k}$ identified as imbedding 1-forms can have only two linearly independent components parallel to the string world sheet. Also the Frobenius conditions stating that the two 1-forms are proportional to gradients of two imbedding space coordinates Φ_i defining also coordinates at string world sheet, must be satisfied. These conditions are rather strong and are expected to select some discrete set of string world sheets.

2. To construct preferred extremal one should fix the partonic 2-surfaces, their light-like orbits defining boundaries of Euclidian and Minkowskian space-time regions, and string world sheets. At string world sheets the boundary condition would be that the normal components of canonical momentum currents for Kähler action vanish. This picture brings in mind strong form of holography and this suggests that might make sense and also solution of Einstein equations with point like sources.
3. The localization of spinor modes at 2-D surfaces would follow from the well-definedness of em charge and one could have situation in which the localization does not occur. For instance, covariantly constant right-handed neutrinos spinor modes at cosmic strings are completely de-localized and one can wonder whether one could give up the localization inside wormhole contacts.
4. String tension is dynamical and physical intuition suggests that induced metric at string world sheet is replaced by the anti-commutator of the K-D gamma matrices and by conformal invariance only the conformal equivalence class of this metric would matter and it could be even equivalent with the induced metric. A possible interpretation is that the energy density of Kähler action has a singularity localized at the string world sheet.

Another interpretation that I proposed for years ago but gave up is that in spirit with the TGD analog of AdS/CFT duality the Noether charges for

Kähler action can be reduced to integrals over string world sheet having interpretation as area in effective metric. In the case of magnetic flux tubes carrying monopole fluxes and containing a string connecting partonic 2-surfaces at its ends this interpretation would be very natural, and string tension would characterize the density of Kähler magnetic energy. String model with dynamical string tension would certainly be a good approximation and string tension would depend on scale of CD.

5. There is also an objection. For M^4 type vacuum extremals one would not obtain any non-vacuum string world sheets carrying fermions but the successes of string model strongly suggest that string world sheets are there. String world sheets would represent a deformation of the vacuum extremal and far from string world sheets one would have vacuum extremal in an excellent approximation. Situation would be analogous to that in general relativity with point particles.
6. The hierarchy of conformal symmetry breakings for K-D action should make string tension proportional to $1/h_{eff}^2$ with $h_{eff} = h_{gr}$ giving correct gravitational Compton length $\Lambda_{gr} = GM/v_0$ defining the minimal size of CD associated with the system. Why the effective string tension of string world sheet should behave like $(\hbar/\hbar_{eff})^2$?

The first point to notice is that the effective metric $G^{\alpha\beta}$ defined as $h^{kl}\Pi_k^\alpha\Pi_l^\beta$, where the canonical momentum current $\Pi_k\alpha = \partial L_K/\partial_{\partial_\alpha}h^k$ has dimension $1/L^2$ as required. Kähler action density must be dimensionless and since the induced Kähler form is dimensionless the canonical momentum currents are proportional to $1/\alpha_K$.

Should one assume that α_K is fundamental coupling strength fixed by quantum criticality to $\alpha_K = 1/137$? Or should one regard g_K^2 as fundamental parameter so that one would have $1/\alpha_K = \hbar_{eff}/4\pi g_K^2$ having spectrum coming as integer multiples (recall the analogy with inverse of critical temperature)?

The latter option is the in spirit with the original idea stating that the increase of h_{eff} reduces the values of the gauge coupling strengths proportional to α_K so that perturbation series converges (Universe is theoretician friendly). The non-perturbative states would be critical states. The non-determinism of Kähler action implying that the 3-surfaces at the boundaries of CD can be connected by large number of space-time sheets forming n conformal equivalence classes. The latter option would give $G^{\alpha\beta} \propto h_{eff}^2$ and $det(G) \propto 1/h_{eff}^2$ as required.

7. It must be emphasized that the string tension has interpretation in terms of gravitational coupling on only at the GRT limit of TGD involving the replacement of many-sheeted space-time with single sheeted one. It can have also interpretation as hadronic string tension or effective string tension associated with magnetic flux tubes and telling the density of Kähler magnetic energy per unit length.

Superstring models would describe only the perturbative Planck scale dynamics for emission and absorption of $h_{eff}/h = 1$ on mass shell gravitons whereas the quantum description of bound states would require $h_{eff}/n > 1$ when the

masses. Also the effective gravitational constant associated with the strings would differ from G .

The natural condition is that the size scale of string world sheet associated with the flux tube mediating gravitational binding is $G(M + m)/v_0$, By expressing string tension in the form $1/T = n^2 \hbar G_1$, $n = h_{eff}/\hbar$, this condition gives $\hbar G_1 = \hbar^2/M_{red}^2$, $M_{red} = Mm/(M + m)$. The effective Planck length defined by the effective Newton's constant G_1 analogous to that appearing in string tension is just the Compton length associated with the reduced mass of the system and string tension equals to $T = [v_0/G(M + m)]^2$ apart from a numerical constant ($2G(M + m)$ is Schwarzschild radius for the entire system). Hence the macroscopic stringy description of gravitation in terms of string differs dramatically from the perturbative one. Note that one can also understand why in the Bohr orbit model of Nottale [?] for the planetary system and in its TGD version [K9] v_0 must be by a factor 1/5 smaller for outer planets rather than inner planets.

3.4.3 Are 4-D spinor modes consistent with associativity?

The condition that octonionic spinors are equivalent with ordinary spinors looks rather natural but in the case of Kähler-Dirac action the non-associativity could leak in. One could of course give up the condition that octonionic and ordinary K-D equation are equivalent in 4-D case. If so, one could see K-D action as related to non-commutative and maybe even non-associative fermion dynamics. Suppose that one does not.

1. K-D action vanishes by K-D equation. Could this save from non-associativity? If the spinors are localized to string world sheets, one obtains just the standard stringy construction of conformal modes of spinor field. The induced spinor connection would have only the holomorphic component A_z . Spinor mode would depend only on z but K-D gamma matrix Γ^z would annihilate the spinor mode so that K-D equation would be satisfied. There are good hopes that the octonionic variant of K-D equation is equivalent with that based on ordinary gamma matrices since quaternionic coordinated reduces to complex coordinate, octonionic quaternionic gamma matrices reduce to complex gamma matrices, sigma matrices are effectively absent by holomorphy.
2. One can consider also 4-D situation (maybe inside wormhole contacts). Could some form of quaternion holomorphy [A13] [K12] allow to realize the K-D equation just as in the case of super string models by replacing complex coordinate and its conjugate with quaternion and its 3 conjugates. Only two quaternion conjugates would appear in the spinor mode and the corresponding quaternionic gamma matrices would annihilate the spinor mode. It is essential that in a suitable gauge the spinor connection has non-vanishing components only for two quaternion conjugate coordinates. As a special case one would have a situation in which only one quaternion coordinate appears in the solution. Depending on the character of quaternion holomorphy the modes would be labelled by one or two integers identifiable as conformal weights.

Even if these octonionic 4-D modes exists (as one expects in the case of cosmic strings), it is far from clear whether the description in terms of them is equivalent with the description using K-D equation based ordinary gamma matrices. The algebraic structure however raises hopes about this. The quaternion coordinate can be represented as sum of two complex coordinates as $q = z_1 + Jz_2$ and the dependence on two quaternion conjugates corresponds to the dependence on two complex coordinates z_1, z_2 . The condition that two quaternion complexified gammas annihilate the spinors is equivalent with the corresponding condition for Dirac equation formulated using 2 complex coordinates. This for wormhole contacts. The possible generalization of this condition to Minkowskian regions would be in terms Hamilton-Jacobi structure.

Note that for cosmic strings of form $X^2 \times Y^2 \subset M^4 \times CP_2$ the associativity condition for S^2 sigma matrix and without assuming localization demands that the commutator of Y^2 imaginary units is proportional to the imaginary unit assignable to X^2 which however depends on point of X^2 . This condition seems to imply correlation between Y^2 and S^2 which does not look physical.

To summarize, the minimal and mathematically most optimistic conclusion is that Kähler-Dirac action is indeed enough to understand gravitational binding without giving up the associativity of the fermionic dynamics. Conformal spinor dynamics would be associative if the spinor modes are localized at string world sheets with vanishing W (and maybe also Z) fields guaranteeing well-definedness of em charge and carrying canonical momentum currents parallel to them. It is not quite clear whether string world sheets are present also inside wormhole contacts: for CP_2 type vacuum extremals the Dirac equation would give only right-handed neutrino as a solution (could they give rise to $N = 2$ SUSY?).

The construction of preferred extremals would realize strong form of holography. By conformal symmetry the effective metric at string world sheet could be conformally equivalent with the induced metric at string world sheets.

Dynamical string tension would be proportional to \hbar/h_{eff}^2 due to the proportionality $\alpha_K \propto 1/h_{eff}$ and predict correctly the size scales of gravitationally bound states for $\hbar_{gr} = \hbar_{eff} = GMm/v_0$. Gravitational constant would be a prediction of the theory and be expressible in terms of α_K and R^2 and \hbar_{eff} ($G \propto R^2/g_K^2$).

In fact, all bound states - elementary particles as pairs of wormhole contacts, hadronic strings, nuclei [K6], molecules, etc. - are described in the same manner quantum mechanically. This is of course nothing new since magnetic flux tubes associated with the strings provide a universal model for interactions in TGD Universe. This also conforms with the TGD counterpart of AdS/CFT duality.

The basic building bricks are symplectic algebra of δCD (this includes CP_2 besides light-cone boundary) and Kac-Moody algebra assignable to the isometries of δCD [K3]. It seems however that the longheld view about the role of Kac-Moody algebra must be modified. Also the earlier realization of super-Hamiltonians and Hamiltonians seems too naive.

1. I have been accustomed to think that Kac-Moody algebra could be regarded as a sub-algebra of symplectic algebra. p-Adic mass calculations however requires

five tensor factors for the coset representation of Super Virasoro algebra naturally assigned to the coset structure G/H of a sector of WCW with fixed zero modes. Therefore Kac-Moody algebra cannot be regarded as a sub-algebra of symplectic algebra giving only single tensor factor and thus inconsistent with interpretation of p-adic mass calculations.

2. The localization of Kac-Moody algebra generators with respect to the internal coordinates of light-like 3-surface taking the role of complex coordinate z in conformal field theory is also questionable: the most economical option relies on localization with respect to light-like radial coordinate of light-cone boundary as in the case of symplectic algebra. Kac-Moody algebra cannot be however sub-algebra of the symplectic algebra assigned with covariantly constant right-handed neutrino in the earlier approach.
3. Right-handed covariantly constant neutrino as a generator of super symmetries plays a key role in the earlier construction of symplectic super-Hamiltonians. What raises doubts is that other spinor modes - both those of right-handed neutrino and electro-weakly charged spinor modes - are absent. All spinor modes should be present and thus provide direct mapping from WCW geometry to WCW spinor fields in accordance with super-symmetry and the general idea that WCW geometry is coded by WCW spinor fields.

Hence it seems that Kac-Moody algebra must be assigned with the modes of the induced spinor field which carry electroweak quantum numbers. It would be natural that the modes of right-handed neutrino having no weak and color interactions would generate the huge symplectic algebra of symmetries and that the modes of fermions with electroweak charges generate much smaller Kac-Moody algebra.

4. The dynamics of Kähler action and Kähler-Dirac action are invisible in the earlier construction. This suggests that the definition of WCW Hamiltonians is too simplistic. The proposal is that the conserved super charges derivable as Noether charges and identifiable as super-Hamiltonians define WCW metric and Hamiltonians as their anti-commutators. Spinor modes would become labels of Hamiltonians and WCW geometry relates directly to the dynamics of elementary particles.
5. Note that light-cone boundary $\delta M_+^4 = S^2 \times R_+$ allows infinite-dimensional group of isometries consisting of conformal transformation of the sphere S^2 with conformal scaling compensated by an S^2 local scaling or the light-like radial coordinate of R_+ . These isometries contain as a subgroup symplectic isometries and could act as gauge symmetries of the theory.

4 About some unclear issues of TGD

TGD has been in the middle of palace revolution during last two years and it is almost impossible to keep the chapters of the books updated. Adelic vision and twistor lift of TGD are the newest developments and there are still many details

to be understood and errors to be corrected. The description of fermions in TGD framework has contained some unclear issues. Hence the motivation for the following brief comments.

There questions about the adelic vision about symmetries. Do the cognitive representations implying number theoretic discretization of the space-time surface lead to the breaking of the basic symmetries and are preferred imbedding space coordinates actually necessary?

In the fermionic sector there are many questions deserving clarification. How quantum classical correspondence (QCC) is realized for fermions? How is SH realized for fermions and how does it lead to the reduction of dimension $D = 4$ to $D = 2$ (apart from number theoretical discretization)? Can scattering amplitudes be really formulated by using only the data at the boundaries of string sheets and what does this mean from the point of view of the modified Dirac equation? Are the spinors at light-like boundaries limiting values or sources? A long-standing issue concerns the fermionic anti-commutation relations: what motivated this article was the solution of this problem. There is also the general problem about whether statistical entanglement is “real”.

4.1 Adelic vision and symmetries

In the adelic TGD SH is weakened: also the points of the space-time surface having imbedding space coordinates in an extension of rationals (cognitive representation) are needed so that data are not precisely 2-D. I have believed hitherto that one must use preferred coordinates for the imbedding space H - a subset of these coordinates would define space-time coordinates. These coordinates are determined apart from isometries. Does the number theoretic discretization imply loss of general coordinate invariance and also other symmetries?

The reduction of symmetry groups to their subgroups (not only algebraic since powers of e define finite-dimensional extension of p-adic numbers since e^p is ordinary p-adic number) is genuine loss of symmetry and reflects finite cognitive resolution. The physics itself has the symmetries of real physics.

The assumption about preferred imbedding space coordinates is actually not necessary. Different choices of H -coordinates means only different and non-equivalent cognitive representations. Spherical and linear coordinates in finite accuracy do not provide equivalent representations.

4.2 Quantum-classical correspondence for fermions

Quantum-classical correspondence (QCC) for fermions is rather well-understood but deserves to be mentioned also here.

QCC for fermions means that the space-time surface as preferred extremal should depend on fermionic quantum numbers. This is indeed the case if one requires QCC in the sense that the fermionic representations of Noether charges in the Cartan algebras of symmetry algebras are equal to those to the classical Noether charges for preferred extremals.

Second aspect of QCC becomes visible in the representation of fermionic states as point like particles moving along the light-like curves at the light-like orbits of the

partonic 2-surfaces (curve at the orbit can be locally only light-like or space-like). The number of fermions and antifermions dictates the number of string world sheets carrying the data needed to fix the preferred extremal by SH. The complexity of the space-time surface increases as the number of fermions increases.

4.3 Strong form of holography for fermions

It seems that scattering amplitudes can be formulated by assigning fermions with the boundaries of strings defining the lines of twistor diagrams [K23, K24]. This information theoretic dimensional reduction from $D = 4$ to $D = 2$ for the scattering amplitudes can be partially understood in terms of strong form of holography (SH): one can construct the theory by using the data at string worlds sheets and/or partonic 2-surfaces at the ends of the space-time surface at the opposite boundaries of causal diamond (CD).

4-D modified Dirac action would appear at fundamental level as supersymmetry demands but would be reduced for preferred extremals to its 2-D stringy variant serving as effective action. Also the value of the 4-D action determining the space-time dynamics would reduce to effective stringy action containing area term, 2-D Kähler action, and topological Kähler magnetic flux term. This reduction would be due to the huge gauge symmetries of preferred extremals. Sub-algebra of super-symplectic algebra with conformal weights coming as n -multiples of those for the entire algebra and the commutators of this algebra with the entire algebra would annihilate the physical states, and the corresponding classical Noether charges would vanish.

One still has the question why not the data at the entire string world sheets is not needed to construct scattering amplitudes. Scattering amplitudes of course need not code for the entire physics. QCC is indeed motivated by the fact that quantum experiments are always interpreted in terms of classical physics, which in TGD framework reduces to that for space-time surface.

4.4 The relationship between spinors in space-time interior and at boundaries between Euclidian and Minkoskian regions

Space-time surface decomposes to interiors of Minkowskian and Euclidian regions. At light-like 3-surfaces at which the four-metric changes, the 4-metric is degenerate. These metrically singular 3-surfaces - partonic orbits- carry the boundaries of string world sheets identified as carriers of fermionic quantum numbers. The boundaries define fermion lines in the twistor lift of TGD [K23, K24]. The relationship between fermions at the partonic orbits and interior of the space-time surface has however remained somewhat enigmatic.

So: What is the precise relationship between induced spinors Ψ_B at light-like partonic 3-surfaces and Ψ_I in the interior of Minkowskian and Euclidian regions? Same question can be made for the spinors Ψ_B at the boundaries of string world sheets and Ψ_I in interior of the string world sheets. There are two options to consider:

- Option I: Ψ_B is the limiting value of Ψ_I .
- Option II: Ψ_B serves as a source of Ψ_I .

For the Option I it is difficult to understand the preferred role of Ψ_B .
I have considered Option II already years ago but have not been able to decide.

1. That scattering amplitudes could be formulated only in terms of sources only, would fit nicely with SH, twistorial amplitude construction, and also with the idea that scattering amplitudes in gauge theories can be formulated in terms of sources of boson fields assignable to vertices and propagators. Now the sources would become fermionic.
2. One can take gauge theory as a guideline. One adds to free Dirac equation source term ${}_k\Psi$. Therefore the natural boundary term in the action would be of the form (forgetting overall scale factor)

$$S_B = \bar{\Psi}_I \Gamma^\alpha (C - S) A_\alpha \Psi_B + c.c. .$$

Here the modified gamma matrix is $\Gamma^\alpha (C - S)$ (contravariant form is natural for light-like 3-surfaces) is most naturally defined by the boundary part of the action - naturally Chern-Simons term for Kähler action. A denotes the Kähler gauge potential.

3. The variation with respect to Ψ_B gives

$$G^\alpha (C - S) A_\alpha \Psi_I = 0$$

at the boundary so that the C-S term and interaction term vanish. This does not however imply vanishing of the source term! This condition can be seen as a boundary condition.

The same argument applies also to string world sheets.

4.5 About second quantization of the induced spinor fields

The anti-commutation relations for the induced spinors have been a long-standing issue and during years I have considered several options. The solution of the problem looks however stupidly simple. The conserved fermion currents are accompanied by super-currents obtained by replacing Ψ with a mode of the induced spinor field to get $\bar{u}_n \Gamma^\alpha \Psi$ or $\bar{\Psi} \Gamma^\alpha u_n$ with the conjugate of the mode. One obtains infinite number of conserved super currents. One can also replace both Ψ and $\bar{\Psi}$ in this manner to get purely bosonic conserved currents $\bar{u}_m \Gamma^\alpha u_n$ to which one can assign a conserved bosonic charges Q_{mn} .

I noticed this years ago but did not realize that these bosonic charges define naturally anti-commutators of fermionic creation and annihilation operators! The ordinary anti-commutators of quantum field theory follow as a special case! By a suitable unitary transformation of the spinor basis one can diagonalize the hermitian

matrix defined by Q_{mn} and by performing suitable scalings one can transform anti-commutation relations to the standard form. An interesting question is whether the diagonalization is needed, and whether the deviation of the diagonal elements from unity could have some meaning and possibly relate to the hierarchy $h_{eff} = n \times h$ of Planck constants - probably not.

4.6 Is statistical entanglement “real” entanglement?

The question about the “reality” of statistical entanglement has bothered me for years. This entanglement is maximal and it cannot be reduced by measurement so that one can argue that it is not “real”. Quite recently I learned that there has been a longstanding debate about the statistical entanglement and that the issue still remains unresolved.

The idea that all electrons of the Universe are maximally entangled looks crazy. TGD provides several variants for solutions of this problem. It could be that only the fermionic oscillator operators at partonic 2-surfaces associated with the space-time surface (or its connected component) inside given CD anti-commute and the fermions are thus indistinguishable. The extremist option is that the fermionic oscillator operators belonging to a network of partonic 2-surfaces connected by string world sheets anti-commute: only the oscillator operators assignable to the same scattering diagram would anti-commute.

What about QCC in the case of entanglement. ER-EPR correspondence introduced by Maldacena and Susskind for 4 years ago proposes that blackholes (maybe even elementary particles) are connected by wormholes. In TGD the analogous statement emerged for more than decade ago - magnetic flux tubes take the role of wormholes in TGD. Magnetic flux tubes were assumed to be accompanied by string world sheets. I did not consider the question whether string world sheets are always accompanied by flux tubes.

What could be the criterion for entanglement to be “real”? “Reality” of entanglement demands some space-time correlate. Could the presence of the flux tubes make the entanglement “real”? If statistical entanglement is accompanied by string connections without magnetic flux tubes, it would not be “real”: only the presence of flux tubes would make it “real”. Or is the presence of strings enough to make the statistical entanglement “real”. In both cases the fermions associated with disjoint space-time surfaces or with disjoint CDs would not be indistinguishable. This looks rather sensible.

The space-time correlate for the reduction of entanglement would be the splitting of a flux tube and fermionic strings inside it. The fermionic strings associated with flux tubes carrying monopole flux are closed and the return flux comes back along parallel space-time sheet. Also fermionic string has similar structure. Reconnection of this flux tube with shape of very long flattened square splitting it to two pieces would be the correlate for the state function reduction reducing the entanglement with other fermions and would indeed decouple the fermion from the network.

5 About The Notion Of Four-Momentum In TGD Framework

The starting point of TGD was the energy problem of General Relativity [K13]. The solution of the problem was proposed in terms of sub-manifold gravity and based on the lifting of the isometries of space-time surface to those of $M^4 \times CP_2$ in which space-times are realized as 4-surfaces so that Poincare transformations act on space-time surface as an 4-D analog of rigid body rather than moving points at space-time surface. It however turned out that the situation is not at all so simple.

There are several conceptual hurdles and I have considered several solutions for them. The basic source of problems has been Equivalence Principle (EP): what does EP mean in TGD framework [K13, K18] ? A related problem has been the interpretation of gravitational and inertial masses, or more generally the corresponding 4-momenta. In General Relativity based cosmology gravitational mass is not conserved and this seems to be in conflict with the conservation of Noether charges. The resolution is in terms of zero energy ontology (ZEO), which however forces to modify slightly the original view about the action of Poincare transformations.

A further problem has been quantum classical correspondence (QCC): are quantal four-momenta associated with super conformal representations and classical four-momenta associated as Noether charges with Kähler action for preferred extremals identical? Could inertial-gravitational duality - that is EP - be actually equivalent with QCC? Or are EP and QCC independent dualities. A powerful experimental input comes p-adic mass calculations [K17] giving excellent predictions provided the number of tensor factors of super-Virasoro representations is five, and this input together with Occam's razor strongly favors QCC=EP identification.

There is also the question about classical realization of EP and more generally, TGD-GRT correspondence.

Twistor Grassmannian approach has meant a technical revolution in quantum field theory (for attempts to understand and generalize the approach in TGD framework see [K12]). This approach seems to be extremely well suited to TGD and I have considered a generalization of this approach from $\mathcal{N} = 4$ SUSY to TGD framework by replacing point like particles with string world sheets in TGD sense and super-conformal algebra with its TGD version: the fundamental objects are now massless fermions which can be regarded as on mass shell particles also in internal lines (but with unphysical helicity). The approach solves old problems related to the realization of stringy amplitudes in TGD framework, and avoids some problems of twistorial QFT (IR divergences and the problems due to non-planar diagrams). The Yangian [A5] [?, ?, ?] variant of 4-D conformal symmetry is crucial for the approach in $\mathcal{N} = 4$ SUSY, and implies the recently introduced notion of amplituhedron [?]. A Yangian generalization of various super-conformal algebras seems more or less a "must" in TGD framework. As a consequence, four-momentum is expected to have characteristic multilocal contributions identifiable as multipart on contributions now and possibly relevant for the understanding of bound states such as hadrons.

5.1 Scale Dependent Notion Of Four-Momentum In Zero Energy Ontology

Quite generally, General Relativity does not allow to identify four-momentum as Noether charges but in GRT based cosmology one can speak of non-conserved mass [K10], which seems to be in conflict with the conservation of four-momentum in TGD framework. The solution of the problem comes in terms of zero energy ontology (ZEO) [K1, K16], which transforms four-momentum to a scale dependent notion: to each causal diamond (CD) one can assign four-momentum assigned with say positive energy part of the quantum state defined as a quantum superposition of 4-surfaces inside CD.

ZEO is necessary also for the fusion of real and various p-adic physics to single coherent whole. ZEO also allows maximal “free will” in quantum jump since every zero energy state can be created from vacuum and at the same time allows consistency with the conservation laws. ZEO has rather dramatic implications: in particular the arrow of thermodynamical time is predicted to vary so that second law must be generalized. This has especially important implications in living matter, where this kind of variation is observed.

More precisely, this superposition corresponds to a spinor field in the “world of classical worlds” (WCW) [K16]: its components - WCW spinors - correspond to elements of fermionic Fock basis for a given 4-surface - or by holography implied by general coordinate invariance (GCI) - for 3-surface having components at both ends of CD. Strong form of GGI implies strong form of holography (SH) so that partonic 2-surfaces at the ends of space-time surface plus their 4-D tangent space data are enough to fix the quantum state. The classical dynamics in the interior is necessary for the translation of the outcomes of quantum measurements to the language of physics based on classical fields, which in turn is reduced to sub-manifold geometry in the extension of the geometrization program of physics provided by TGD.

Holography is very much reminiscent of QCC suggesting trinity: GCI-holography-QCC. Strong form of holography has strongly stringy flavor: string world sheets connecting the wormhole throats appearing as basic building bricks of particles emerge from the dynamics of induced spinor fields if one requires that the fermionic mode carries well-defined electromagnetic charge [K15].

5.2 Are The Classical And Quantal Four-Momenta Identical?

One key question concerns the classical and quantum counterparts of four-momentum. In TGD framework classical theory is an exact part of quantum theory. Classical four-momentum corresponds to Noether charge for preferred extremals of Kähler action. Quantal four-momentum in turn is assigned with the quantum superposition of space-time sheets assigned with CD - actually WCW spinor field analogous to ordinary spinor field carrying fermionic degrees of freedom as analogs of spin. Quantal four-momentum emerges just as it does in super string models - that is as a parameter associated with the representations of super-conformal algebras. The precise action of translations in the representation remains poorly specified. Note that quantal four-momentum does not emerge as Noether charge: at least it is

not at all obvious that this could be the case.

Are these classical and quantal four-momenta identical as QCC would suggest? If so, the Noether four-momentum should be same for all space-time surfaces in the superposition. QCC suggests that also the classical correlation functions for various general coordinate invariant local quantities are same as corresponding quantal correlation functions and thus same for all 4-surfaces in quantum superposition - this at least in the measurement resolution used. This would be an extremely powerful constraint on the quantum states and to a high extend could determined the U-, M-, and S-matrices.

QCC seems to be more or less equivalent with SH stating that in some respects the descriptions based on classical physics defined by Kähler action in the interior of space-time surface and the quantal description in terms of quantum states assignable to the intersections of space-like 3-surfaces at the boundaries of CD and light-like 3-surfaces at which the signature of induced metric changes. SH means effective 2-dimensionality since the four-dimensional tangent space data at partonic 2-surfaces matters. SH could be interpreted as Kac-Mody and symplectic symmetries meaning that apart from central extension they act almost like gauge symmetries in the interiors of space-like 3-surfaces at the ends of CD and in the interiors of light-like 3-surfaces representing orbits of partonic 2-surfaces. Gauge conditions are replaced with Super Virasoro conditions. The word “almost” is of course extremely important.

5.3 What Equivalence Principle (EP) Means In Quantum TGD?

EP states the equivalence of gravitational and inertial masses in Newtonian theory. A possible generalization would be equivalence of gravitational and inertial four-momenta. In GRT this correspondence cannot be realized in mathematically rigorous manner since these notions are poorly defined and EP reduces to a purely local statement in terms of Einstein’s equations.

What about TGD? What could EP mean in TGD framework?

1. Is EP realized at both quantum and space-time level? This option requires the identification of inertial and gravitational four-momenta at both quantum and classical level. It is now clear that at classical level EP follows from very simple assumption that GRT space-time is obtained by lumping together the space-time sheets of the many-sheeted space-time and by the identification the effective metric as sum of M^4 metric and deviations of the induced metrics of space-time sheets from M^2 metric: the deviations indeed define the gravitational field defined by multiply topologically condensed test particle. Similar description applies to gauge fields. EP as expressed by Einstein’s equations would follow from Poincare invariance at microscopic level defined by TGD space-time. The effective fields have as sources the energy momentum tensor and YM currents defined by topological inhomogenities smaller than the resolution scale.
2. QCC would require the identification of quantal and classical counterparts of both gravitational and inertial four-momenta. This would give three indepen-

dent equivalences, say $P_{I,class} = P_{I,quant}$, $P_{gr,class} = P_{gr,quant}$, $P_{gr,class} = P_{I,quant}$, which imply the remaining ones.

Consider the condition $P_{gr,class} = P_{I,class}$. At classical level the condition that the standard energy momentum tensor associated with Kähler action has a vanishing divergence is guaranteed if Einstein's equations with cosmological term are satisfied. If preferred extremals satisfy this condition they are constant curvature spaces for non-vanishing cosmological constant. A more general solution ansatz involves several functions analogous to cosmological constant corresponding to the decomposition of energy momentum tensor to terms proportional to Einstein tensor and several lower-dimensional projection operators [K18]. It must be emphasized that field equations are extremely non-linear and one must also consider preferred extremals (which could be identified in terms of space-time regions having so called Hamilton-Jacobi structure): hence these proposals are guesses motivated by what is known about exact solutions of field equations.

Consider next $P_{gr,class} = P_{I,class}$. At quantum level I have proposed coset representations for the pair of super conformal algebras g and $h \subset g$ which correspond to the coset space decomposition of a given sector of WCW with constant values of zero modes. The coset construction would state that the differences of super-Virasoro generators associated with g resp. h annihilate physical states.

The identification of the algebras g and h is not straightforward. The algebra g could be formed by the direct sum of super-symplectic and super Kac-Moody algebras and its sub-algebra h for which the generators vanish at partonic 2-surface considered. This would correspond to the idea about WCW as a coset space G/H of corresponding groups (consider as a model $CP_2 = SU(3)/U(2)$ with $U(2)$ leaving preferred point invariant). The sub-algebra h in question includes or equals to the algebra of Kac-Moody generators vanishing at the partonic 2-surface. A natural choice for the preferred WCW point would be as maximum of Kähler function in Euclidian regions: positive definiteness of Kähler function allows only single maximum for fixed values of zero modes). Coset construction states that differences of super Virasoro generators associated with g and h annihilate physical states. This implies that corresponding four-momenta are identical that is Equivalence Principle.

3. Does EP at quantum level reduce to one aspect of QCC? This would require that classical Noether four-momentum identified as inertial momentum equals to the quantal four-momentum assignable to the states of super-conformal representations and identifiable as gravitational four-momentum. There would be only one independent condition: $P_{class} \equiv P_{I,class} = P_{gr,quant} \equiv P_{quant}$.

Holography realized as AdS/CFT correspondence states the equivalence of descriptions in terms of gravitation realized in terms of strings in 10-D space-time and gauge fields at the boundary of AdS. What is disturbing is that this picture is not completely equivalent with the proposed one. In this case the super-conformal algebra would be direct sum of super-symplectic and super Kac-Moody parts.

Which of the options looks more plausible? The success of p-adic mass calculations [K17] have motivated the use of them as a guideline in attempts to understand TGD. The basic outcome was that elementary particle spectrum can be understood if Super Virasoro algebra has five tensor factors. Can one decide the fate of the two approaches to EP using this number as an input?

This is not the case. For both options the number of tensor factors is five as required. Four tensor factors come from Super Kac-Moody and correspond to translational Kac-Moody type degrees of freedom in M^4 , to color degrees of freedom and to electroweak degrees of freedom ($SU(2) \times U(1)$). One tensor factor comes from the symplectic degrees of freedom in $\Delta CD \times CP_2$ (note that Hamiltonians include also products of δCD and CP_2 Hamiltonians so that one does not have direct sum!).

The reduction of EP to the coset structure of WCW sectors is extremely beautiful property. But also the reduction of EP to QCC looks very nice and deep. It is of course possible that the two realizations of EP are equivalent and the natural conjecture is that this is the case.

For QCC option the GRT inspired interpretation of Equivalence Principle at space-time level remains to be understood. Is it needed at all? The condition that the energy momentum tensor of Kähler action has a vanishing divergence leads in General Relativity to Einstein equations with cosmological term. In TGD framework preferred extremals satisfying the analogs of Einstein's equations with several cosmological constant like parameters can be considered.

Should one give up this idea, which indeed might be wrong? Could the divergence of energy momentum tensor vanish only asymptotically as was the original proposal? Or should one try to generalize the interpretation? QCC states that quantum physics has classical correlate at space-time level and implies EP. Could also quantum classical correspondence itself have a correlate at space-time level. If so, space-time surface would be able to represent abstractions as statements about statements about.... as the many-sheeted structure and the vision about TGD physics as analog of Turing machine able to mimic any other Turing machine suggest.

5.4 TGD-GRT Correspondence And Equivalence Principle

One should also understand how General Relativity and EP emerge at classical level. The understanding comes from the realization that GRT is only an effective theory obtained by endowing M^4 with effective metric.

1. The replacement of superposition of fields with superposition of their effects means replacing superposition of fields with the set-theoretic union of space-time surfaces. Particle experiences sum of the effects caused by the classical fields at the space-time sheets (see **Fig.** <http://tgdtheory.fi/appfigures/fieldsuperpose.jpg> or **Fig. ??** in the appendix of this book).
2. This is true also for the classical gravitational field defined by the deviation from flat Minkowski metric in standard M^4 coordinates for the space-time sheets. One can define effective metric as sum of M^4 metric and deviations. This effective metric would correspond to that of General Relativity. This resolves long standing issues relating to the interpretation of TGD.

3. Einstein's equations could hold true for the effective metric. They are motivated by the underlying Poincare invariance which cannot be realized as global conservation laws for the effective metric. The conjecture vanishing of divergence of Kähler energy momentum tensor can be seen as the microscopic justification for the claim that Einstein's equations hold true for the effective space-time.
4. The breaking of Poincare invariance could have interpretation as effective breaking in zero energy ontology (ZEO), in which various conserved charges are length dependent and defined separately for each causal diamond (CD).

One can of course consider the possibility that Einstein's equations generalize for preferred extremals of Kähler action. This would actually represent at space-time level the notion of QCC rather than realise QCC interpreted as EP. The condition that the energy momentum tensor for Kähler action has vanishing covariant divergence would be satisfied in GRT if Einstein's equations with cosmological term hold true. This is the case also now but one can consider also more general solutions in which one has two cosmological constants which are not genuine constants anymore [K18].

5.5 How Translations Are Represented At The Level Of WCW ?

The four-momentum components appearing in the formulas of super conformal generators correspond to infinitesimal translations. In TGD framework one must be able to identify these infinitesimal translations precisely. As a matter of fact, finite measurement resolution implies that it is probably too much to assume infinitesimal translations. Rather, finite exponentials of translation generators are involved and translations are discretized. This does not have practical significance since for optimal resolution the discretization step is about CP_2 length scale.

Where and how do these translations act at the level of WCW ? ZEO provides a possible answer to this question.

5.5.1 Discrete Lorentz transformations and time translations act in the space of CDs: inertial four-momentum

Quantum state corresponds also to wave function in moduli space of CDs. The moduli space is obtained from given CD by making all boosts for its non-fixed boundary: boosts correspond to a discrete subgroup of Lorentz group and define a lattice-like structure at the hyperboloid for which proper time distance from the second tip of CD is fixed to $T_n = n \times T(CP_2)$. The quantization of cosmic redshift for which there is evidence, could relate to this lattice generalizing ordinary 3-D lattices from Euclidian to hyperbolic space by replacing translations with boosts (velocities).

The additional degree of freedom comes from the fact that the integer $n > 0$ obtains all positive values. One has wave functions in the moduli space defined as a pile of these lattices defined at the hyperboloid with constant value of $T(CP_2)$:

one can say that the points of this pile of lattices correspond to Lorentz boosts and scalings of CDs defining sub- WCW : s.

The interpretation in terms of group which is product of the group of shifts $T_n(CP_2) \rightarrow T_{n+m}(CP_2)$ and discrete Lorentz boosts is natural. This group has same Cartesian product structure as Galilean group of Newtonian mechanics. This would give a discrete rest energy and by Lorentz boosts discrete set of four-momenta giving a contribution to the four-momentum appearing in the super-conformal representation.

What is important that each state function reduction would mean localisation of either boundary of CD (that is its tip). This localization is analogous to the localization of particle in position measurement in E^3 but now discrete Lorentz boosts and discrete translations $T_n \rightarrow T_{n+m}$ replace translations. Since the second end of CD is necessary del-ocalized in moduli space, one has kind of flip-flop: localization at second end implies de-localization at the second end. Could the localization of the second end (tip) of CD in moduli space correspond to our experience that momentum and position can be measured simultaneously? This apparent classicality would be an illusion made possible by ZEO.

The flip-flop character of state function reduction process implies also the alternation of the direction of the thermodynamical time: the asymmetry between the two ends of CDs would induce the quantum arrow of time. This picture also allows to understand what the experience growth of geometric time means in terms of CDs.

5.5.2 The action of translations at space-time sheets

The action of imbedding space translations on space-time surfaces possibly becoming trivial at partonic 2-surfaces or reducing to action at δCD induces action on space-time sheet which becomes ordinary translation far enough from end end of space-time surface. The four-momentum in question is very naturally that associated with Kähler action and would therefore correspond to inertial momentum for $P_{I,class} = P_{quant,gr}$ option. Indeed, one cannot assign quantal four-momentum to Kähler action as an operator since canonical quantization badly fails. In finite measurement infinitesimal translations are replaced with their exponentials for $P_{I,class} = P_{quant,gr}$ option.

What looks like a problem is that ordinary translations in the general case lead out from given CD near its boundaries. In the interior one expects that the translation acts like ordinary translation. The Lie-algebra structure of Poincare algebra including sums of translation generators with positive coefficient for time translation is preserved if only time-like superpositions if generators are allowed also the commutators of time-like translation generators with boost generators give time like translations. This defines a Lie-algebraic formulation for the arrow of geometric time. The action of time translation on preferred extremal would be ordinary translation plus continuation of the translated preferred extremal backwards in time to the boundary of CD. The transversal space-like translations could be made Kac-Moody algebra by multiplying them with functions which vanish at δCD .

A possible interpretation would be that $P_{quant,gr}$ corresponds to the momentum assignable to the moduli degrees of freedom and $P_{cl,I}$ to that assignable to the time like translations. $P_{quant,gr} = P_{cl,I}$ would code for QCC. Geometrically quantum

classical correspondence would state that time-like translation shift both the interior of space-time surface and second boundary of CD to the geometric future/past while keeping the second boundary of space-time surface and CD fixed.

5.6 Yangian And Four-Momentum

Yangian symmetry implies the marvellous results of twistor Grassmannian approach to $\mathcal{N} = 4$ SUSY culminating in the notion of amplituhedron which promises to give a nice projective geometry interpretation for the scattering amplitudes [?]. Yangian symmetry is a multilocal generalization of ordinary symmetry based on the notion of co-product and implies that Lie algebra generators receive also multilocal contributions. I have discussed these topics from slightly different point of view in [K12], where also references to the work of pioneers can be found.

5.6.1 Yangian symmetry

The notion equivalent to that of Yangian was originally introduced by Faddeev and his group in the study of integrable systems. Yangians are Hopf algebras which can be assigned with Lie algebras as the deformations of their universal enveloping algebras. The elegant but rather cryptic looking definition is in terms of the modification of the relations for generating elements [K12]. Besides ordinary product in the enveloping algebra there is co-product Δ which maps the elements of the enveloping algebra to its tensor product with itself. One can visualize product and co-product in terms of particle reactions. Particle annihilation is analogous to annihilation of two particles into a single one and co-product is analogous to the decay of a particle into two. Δ allows to construct higher generators of the algebra.

Lie-algebra can mean here ordinary finite-dimensional simple Lie algebra, Kac-Moody algebra or Virasoro algebra. In the case of SUSY it means conformal algebra of M^4 - or rather its super counterpart. Witten, Nappi and Dolan have described the notion of Yangian for super-conformal algebra in very elegant and concrete manner in the article *Yangian Symmetry in $D=4$ superconformal Yang-Mills theory* [?]. Also Yangians for gauge groups are discussed.

In the general case Yangian resembles Kac-Moody algebra with discrete index n replaced with a continuous one. Discrete index poses conditions on the Lie group and its representation (adjoint representation in the case of $\mathcal{N} = 4$ SUSY). One of the conditions is that the tensor product $R \otimes R^*$ for representations involved contains adjoint representation only once. This condition is non-trivial. For $SU(n)$ these conditions are satisfied for any representation. In the case of $SU(2)$ the basic branching rule for the tensor product of representations implies that the condition is satisfied for the product of any representations.

Yangian algebra with a discrete basis is in many respects analogous to Kac-Moody algebra. Now however the generators are labelled by non-negative integers labeling the light-like incoming and outgoing momenta of scattering amplitude whereas in the case of Kac-Moody algebra also negative values are allowed. Note that only the generators with non-negative conformal weight appear in the construction of states of Kac-Moody and Virasoro representations so that the extension to Yangian makes sense.

The generating elements are labelled by the generators of ordinary conformal transformations acting in M^4 and their duals acting in momentum space. These two sets of elements can be labelled by conformal weights $n = 0$ and $n = 1$ and their mutual commutation relations are same as for Kac-Moody algebra. The commutators of $n = 1$ generators with themselves are however something different for a non-vanishing deformation parameter h . Serre's relations characterize the difference and involve the deformation parameter h . Under repeated commutations the generating elements generate infinite-dimensional symmetric algebra, the Yangian. For $h = 0$ one obtains just one half of the Virasoro algebra or Kac-Moody algebra. The generators with $n > 0$ are $n + 1$ -local in the sense that they involve $n + 1$ -forms of local generators assignable to the ordered set of incoming particles of the scattering amplitude. This non-locality generalizes the notion of local symmetry and is claimed to be powerful enough to fix the scattering amplitudes completely.

5.6.2 How to generalize Yangian symmetry in TGD framework?

As far as concrete calculations are considered, it is not much to say. It is however possible to keep discussion at general level and still say something interesting (as I hope!). The key question is whether it could be possible to generalize the proposed Yangian symmetry and geometric picture behind it to TGD framework.

1. The first thing to notice is that the Yangian symmetry of $\mathcal{N} = 4$ SUSY in question is quite too limited since it allows only single representation of the gauge group and requires massless particles. One must allow all representations and massive particles so that the representation of symmetry algebra must involve states with different masses, in principle arbitrary spin and arbitrary internal quantum numbers. The candidates are obvious: Kac-Moody algebras [A2] and Virasoro algebras [A3] and their super counterparts. Yangians indeed exist for arbitrary super Lie algebras. In TGD framework conformal algebra of Minkowski space reduces to Poincare algebra and its extension to Kac-Moody allows to have also massive states.
2. The formal generalization looks surprisingly straightforward at the formal level. In zero energy ontology one replaces point like particles with partonic two-surfaces appearing at the ends of light-like orbits of wormhole throats located to the future and past light-like boundaries of causal diamond ($CD \times CP_2$ or briefly CD). Here CD is defined as the intersection of future and past directed light-cones. The polygon with light-like momenta is naturally replaced with a polygon with more general momenta in zero energy ontology and having partonic surfaces as its vertices. Non-point-likeness forces to replace the finite-dimensional super Lie-algebra with infinite-dimensional Kac-Moody algebras and corresponding super-Virasoro algebras assignable to partonic 2-surfaces.
3. This description replaces disjoint holomorphic surfaces in twistor space with partonic 2-surfaces at the boundaries of $CD \times CP_2$ so that there seems to be a close analogy with Cachazo-Svrcek-Witten picture. These surfaces are connected by either light-like orbits of partonic 2-surface or space-like 3-surfaces at the ends of CD so that one indeed obtains the analog of polygon.

What does this then mean concretely (if this word can be used in this kind of context)?

1. At least it means that ordinary Super Kac-Moody and Super Virasoro algebras associated with isometries of $M^4 \times CP_2$ annihilating the scattering amplitudes must be extended to a co-algebras with a non-trivial deformation parameter. Kac-Moody group is thus the product of Poincare and color groups. This algebra acts as deformations of the light-like 3-surfaces representing the light-like orbits of particles which are extremals of Chern-Simon action with the constraint that weak form of electric-magnetic duality holds true. I know so little about the mathematical side that I cannot tell whether the condition that the product of the representations of Super-Kac-Moody and Super-Virasoro algebras contains adjoint representation only once, holds true in this case. In any case, it would allow all representations of finite-dimensional Lie group in vertices whereas $\mathcal{N} = 4$ SUSY would allow only the adjoint.
2. Besides this ordinary kind of Kac-Moody algebra there is the analog of Super-Kac-Moody algebra associated with the light-cone boundary which is metrically 3-dimensional. The finite-dimensional Lie group is in this case replaced with infinite-dimensional group of symplectomorphisms of $\delta M^4_{+/-}$ made local with respect to the internal coordinates of the partonic 2-surface. This picture also justifies p-adic thermodynamics applied to either symplectic or isometry Super-Virasoro and giving thermal contribution to the vacuum conformal and thus to mass squared.
3. The construction of TGD leads also to other super-conformal algebras and the natural guess is that the Yangians of all these algebras annihilate the scattering amplitudes.
4. Obviously, already the starting point symmetries look formidable but they still act on single partonic surface only. The discrete Yangian associated with this algebra associated with the closed polygon defined by the incoming momenta and the negatives of the outgoing momenta acts in multi-local manner on scattering amplitudes. It might make sense to speak about polygons defined also by other conserved quantum numbers so that one would have generalized light-like curves in the sense that state are massless in 8-D sense.

5.6.3 Could Yangian symmetry provide a new view about conserved quantum numbers?

The Yangian algebra has some properties which suggest a new kind of description for bound states. The Cartan algebra generators of $n = 0$ and $n = 1$ levels of Yangian algebra commute. Since the co-product Δ maps $n = 0$ generators to $n = 1$ generators and these in turn to generators with high value of n , it seems that they commute also with $n \geq 1$ generators. This applies to four-momentum, color isospin and color hyper charge, and also to the Virasoro generator L_0 acting on Kac-Moody algebra of isometries and defining mass squared operator.

Could one identify total four momentum and Cartan algebra quantum numbers as sum of contributions from various levels? If so, the four momentum and mass

squared would involve besides the local term assignable to wormhole throats also n -local contributions. The interpretation in terms of n -parton bound states would be extremely attractive. n -local contribution would involve interaction energy. For instance, string like object would correspond to $n = 1$ level and give $n = 2$ -local contribution to the momentum. For baryonic valence quarks one would have 3-local contribution corresponding to $n = 2$ level. The Yangian view about quantum numbers could give a rigorous formulation for the idea that massive particles are bound states of massless particles.

6 Generalization Of Ads/CFT Duality To TGD Framework

AdS/CFT duality has provided a powerful approach in the attempts to understand the non-perturbative aspects of super-string theories. The duality states that conformal field theory in n -dimensional Minkowski space M^n identifiable as a boundary of $n + 1$ -dimensional space AdS_{n+1} is dual to a string theory in $AdS_{n+1} \times S^{9-n}$.

As a mathematical discovery the duality is extremely interesting but it seems that it need not have much to do with physics. From TGD point of view the reason is obvious: the notion of conformal invariance is quite too limited. In TGD framework conformal invariance is extended to a super-symplectic symmetry in $\delta M_{\pm}^4 \times CP_2$, whose Lie-algebra has the structure of conformal algebra. Also ordinary super-conformal symmetries associated with string world sheets are present as well as generalization of 2-D conformal symmetries to their analogs at light-cone boundary and light-like orbits of partonic 2-surfaces. In this framework AdS/CFT duality is expected to be modified and this seems to be the case.

The matrix elements of Kähler metric of WCW can be expressed in two manners. As contractions of the derivatives $\partial_K \partial_{\bar{L}} K$ of the Kähler function of WCW with isometry generators or as anticommutators of WCW gamma matrices identified as supersymplectic Noether super charges assignable to fermion strings connecting partonic 2-surfaces. Kähler function is identified as Kähler action for the Euclidian space-time regions with 4-D CP_2 projection. Kähler action defines the Kähler-Dirac gamma matrices appearing in K-D action as contractions of canonical momentum currents with imbedding space gamma matrices. Bosonic and fermionic degrees of freedom are therefore dual in a well-defined sense.

This observation suggests various generalizations. There is super-symmetry between Kähler action and Kähler-Dirac action. The problem is that induced spinor fields are localized at 2-D string world sheets. Strong form of holography implying effective 2-dimensionality suggests the solution to the paradox. The paradox disappears if the Kähler action is expressible as string area for the effective metric defined by the anti-commutators of K-D gamma matrices at string world sheet. This expression allows to understand how strings connecting partonic 2-surfaces give rise to the formation of gravitationally bound states. Bound states of macroscopic size are however possible only if one allows hierarchy of Planck constants. This representation of Kähler action can be seen as one aspect of the analog of AdS/CFT duality in TGD framework.

One can imagine also other realizations. For instance, Dirac determinant for the

spinors associated with string world sheets should reduce to the exponent of Kähler action.

6.1 Does The Exponent Of Chern-Simons Action Reduce To The Exponent Of The Area Of Minimal Surfaces?

As I scanned of hep-th I found an interesting article by Giordano, Peschanski, and Seki [?] based on AdS/CFT correspondence. What is studied is the high energy behavior of the gluon-gluon and quark-quark scattering amplitudes of $\mathcal{N} = 4$ SUSY.

1. The proposal made earlier by Aldaya and Maldacena [?] is that gluon-gluon scattering amplitudes are proportional to the imaginary exponent of the area of a minimal surface in AdS_5 whose boundary is identified as *momentum space*. The boundary of the minimal surface would be polygon with light-like edges: this polygon and its dual are familiar from twistor approach.
2. Giordano, Peschanski, and Seki claim that quark-quark scattering amplitude for heavy quarks corresponds to the exponent of the area for a minimal surface in the *Euclidian* version of AdS_5 which is hyperbolic space (space with a constant negative curvature): it is interpreted as a counterpart of WCW rather than momentum space and amplitudes are obtained by analytic continuation. For instance, a universal Regge behavior is obtained. For general amplitudes the exponent of the area alone is not enough since it does not depend on gluon quantum numbers and vertex operators at the edges of the boundary polygon are needed.

In the following my intention is to consider the formulation of this conjecture in quantum TGD framework. I hasten to inform that I am not a specialist in AdS/CFT and can make only general comments inspired by analogies with TGD and the generalization of AdS/CFT duality to TGD framework based on the localization of induced spinors at string world sheets, super-symmetry between bosonic and fermionic degrees of freedom at the level of WCW, and the notion of effective metric at string world sheets.

6.2 Does Kähler Action Reduce To The Sum Of Areas Of Minimal Surfaces In Effective Metric?

Minimal surface conjectures are highly interesting from TGD point of view. The weak form of electric magnetic duality implies the reduction of Kähler action to 3-D Chern-Simons terms. Effective 2-dimensionality implied by the strong form of General Coordinate Invariance suggests a further reduction of Chern-Simons terms to 2-D terms and the areas of string world sheet and of partonic 2-surface are the only non-topological options that one can imagine. Skeptic could of course argue that the exponent of the minimal surface area results as a characterizer of the quantum state rather than vacuum functional. In the following I end up with the proposal that the Kähler action should reduce to the sum of string world sheet areas in the effective metric defines by the anticommutators of Kähler-Dirac gamma matrices at string world sheets.

Let us look this conjecture in more detail.

1. In zero energy ontology twistor approach is very natural since all physical states are bound states of massless particles. Also virtual particles are composites of massless states. The possibility to have both signs of energy makes possible space-like momenta for wormhole contacts. Mass shell conditions at internal lines imply extremely strong constraints on the virtual momenta and both UV and IR finiteness are expected to hold true.
2. The weak form of electric magnetic duality [K15] implies that the exponent of Kähler action reduces to the exponent of Chern-Simons term for 3-D space-like surfaces at the ends of space-time surface inside CD and for light-like 3-surfaces. The coefficient of this term is complex since the contribution of Minkowskian regions of the space-time surface is imaginary ($\sqrt{g_4}$ is imaginary) and that of Euclidian regions (generalized Feynman diagrams) real. The Chern-Simons term from Minkowskian regions is like Morse function and that from Euclidian regions defines Kähler function and stationary phase approximation makes sense. The two contributions are different since the space-like 3-surfaces contributing to Kähler function and Morse function are different.
3. Electric magnetic duality [K15] leads also to the conclusion that wormhole throats carrying elementary particle quantum numbers are Kähler magnetic monopoles. This forces to identify elementary particles as string like objects with ends having opposite monopole charges. Also more complex configurations are possible.

It is not quite clear what the scale of the stringyness is. The natural first guess inspired by quantum classical correspondence is that it corresponds to the p-adic length scale of the particle characterizing its Compton length. Second possibility is that it corresponds to electroweak scale. For leptons stringyness in Compton length scale might not have any fatal implications since the second end of string contains only neutrinos neutralizing the weak isospin of the state. This kind of monopole pairs could appear even in condensed matter scales: in particular if the proposed hierarchy of Planck constants [K4] is realized.

4. Strong form of General Coordinate Invariance requires effective 2-dimensionality. In given UV and IR resolutions either partonic 2-surfaces or string world sheets form a finite hierarchy of CDs inside CDs with given CD characterized by a discrete scale coming as an integer multiple of a fundamental scale (essentially CP_2 size). The string world sheets have boundaries consisting of either light-like curves in induced metric at light-like wormhole throats and space-like curves at the ends of CD whose M^4 projections are light-like. These braids intersect partonic 2-surfaces at discrete points carrying fermionic quantum numbers.

This implies a rather concrete analogy with $AdS_5 \times S_5$ duality, which describes gluons as open strings. In zero energy ontology (ZEO) string world sheets are indeed a fundamental notion and the natural conjecture is that these surfaces are minimal surfaces whose area by quantum classical correspondence depends

on the quantum numbers of the external particles. String tension in turn should depend on gauge couplings -perhaps only Kähler coupling strength- and geometric parameters like the size scale of CD and the p-adic length scale of the particle.

5. One can of course ask whether the metric defining the string area is induced metric or possibly the metric defined by the anti-commutators of Kähler-Dirac gamma matrices. The recent view does not actually leave any other alternative. The analog of AdS/CFT duality together with supersymmetry demands that Kähler action is proportional to the sum of the areas of string world sheets in this effective metric. Whether the vanishing of induced W fields (and possibly also Z^0) making possible well-defined em charge for the spinor nodes is realized by the condition that the string world sheet is a minimal surface in the effective metric remains an open question.

The assumption that ordinary minimal surfaces are in question is not consistent with the TGD view about the formation of gravitational bound states and if string tension is $1/\hbar G$ as in string models, only bound states with size of order Planck length are possible. This strongly favors effective metric giving string tension proportional to $1/h_{eff}^2$. How $1/h_{eff}^2$ proportionality might be understood is discussed in [K19] in terms electric-magnetic duality.

6. One can of course still consider also the option that ordinary minimal surfaces are in question. Are the minimal surfaces in question minimal surfaces of the imbedding space $M^4 \times CP_2$ or of the space-time surface X^4 ? All possible 2-surfaces at the boundary of CD must be allowed so that they cannot correspond to minimal surfaces in $M^4 \times CP_2$ unless one assumes that they emerge in stationary phase approximation only. The boundary conditions at the ends of CD could however be such that *any* partonic 2-surface correspond to a minimal surfaces in X^4 . Same applies to string world sheets. One might even hope that these conditions combined with the weak form of electric magnetic duality fixes completely the boundary conditions at wormhole throats and space-like ends of space-time surface.

The trace of the second fundamental form orthogonal to the string world sheet/partonic 2-surface as sub-manifold of space-time surface would vanish: this is nothing but a generalization of the geodesic motion obtained by replacing word line with a 2-D surface. It does not imply the vanishing of the trace of the second fundamental form in $M^4 \times CP_2$ having interpretation as a generalization of particle acceleration [K13]. Effective 2-dimensionality would be realized if Chern-Simons terms reduce to a sum of the areas of these minimal surfaces.

These arguments suggest that scattering amplitudes are proportional to the product of exponents of 2-dimensional actions which can be either imaginary or real. Imaginary exponent would be proportional to the total area of string world sheets and the imaginary unit would come naturally from $\sqrt{g_2}$, where g_2 is effective metric most naturally. Real exponent proportional to the total area of partonic 2-surfaces. The coefficient of these areas would not in general be same.

The reduction of the Kähler action from Minkowskian regions to Chern-Simons terms means that Chern-Simons terms reduce to actions assignable to string world sheets. The equality of the Minkowskian and Euclidian Chern-Simons terms is suggestive but not necessarily true since there could be also other Chern-Simons contributions than those assignable to wormhole throats and the ends of space-time. The equality would imply that the total area of string world sheets equals to the total area of partonic 2-surfaces suggesting strongly a duality meaning that either Euclidian or Minkowskian regions carry the needed information.

6.3 Surface Area As Geometric Representation Of Entanglement Entropy?

In Thinking Allowed Original there was a link to a talk by James Sully and having the title Geometry of Compression. I must admit that I understood very little about the talk. My not so educated guess is however that information is compressed: UV or IR cutoff eliminating entanglement in short length scales and describing its presence in terms of density matrix - that is thermodynamically - is another manner to say it. The TGD inspired proposal for the interpretation of the inclusions of hyper-finite factors of type II_1 (HFFs) [K14] is in spirit with this.

The space-time counterpart for the compression would be in TGD framework discretization. Discretizations using rational points (or points in algebraic extensions of rationals) make sense also p-adically and thus satisfy number theoretic universality. Discretization would be defined in terms of intersection (rational or in algebraic extension of rationals) of real and p-adic surfaces. At the level of “world of classical worlds” the discretization would correspond to - say - surfaces defined in terms of polynomials, whose coefficients are rational or in some algebraic extension of rationals. Pinary UV and IR cutoffs are involved too. The notion of p-adic manifold allows to interpret the p-adic variants of space-time surfaces as cognitive representations of real space-time surfaces.

Finite measurement resolution does not allow state function reduction reducing entanglement totally. In TGD framework also negentropic entanglement stable under Negentropy Maximization Principle (NMP) is possible [K5]. For HFFs the projection into single ray of Hilbert space is indeed impossible: the reduction takes always to infinite-D sub-space.

The visit to the URL was however not in vain. There was a link to an article [?] discussing the geometrization of entanglement entropy inspired by the AdS/CFT hypothesis.

Quantum classical correspondence is basic guiding principle of TGD and suggests that entanglement entropy should indeed have space-time correlate, which would be the analog of Hawking-Bekenstein entropy.

6.3.1 Generalization of AdS/CFT to TGD context

AdS/CFT generalizes to TGD context in non-trivial manner. There are two alternative interpretations, which both could make sense. These interpretations are not mutually exclusive. The first interpretation makes sense at the level of “world of

classical worlds" (WCW) with symplectic algebra and extended conformal algebra associated with δM_{\pm}^4 replacing ordinary conformal and Kac-Moody algebras. Second interpretation at the level of space-time surface with the extended conformal algebras of the light-likes orbits of partonic 2-surfaces replacing the conformal algebra of boundary of AdS^n .

1. *First interpretation*

For the first interpretation 2-D conformal invariance is generalised to 4-D conformal invariance relying crucially on the 4-dimensionality of space-time surfaces and Minkowski space.

1. One has an extension of the conformal invariance provided by the symplectic transformations of $\delta CD \times CP_2$ for which Lie algebra has the structure of conformal algebra with radial light-like coordinate of δM_+^4 replacing complex coordinate z .
2. One could see the counterpart of AdS_n as imbedding space $H = M^4 \times CP_2$ completely unique by twistorial considerations and from the condition that standard model symmetries are obtained and its causal diamonds defined as sub-sets $CD \times CP_2$, where CD is an intersection of future and past directed light-cones. I will use the shorthand CD for $CD \times CP_2$. Strings in $AdS_5 \times S^5$ are replaced with space-time surfaces inside 8-D CD.
3. For this interpretation 8-D CD replaces the 10-D space-time $AdS_5 \times S^5$. 7-D light-like boundaries of CD correspond to the boundary of say AdS_5 , which is 4-D Minkowski space so that zero energy ontology (ZEO) allows rather natural formulation of the generalization of AdS/CFT correspondence since the positive and negative energy parts of zero energy states are localized at the boundaries of CD.

6.3.2 Second interpretation

For the second interpretation relies on the observation that string world sheets as carriers of induced spinor fields emerge in TGD framework from the condition that electromagnetic charge is well-defined for the modes of induced spinor field.

1. One could see the 4-D space-time surfaces X^4 as counterparts of AdS_4 . The boundary of AdS_4 is replaced in this picture with 3-surfaces at the ends of space-time surface at opposite boundaries of CD and by strong form of holography the union of partonic 2-surfaces defining the intersections of the 3-D boundaries between Euclidian and Minkowskian regions of space-time surface with the boundaries of CD. Strong form of holography in TGD is very much like ordinary holography.
2. Note that one has a dimensional hierarchy: the ends of the boundaries of string world sheets at boundaries of CD as point-like partices, boundaries as fermion number carrying lines, string world sheets, light-like orbits of partonic 2-surfaces, 4-surfaces, imbedding space $M^4 \times CP_2$. Clearly the situation is more complex than for AdS/CFT correspondence.

3. One can restrict the consideration to 3-D sub-manifolds X^3 at either boundary of causal diamond (CD): the ends of space-time surface. In fact, the position of the other boundary is not well-defined since one has superposition of CDs with only one boundary fixed to be piece of light-cone boundary. The delocalization of the other boundary is essential for the understanding of the arrow of time. The state function reductions at fixed boundary leave positive energy part (say) of the zero energy state at that boundary invariant (in positive energy ontology entire state would remain unchanged) but affect the states associated with opposite boundaries forming a superposition which also changes between reduction: this is analog for unitary time evolution. The average for the distance between tips of CDs in the superposition increases and gives rise to the flow of time.
4. One wants an expression for the entanglement entropy between X^3 and its partner. Bekenstein area law allows to guess the general expression for the entanglement entropy: for the proposal discussed in the article the entropy would be the area of the boundary of X^3 divided by gravitational constant: $S = A/4G$. In TGD framework gravitational constant might be replaced by the square of CP_2 radius apart from numerical constant. How gravitational constant emerges in TGD framework is not completely understood although one can deduce for it an estimate using dimensional analyses. In any case, gravitational constant is a parameter which characterizes GRT limit of TGD in which many-sheeted space-time is in long scales replaced with a piece of Minkowski space such that the classical gravitational fields and gauge potentials for sheets are summed. The physics behind this relies on the generalization of linear superposition of fields: the effects of different space-time sheets particle touching them sum up rather than fields.
5. The counterpart for the boundary of X^3 appearing in the proposal for the geometrization of the entanglement entropy naturally corresponds to partonic 2-surface or a collection of them if strong form of holography holds true.

There is however also another candidate to be considered! Partonic 2-surfaces are basic objects, and one expects that the entanglement between fundamental fermions associated with distinct partonic 2-surfaces has string world sheets as space-time correlates. Could the area of the string world sheet in the effective metric defined by the anti-commutators of K-D gamma matrices at string world sheet provide a measure for entanglement entropy? If this conjecture is correct: the entanglement entropy would be proportional to Kähler action. Also negative values are possible for Kähler action in Minkowskian regions but in TGD framework number theoretic entanglement entropy having also negative values emerges naturally.

Which of these guesses is correct, if any? Or are they equivalent?

6.3.3 With what kind of systems 3-surfaces can entangle?

With what system X^3 is entangled/can entangle? There are several options to consider and they could correspond to the two TGD variants for the AdS/CFT

correspondence.

1. X^3 could correspond to a wormhole contact with Euclidian signature of induced metric. The entanglement would be between it and the exterior region with Minkowskian signature of the induced metric.
2. X^3 could correspond to single sheet of space-time surface connected by wormhole contacts to a larger space-time sheet defining its environment. More precisely, X^3 and its complement would be obtained by throwing away the wormhole contacts with Euclidian signature of induce metric. Entanglement would be between these regions. In the generalization of the formula

$$S = \frac{A}{4\hbar G}$$

area A would be replaced by the total area of partonic 2-surfaces and G perhaps with CP_2 length scale squared.

3. In ZEO the entanglement could also correspond to time-like entanglement between the 3-D ends of the space-time surface at opposite light-like boundaries of CD. M-matrix, which can be seen as the analog of thermal S-matrix, decomposes to a product of hermitian square root of density matrix and unitary S-matrix and this hermitian matrix could also define p-adic thermodynamics. Note that in ZEO quantum theory can be regarded as square root of thermodynamics.

6.3.4 Minimal surface property is not favored in TGD framework

Minimal surface property for the 3-surfaces X^3 at the ends of space-time surface looks at first glance strange but a proper generalization of this condition makes sense if one assumes strong form of holography. Strong form of holography realizes General Coordinate Invariance (GCI) in strong sense meaning that light-like parton orbits and space-like 3-surfaces at the ends of space-time surfaces are equivalent physically. As a consequence, partonic 2-surfaces and their 4-D tangent space data must code for the quantum dynamics.

The mathematical realization is in terms of conformal symmetries accompanying the symplectic symmetries of $\delta M_{\pm}^4 \times CP_2$ and conformal transformations of the light-like partonic orbit [K15]. The generalizations of ordinary conformal algebras correspond to conformal algebra, Kac-Moody algebra at the light-like parton orbits and to symplectic transformations $\delta M^4 \times CP_2$ acting as isometries of WCW and having conformal structure with respect to the light-like radial coordinate plus conformal transformations of δM_+^4 , which is metrically 2-dimensional and allows extended conformal symmetries.

1. If the conformal realization of the strong form of holography works, conformal transformations act at quantum level as gauge symmetries in the sense that generators with no-vanishing conformal weight are zero or generate zero norm states. Conformal degeneracy can be eliminated by fixing the gauge somehow. Classical conformal gauge conditions analogous to Virasoro and Kac-Moody

conditions satisfied by the 3-surfaces at the ends of CD are natural in this respect. Similar conditions would hold true for the light-like partonic orbits at which the signature of the induced metric changes.

2. What is also completely new is the hierarchy of conformal symmetry breakings associated with the hierarchy of Planck constants $h_{eff}/h = n$ [K4]. The deformations of the 3-surfaces which correspond to non-vanishing conformal weight in algebra or any sub-algebra with conformal weights vanishing modulo n give rise to vanishing classical charges and thus do not affect the value of the Kähler action [K15].

The inclusion hierarchies of conformal sub-algebras are assumed to correspond to those for hyper-finite factors. There is obviously a precise analogy with quantal conformal invariance conditions for Virasoro algebra and Kac-Moody algebra. There is also hierarchy of inclusions which corresponds to hierarchy of measurement resolutions. An attractive interpretation is that singular conformal transformations relate to each other the states for broken conformal symmetry. Infinitesimal transformations for symmetry broken phase would carry fractional conformal weights coming as multiples of $1/n$.

3. Conformal gauge conditions need not reduce to minimal surface conditions holding true for all variations.
4. Note that Kähler action reduces to Chern-Simons term at the ends of CD if weak form of electric magnetic duality holds true. The conformal charges at the ends of CD cannot however reduce to Chern-Simons charges by this condition since only the charges associated with CP_2 degrees of freedom would be non-trivial.

The way out of the problem is provided by the generalization of AdS/CFT conjecture. String area is estimated in the effective metric provided by the anti-commutator of K-D gamma matrices at string world sheet.

6.4 Related Ideas

p-Adic mass calculations led to the introduction of the p-adic variant of Bekenstein-Hawking law in which Planck length is replaced by p-adic length scale. This generalization is in spirit with the idea that string world sheet area is estimated in effective rather than induced metric.

6.4.1 p-Adic variant of Bekenstein-Hawking law

When the 3-surface corresponds to elementary particle, a direct connection with p-adic thermodynamics suggests itself and allows to answer the questions above. p-Adic thermodynamics could be interpreted as a description of the entanglement with environment. In ZEO the entanglement could also correspond to time-like entanglement between the 3-D ends of the space-time surface at opposite light-like boundaries of CD. M-matrix, which can be seen as the analog of thermal S-matrix, decomposes to a product of hermitian square root of density matrix and unitary S-matrix and this hermitian matrix could also define p-adic thermodynamics.

1. p-Adic thermodynamics [K17] would not be for energy but for mass squared (or scaling generator L_0) would describe the entanglement of the particle with environment defined by the larger space-time sheet. Conformal weights would come as positive powers of integers (p_0^L would replace $\exp(-H/T)$ to guarantee the number theoretical existence and convergence of the Boltzmann weight: note that conformal invariance that is integer spectrum of L_0 is also essential).
2. The interactions with environment would excite very massive CP_2 mass scale excitations (mass scale is about 10^{-4} times Planck mass) of the particle and give it thermal mass squared identifiable as the observed mass squared. The Boltzmann weights would be extremely small having p-adic norm about $1/p^n$, p the p-adic prime: $M_{127} = 2^{127} - 1$ for electron.
3. One of the first ideas inspired by p-adic vision was that p-adic entropy could be seen as a p-adic counterpart of Bekenstein-Hawking entropy [K7]. $S = (R^2/\hbar^2) \times M^2$ holds true identically apart from numerical constant. Note that one could interpret $R^2 M/\hbar$ as the counterpart of Schwarzschild radius. Note that this radius is proportional to $1/\sqrt{p}$ so that the area A would correspond to the area defined by Compton length. This is in accordance with the third option.

6.4.2 What is the space-time correlate for negentropic entanglement?

The new element brought in by TGD framework is that number theoretic entanglement entropy is negative for negentropic entanglement assignable to unitary entanglement (in the sense that entanglement matrix is proportional to a unitary matrix) and NMP states that this negentropy increases [K5]. Since entropy is essentially number of energy degenerate states, a good guess is that the number $n = h_{eff}/\hbar$ of space-time sheets associated with h_{eff} defines the negentropy. An attractive space-time correlate for the negentropic entanglement is braiding. Braiding defines unitary S-matrix between the states at the ends of braid and this entanglement is negentropic. This entanglement gives also rise to topological quantum computation.

6.5 The Importance Of Being Light-Like

The singular geometric objects associated with the space-time surface have become increasingly important in TGD framework. In particular, the recent progress has made clear that these objects might be crucial for the understanding of quantum TGD. The singular objects are associated not only with the induced metric but also with the effective metric defined by the anti-commutators of the Kähler-Dirac gamma matrices appearing in the Kähler-Dirac equation and determined by the Kähler action.

6.5.1 The singular objects associated with the induced metric

Consider first the singular objects associated with the induced metric.

1. At light-like 3-surfaces defined by wormhole throats the signature of the induced metric changes from Euclidian to Minkowskian so that 4-metric is degenerate. These surfaces are carriers of elementary particle quantum numbers and the 4-D induced metric degenerates locally to 3-D one at these surfaces.
2. Braid strands at partonic orbits - fermion lines identified as boundaries of string world sheets in the more recent terminology - are most naturally light-like curves: this correspond to the boundary condition for open strings. One can assign fermion number to the braid strands. Braid strands allow an identification as curves along which the Euclidian signature of the string world sheet in Euclidian region transforms to Minkowskian one. Number theoretic interpretation would be as a transformation of complex regions to hyper-complex regions meaning that imaginary unit i satisfying $i^2 = -1$ becomes hyper-complex unit e satisfying $e^2 = 1$. The complex coordinates (z, \bar{z}) become hyper-complex coordinates $(u = t + ex, v = t - ex)$ giving the standard light-like coordinates when one puts $e = 1$.

6.5.2 The singular objects associated with the effective metric

There are also singular objects assignable to the effective metric. According to the simple arguments already developed, string world sheets and possibly also partonic 2-surfaces are singular objects with respect to the effective metric defined by the anti-commutators of the Kähler-Dirac gamma matrices rather than induced gamma matrices. Therefore the effective metric might be more than a mere formal structure. The following is of course mere speculation and should be taken as such.

1. For instance, quaternionicity of the space-time surface *might* allow an elegant formulation in terms of the effective metric avoiding the problems due to the Minkowski signature. This is achieved if the effective metric has Euclidian signature $\epsilon \times (1, 1, 1, 1)$, $\epsilon = \pm 1$ or a complex counterpart of the Minkowskian signature $\epsilon(1, 1, -1, -1)$.
2. String world sheets and perhaps also partonic 2-surfaces might be understood as singularities of the effective metric. What happens that the effective metric with Euclidian signature $\epsilon \times (1, 1, 1, 1)$ transforms to the signature $\epsilon(1, 1, -1, -1)$ (say) at string world sheet so that one would have the degenerate signature $\epsilon \times (1, 1, 0, 0)$ at the string world sheet.

What is amazing is that this works also number theoretically. It came as a total surprise to me that the notion of hyper-quaternions as a closed algebraic structure indeed exists. The hyper-quaternionic units would be given by $(1, I, iJ, iK)$, where i is a commuting imaginary unit satisfying $i^2 = -1$. Hyper-quaternionic numbers defined as combinations of these units with real coefficients do form a closed algebraic structure which however fails to be a number field just like hyper-complex numbers do. Note that the hyper-quaternions obtained with real coefficients from the basis $(1, iI, iJ, iK)$ fail to form an algebra since the product is not hyper-quaternion in this sense but belongs to the algebra of complexified quaternions. The same problem is encountered in the case of hyper-octonions defined in this manner. This has

been a stone in my shoe since I feel strong disrelish towards Wick rotation as a trick for moving between different signatures.

3. Could also partonic 2-surfaces correspond to this kind of singular 2-surfaces? In principle, 2-D surfaces of 4-D space intersect at discrete points just as string world sheets and partonic 2-surfaces do so that this might make sense. By complex structure the situation is algebraically equivalent to the analog of plane with non-flat metric allowing all possible signatures (ϵ_1, ϵ_2) in various regions. At light-like curve either ϵ_1 or ϵ_2 changes sign and light-like curves for these two kinds of changes can intersect as one can easily verify by drawing what happens. At the intersection point the metric is completely degenerate and simply vanishes.
4. Replacing real 2-dimensionality with complex 2-dimensionality, one obtains by the universality of algebraic dimension the same result for partonic 2-surfaces and string world sheets. The braid ends at partonic 2-surfaces representing the intersection points of 2-surfaces of this kind would have completely degenerate effective metric so that the Kähler-Dirac gamma matrices would vanish implying that energy momentum tensor vanishes as does also the induced Kähler field.
5. The effective metric suffers a local conformal scaling in the critical deformations identified in the proposed manner. Since ordinary conformal group acts on Minkowski space and leaves the boundary of light-cone invariant, one has two conformal groups. It is not however clear whether the M^4 conformal transformations can act as symmetries in TGD, where the presence of the induced metric in Kähler action breaks M^4 conformal symmetry. As found, also in TGD framework the Kac-Moody currents assigned to the braid strands generate Yangian: this is expected to be true also for the Kac-Moody counterparts of the conformal algebra associated with quantum criticality. On the other hand, in twistor program one encounters also two conformal groups and the space in which the second conformal group acts remains somewhat mysterious object. The Lie algebras for the two conformal groups generate the conformal Yangian and the integrands of the scattering amplitudes are Yangian invariants. Twistor approach should apply in TGD if zero energy ontology is right. Does this mean a deep connection?

What is also intriguing that twistor approach in principle works in strict mathematical sense only at signatures $\epsilon \times (1, 1, -1-1)$ and the scattering amplitudes in Minkowski signature are obtained by analytic continuation. Could the effective metric give rise to the desired signature? Note that the notion of massless particle does not make sense in the signature $\epsilon \times (1, 1, 1, 1)$.

These arguments provide genuine a support for the notion of quaternionicity and suggest a connection with the twistor approach.

7 Could One Define Dynamical Homotopy Groups In WCW?

Agostino Prastaro - working as professor at the University of Rome - has done highly interesting work with partial differential equations, also those assignable to geometric variational principles such as Kähler action in TGD [A6, A7]. I do not understand the mathematical details but the key idea is a simple and elegant generalization of Thom's cobordism theory, and it is difficult to avoid the idea that the application of Prastaro's idea might provide insights about the preferred extremals, whose identification is now on rather firm basis [K22].

One could also consider a definition of what one might call dynamical homotopy groups as a genuine characteristics of WCW topology. The first prediction is that the values of conserved classical Noether charges correspond to disjoint components of WCW. Could this mean that the natural topology in the parameter space of Noether charges zero modes of WCW metric) is p-adic? An analogous conjecture was made on basis of spin glass analogy long time ago. Second surprise is that the only the six lowest dynamical homotopy groups of WCW would be non-trivial. The finite number of these groups dictate by the dimension of imbedding space suggests also an interpretation as analogs of homology groups.

In the following the notion of cobordism is briefly discussed and the idea of Prastaro about assigning cobordism with partial differential equations is discussed.

7.1 About Cobordism As A Concept

To get some background consider first the notion of cobordism (<https://en.wikipedia.org/wiki/Cobordism>).

1. Thom's cobordism theory [A11] is inspired by the question "When an n -manifold can be represented as a boundary of $n + 1$ -manifold". One can also pose additional conditions such as continuity, smoothness, orientability, one can add bundles structures and require that they are induced to n -manifold from that of $n + 1$ -manifold. One can also consider sub-manifolds of some higher-dimensional manifold.

One can also fix n -manifold M and ask "What is the set of n -manifolds N with the property that there exists $n + 1$ -manifold W having union of $M \cup N$ as its boundary". One can also allow M to have boundary and pose the same question by allowing also the boundary of connecting $n+1$ -manifold W contain also the orbits of boundaries of M and N .

The cobordism class of M can be defined as the set of manifolds N cobordant with M - that is connectable in this manner. They have same cobordism class since cobordism is equivalence relation. The classes form also a group with respect to disjoint union. Cobordism is much rougher equivalence relation than diffeomorphy or homeomorphy since topology changes are possible. For instance, every 3-D closed un-oriented manifold is a boundary of a 4-manifold! Same is true for orientable cobordisms. Cobordism defines a category: objects are (say closed) manifolds and morphisms are cobordisms.

2. The basic result of Morse, Thom, and Milnor is that cobordism as topology changes can be engineered from elementary cobordisms. One take manifold $M \times I$ and imbeds to its other n -dimensional end the manifold $S^p \times D^q$, $n = p + q$, removes its interior and glues back $D^{p+1} \times S^{q-1}$ along its boundary to the boundary of the resulting hole. This gives n -manifold with different topology, call it N . The outcome is a cobordism connecting M and N unless there are some obstructions.

There is a connection with Morse theory (https://en.wikipedia.org/wiki/Morse_theory) in which cobordism can be seen as a mapping of W to a unit interval such that the inverse images define a slicing of W and the inverse images at ends correspond to M and N .

3. One can generalize the abstract cobordism to that for n -sub-manifolds of a given imbedding space. This generalization is natural in TGD framework. This might give less trivial results since not all connecting manifolds are imbeddable into a given imbedding space. If connecting 4-manifolds connecting 3-manifolds with Euclidian signature (of induced metric) are assumed to have a Minkowskian signature, one obtains additional conditions, which might be too strong (the classical result of Geroch [A12] implies that non-trivial cobordism implies closed time loops - impossible in TGD).

From TGD point of view this is too strong a condition and in TGD framework space-time surfaces with both Euclidian and Minkowskian signature of the induced metric are allowed. Also cobordisms singular as 4-surfaces are analogous to 3-vertices of Feynman diagrams are allowed.

7.2 Prastaro's Generalization Of Cobordism Concept To The Level Of Partial Differential Equations

I am not enough mathematician in technical sense of the word to develop overall view about what Prastaro has done and I have caught only the basic idea. I have tried to understand the articles [A6, A7] with title "Geometry of PDE's. I/II: Variational PDE's and integral bordism groups" (<http://www.sciencedirect.com/science/article/pii/S0022247X05005998> and <http://www.sciencedirect.com/science/article/pii/S0022247X05008115>), which seem to correspond to my needs. The key idea is to generalize the cobordism concept also to partial differential equations with cobordism replaced with the time evolution defined by partial differential equation. In particular, to geometric variational principles defining as their extremals the counterparts of cobordisms.

Quite generally, and especially so in the case of the conservation of Noether charges give rise to strong selection rules since two n -surfaces with different classical charges cannot be connected by extremals of the variational principle. Note however that the values of the conserved charges depend on the normal derivatives of the imbedding space coordinates at the n -dimensional ends of cobordism. If one poses additional conditions fixing these normal derivatives, the selection rules become even stronger. In TGD framework Bohr orbit property central for the notion of WCW geometry and holography allows to hope that conserved charges depend on 3-surfaces only.

What is so beautiful in this approach that it promises to generalize the notion of cobordism and perhaps also the notions of homotopy/homology groups so that they would apply to partial differential equations quite generally, and especially so in the case of geometric variational principles giving rise to $n + 1$ -surfaces connecting n -surfaces characterizing the initial and final states classically. TGD with $n = 3$ seems to be an ideal applications for these ideas.

Prastaro also proposes a generalization of cobordism theory to super-manifolds and quantum super-manifolds. The generalization in the case of quantum theory utilizing path integral does not pose conditions on classical connecting field configurations. In TGD framework these generalizations are not needed since fermion number is geometrized in terms of imbedding space gamma matrices and super(symplectic) symmetry is realized differently.

7.3 Why Prastaro's Idea Resonates So Strongly With TGD

Before continuing I want to make clear why Prastaro's idea resonates so strongly with TGD.

1. One of the first ideas as I started to develop TGD was that there might be selection rules analogous to those of quantum theory telling which 3-surfaces can be connected by a space-time surface. At that time I still believed in path integral formalism assuming that two 3-surfaces at different time slices with different values of Minkowski time can be connected by any space-time surface for which imbedding space coordinates have first derivatives.

I soon learned about Thom's theory but was greatly disappointed since no selection rules were involved in the category of abstract 3-manifolds. I thought that possible selection rules should result from the imbeddability of the connecting four-manifold to $H = M^4 \times CP_2$ but my gut feeling was that these rules are more or less trivial since so many connecting 4-manifolds exist and some of them are very probably imbeddable.

One possible source of selection rules could have been the condition that the induced metric has Minkowskian signature - one could justify it in terms of classical causality. This restricts strongly topology change in general relativity (<http://arxiv.org/pdf/1406.6194v1.pdf>). Geroch's classical result [A12] states that non-trivial smooth Lorentz cobordism between compact 3-surfaces implies the existence of closed time loop - not possible in TGD framework. Second non-encouraging result is that scalar field propagating in trouser topology leads to an occurrence of infinite energy burst (<http://journals.aps.org/prd/abstract/10.1103/PhysRevD.58.124019>).

In the recent formulation of TGD however also Euclidian signature of the induced metric is allowed. For space-time counterparts of 3-particle vertices three space-time surfaces are glued along their smooth 3-D ends whereas space-time surface fails to be everywhere smooth manifold. This picture fits nicely with the idea that one can engineer space-time surfaces by gluing them together along their ends.

2. At that time (before 1980) the discovery of the geometry of the “World of Classical Worlds” (WCW) as a possible solution to the failures of canonical quantization and path integral formalism was still at distance of ten years in future. Around 1985 I discovered the notion of WCW. I made some unsuccessful trials to construct its geometry, and around 1990 finally realized that 4-D general coordinate invariance is needed although basic objects are 3-D surfaces.

This is realized if classical physics is an exact part of quantum theory - not only something resulting in a stationary phase approximation. Classical variational principle should assign to a 3-surface a physically unique space-time surface - the analog of Bohr orbit - and the action for this surface would define Kähler function defining the Kähler geometry of WCW using standard formula.

This led to a notion of preferred extremal: absolute minimum of Kähler action was the first guess and might indeed make sense in the space-time regions with Euclidian signature of induced metric but not in Minkowskian regions, which give to the vacuum functional and exponential of Minkowskian Kähler action multiplied by imaginary unit coming from \sqrt{g} - just as in quantum field theories. Euclidian regions give the analog of the free energy exponential of thermodynamics and transform path integral to mathematically well-defined functional integral.

3. After having discovered the notion of preferred extremal, I should have also realized that an interesting generalization of cobordism theory might make sense after all, and could even give rise to the classical counterparts of the selection rules! For instance, conservation of isometry charges defines equivalence classes of 3-surfaces endowed with tangent space data. Bohr orbit property could fix the tangent space data (normal derivatives of imbedding space coordinates) so that conserved classical charges would characterize 3-surfaces alone and thus cobordism equivalence classes and become analogous to topological invariants. This would be in spirit with the attribute “Topological” in TGD!

7.4 What Preferred Extremals Are?

The topology of WCW has remained mystery hitherto - partly due to my very limited technical skills and partly by the lack of any real physical idea. The fact, that p-adic topology seems to be natural at least as an effective topology for the maxima of Kähler function of WCW gave a hint but this was not enough.

I hope that the above summary has made clear why the idea about dynamical cobordism and even dynamical homotopy theory is so attractive in TGD framework. One could even hope that dynamics determines not only Kähler geometry but also the topology of WCW to some extent at least! To get some idea what might be involved one must however first tell about the recent situation concerning the notion of preferred extremal.

1. The recent formulation for the notion of preferred extremal relies on strong form of General Coordinate Invariance (SGCI). SGCI states that two kinds of 3-surfaces can be identified as fundamental objects. Either the light-light 3-D

orbits of partonic 2-surfaces defining boundaries between Minkowskian and Euclidian space-time regions or the space-like 3-D ends of space-time surfaces at boundaries of CD. Since both choices are equally good, partonic 2-surfaces and their tangent space-data at the ends of space-time should be the most economic choice.

This eventually led to the realization that partonic 2-surfaces and string world sheets should be enough for the formulation of quantum TGD. Classical fields in the interior of space-time surface would be needed only in quantum measurement theory, which demands classical physics in order to interpret the experiments.

2. The outcome is strong form of holography (SH) stating that quantum physics should be coded by string world sheets and partonic 2-surfaces inside given causal diamond (CD). SH is very much analogous to the AdS/CFT correspondence but is much simpler: the simplicity is made possible by much larger group of conformal symmetries.

If these 2-surfaces satisfy some consistency conditions one can continue them to 4-D space-time surface inside CD such that string world sheets are surfaces inside them satisfying the condition that charged (possibly all) weak gauge potentials identified as components of the induced spinor connection vanish at the string world sheets and also that energy momentum currents flow along these surfaces. String world sheets carry second quantized free induced spinor fields and fermionic oscillator operator basis is used to construct WCW gamma matrices.

3. The 3-surfaces at the ends of WCW must satisfy strong conditions to guarantee effective 2-dimensionality. Quantum criticality suggests the identification of these conditions. All Noether charges assignable to a sub-algebra of super-symplectic algebra isomorphic to it and having conformal weights which are n -multiples of those of entire algebra vanish/annihilate quantum states. One has infinite fractal hierarchy of broken super-conformal symmetries with the property that the the sub-algebra is isomorphic with the entire algebra. This like a ball at the top of ball at the top of

The speculative vision is that super-symplectic subalgebra with weights coming as n -ples of those for the entire algebra acts as an analog of conformal gauge symmetries on light-like orbits of partonic 2-surfaces, and gives rise to a pure gauge degeneracy whereas other elements of super-symplectic algebra act as dynamical symmetries. The hierarchy of quantum criticalities defines hierarchies of symmetry breakings characterized by hierarchies of sub-algebras for which one n_{i+1} is divisible by n_i . The proposal is that conformal gauge invariance means that the analogs of Bohr orbits are determined only apart from conformal gauge transformations forming to n_i conformal equivalence classes so that effectively one has n_i discrete degrees of freedom assignable to light-like partonic orbits.

4. In this framework manifolds M and N would correspond the 3-surfaces at the boundaries of CD and containing a collection strings carrying induced

spinor fields. The connecting 4-surface W would contain string world sheets and the light-like orbits of partonic 2-surfaces as simultaneous boundaries for Minkowskian and Euclidian regions.

Propagator line has several meanings depending on whether one considers particles as strings, as single fermion states localizable at the ends of strings, or as Euclidian space-time regions or their light-like boundaries with singular induced metric having vanishing determinant. Vertices appear as generalizations of the stringy vertices and as generalization of the vertices of Feynman diagrams in which the incoming 4-surfaces meet along their ends.

1. Propagator line has several meanings depending on whether one considers particles as strings, as single fermion states localizable at the ends of strings, or as Euclidian space-time regions or their light-like boundaries with degenerate induced metric with vanishing determinant. Vertices appear as generalizations of the stringy vertices and as generalization of the vertices of Feynman diagrams in which the incoming 4-surfaces meet along their ends.
 - (a) The lines of generalized Feynman graphs defined in topological sense are identified as slightly deformed pieces of CP_2 defining wormhole contacts connecting two Minkowskian regions and having wormhole throats identified as light-like parton orbits as boundaries. Since there is a magnetic monopole flux through the wormhole contacts they must appear as pairs (also larger number is possible) in order that magnetic field lines can close. Elementary particles correspond to pairs of wormhole contacts. At both space-time sheets the throats are connected by magnetic flux tubes carrying monopole flux so that a closed flux tube results having a shape of an extremely flattened square and having wormhole contacts at its ends. It is a matter of taste, whether to call the light-like wormhole throats or their interiors as lines of the generalized Feynman/twistor diagrams.

The light-like orbits of partonic 2-surfaces bring strongly in mind the light-like 3-surfaces along which radiation fields can be restricted - kind of shockwaves at which the signature of the induced space-time metric changes its signature.

- (b) String world sheets as orbits of strings are also in an essential role and could be seen as particle like objects. String world sheets could as kind of singular solutions of field equations analogous to characteristics of hyperbolic differential equations. The isometry currents of Kähler action flow along string world sheets and field equations restricted to them are satisfied. As if one would have 2-dimensional solution. $\sqrt{g_4}$ would of course vanishes for genuinely 2-D solution but this one can argue that this is not a problem since $\sqrt{g_4}$ can be eliminated from field equations. String world sheets could serve as 2-D a analoga for a solution of hyperbolic field equations defining expanding wave front localized at 3-D light-like surface.

- (c) Propagation in the third sense of word is assignable to the ends of string world sheets at the light-like orbits of partonic 2-surfaces and possibly carrying fermion number. One could say that in TGD one has both fundamental fermions serving as building bricks of elementary particles and strings characterizing interactions between particles. Fermion lines are massless in 8-D sense. By strong form of holography this quantum description has 4-D description space-time description as a classical dual.
2. The topological description of interaction vertices brings in the most important deviation from the standard picture behind cobordism: space-time surfaces are not smooth in TGD framework. One allows topological analogs of 3-vertices of Feynman diagrams realized by connecting three 4-surfaces along their smooth 3-D ends. 3-vertex is also an analog (actually much more!) for the replication in biology. This vertex is *not* the analog of stringy trouser vertex for which space-time surface is continuous whereas 3-surface at the vertex is singular (also trouser vertex could appear in TGD).

The analog of trouser vertex for string world sheets means splitting of string and fermionic field modes decompose into superposition of modes propagating along the two branches. For instance, the propagation of photon along two paths could correspond to its geometric decay at trouser vertex not identifiable as “decay” to two separate particles.

For the analog of 3-vertex of Feynman diagram the 3-surface at the vertex is non-singular but space-time surface is singular. The gluing along ends corresponds to genuine 3-particle vertex.

The view about solution of PDEs generalizes dramatically but the general idea about cobordism might make sense also in the generalized context.

7.5 Could Dynamical Homotopy/Homology Groups Characterize WCW Topology?

The challenge is to at least formulate (with my technical background one cannot dream of much more) the analog of cobordism theory in this framework. One can actually hope even the analog of homotopy/homology theory.

1. To a given 3-surface one can assign its cobordism class as the set of 3-surfaces at the opposite boundary of CD connected by a preferred extremal. The 3-surfaces in the same cobordism class are characterized by same conserved classical Noether charges, which become analogs of topological invariants.

One can also consider generalization of cobordisms as analogs to homotopies by allowing return from the opposite boundary of CD. This would give rise to first homotopy groupoid. One can even go back and forth several times. These dynamical cobordisms allow to divide 3-surfaces at given boundary of CD in equivalence classes characterized among other things by same values of conserved charges. One can also return to the original 3-surface. This could give rise to the analog of the first homotopy group Π_1 .

2. If one takes the homotopy interpretation literally one must conclude that the 3-surfaces with different conserved Noether charges cannot be connected by any path in WCW - they belong to disjoint components of the WCW! The zeroth dynamical homotopy group Π_0 of WCW would be non-trivial and its elements would be labelled by the conserved Noether charges defining topological invariants!

The values of the classical Noether charges would label disjoint components of WCW. The topology for the space of these parameters would be totally disconnected - no two points cannot be connected by a continuous path. p-Adic topologies are indeed totally disconnected. Could it be that p-adic topology is natural for the conserved classical Noether charges and the sectors of WCW are characterized by p-adic number fields and their algebraic extensions?

Long time ago I noticed that the 4-D spin glass degeneracy induced by the huge vacuum degeneracy of Kähler action implies analogy between the space of maxima of Kähler function and the energy landscape of spin glass systems [K7]. Ultrametricity (https://en.wikipedia.org/wiki/Ultrametric_space) is the basic property of the topology of the spin glass energy landscape. p-Adic topology is ultrametric and the proposal was that the effective topology for the space of maxima could be p-adic.

3. Isometry charges are the most important Noether charges. These Noether charges are very probably not the only conserved charges. Also the generators in the complement of the gauge sub-algebra of symplectic algebra acting as gauge conformal symmetries could be conserved. All these conserved Noether charges would define a parameter space with a natural p-adic topology.

Since integration is problematic p-adically, one can ask whether only discrete quantum superpositions of 3-surfaces with different classical charges are allowed or whether one should even assume fixed values for the total classical Noether charges appearing in the scattering amplitudes.

I have proposed this kind of approach for the zero modes of WCW geometry not contributing to the Kähler metric except as parameters. The integration for zero modes is also problematic because there is no metric, which would define the integration measure. Since classical charges do not correspond to quantum fluctuating degrees of freedom they should correspond to zero modes. Hence these arguments are equivalent.

The above argument led to the identification of the analogs of the homotopy group Π_0 and led to the idea about homotopy groupoid/group Π_1 . The elements of Π_1 would correspond to space-time surfaces, which run arbitrary number of times fourth and back and return to the initial 3-surface at the boundary of CD. If the two preferred extremals connecting same pair of 3-surfaces can be deformed to each other, one can say that they are equivalent as dynamical homotopies (or cobordisms). What could be the allowed deformations? Are they cobordisms of cobordisms? What this could mean? Could they define the analog of homotopy groupoid Π_2 as foliations of preferred extremals connecting the same 3-surfaces?

1. The number theoretic vision about generalized Feynman diagrams suggests a possible approach. Number theoretic ideas combined with the generalization of twistor approach [K22, K12] led to the vision that generalized Feynman graphs can be identified as sequences or webs of algebraic operations in the co-algebra defined by the Yangian assignable to super-symplectic algebra [A5] [?, ?, ?] and acting as symmetries of TGD. Generalized Feynman graphs would represent algebraic computations. Computations can be done in very many different manners and each of them corresponds to a generalized Feynman diagram. These computations transform give same final collection of “numbers” when the initial collection of “numbers” is given. Does this mean that the corresponding scattering amplitudes must be identical?

If so, a huge generalization of the duality symmetry of the hadronic string models would suggest itself. All computations can be reduced to minimal computations. Accordingly, generalized Feynman diagrams can be reduced to trees by eliminating loops by moving the ends of the loops to same point and snipping the resulting tadpole out! The snipped of tadpole would give a mere multiplicative factor to the amplitude contributing nothing to the scattering rate - just like vacuum bubbles contribute nothing in the case of ordinary Feynman diagrams.

2. How this symmetry could be realized? Could one just assume that only the minimal generalized Feynman diagrams contribute? - not a very attractive option. Or could one hope that only tree diagrams are allowed by the classical dynamics: this was roughly the original vision? The huge vacuum degeneracy of Kähler action implying non-determinism does not encourage this option. The most attractive and most predictive realization conforming with the idea about generalized Feynman diagrammatics as arithmetics would be that all the diagrams differing by these moves give the same result. An analogous symmetry has been discovered for twistor diagrams.
3. Suppose one takes seriously the snipping of a tadpole away from diagram as a move, which does not affect the scattering amplitude. Could this move correspond to an allowed elementary cobordism of preferred extremal? If so, scattering amplitudes would have purely topological meaning as representations of the elements of cobordism classes! TGD would indeed be what it was proposed to be but in much deeper sense than I thought originally. This could also conform with the interpretation of classical charges as topological invariants, realize adelic physics at the level of WCW, and conform with the idea about TGD as almost topological QFT and perhaps generalizing it to topological QFT in generalized sense.
4. One can imagine several interpretations for the snipping operation at space-time level. TGD allows a huge classical vacuum degeneracy: all space-time surfaces having Lagrangian manifold of CP_2 as their CP_2 projection are vacuum extremals of Kähler action. Also all CP_2 extremals having 1-D light-like curve as M^4 projection are vacuum extremals but have non-vanishing Kähler action. This would not matter if one does not have superpositions since mul-

tiplicative factors are eliminated in scattering amplitudes. Could the tadpoles correspond to CP_2 type vacuum extremals at space-time level?

There is also an alternative interpretation. In ZEO causal diamonds (CD) form a hierarchy and one can imagine that the sub-CDs of given CD correspond to quantum fluctuations. Could tadpoles be assigned to sub-CDs of CD be considered+

5. In this manner one could perhaps define elements of homotopy groupoid Π_2 as foliations preferred extremals with same ends - these would be 5-D surfaces. If one has two such 5-D foliations with the same 4-D ends, one can form the reverse of the other and form a closed surface. This would be analogous to a map of S^2 to WCW. If the two 5-D foliations cannot be transformed to each other, one would have something, which might be regarded as a non-trivial element of dynamical homotopy group Π_2 .

One can ask whether one could define also the analogs of higher homology or homotopy groupoids and groupoids Π_3 up to Π_5 - the upper bound $n = 5 = 8 - 3$ comes from the fact that foliations of foliations.. can have maximum dimension $D = 8$ and from the dimension of $D = 3$ of basic objects.

1. One could form a foliation of the foliations of preferred extremals as the element of the homotopy groupoid Π_3 . Could allowed moves reduce to the snipping operation for generalized Feynman diagrams but performed along direction characterized by a new foliation parameter.
2. The topology of the zero mode sector of WCW parameterized by fixed values of conserved Noether charges as element of Π_0 could be characterized by dynamical homotopy groups Π_n , $n = 1, \dots, 5$ - at least partially. These degrees of freedom could correspond to quantum fluctuating degrees of freedom. The Kähler structure of WCW and finite-D analogy suggests that all odd dynamical homotopy groups vanish so that Π_0 , Π_2 and Π_4 would be the only non-trivial dynamical homotopy groups. The vanishing of Π_1 would imply that there is only single minimal generalized Feynman diagram contributing to the scattering amplitude. This also true if Feynman diagrams correspond to arithmetic operations.
3. Whether one should call these groups homotopy groups or homology groups is not obvious. The construction means that the foliations of foliations of ... can be seen as images of spheres suggesting "homotopy". The number of these groups is determined by the dimension of imbedding space, which suggests "homology".
4. Clearly, the surfaces defining the dynamical homotopy groups/groupoids would be analogs of branes of M-theory but would be obtained constructing paths of paths of paths... by starting from preferred extremals. The construction of so called n -groups ([https://en.wikipedia.org/wiki/N-group_\(category_theory\)](https://en.wikipedia.org/wiki/N-group_(category_theory))) brings strongly in mind this construction.

7.6 Appendix: About Field Equations Of TGD In Jet Bundle Formulation

Prastaro utilizes jet bundle (https://en.wikipedia.org/wiki/Jet_bundle) formulation of partial differential equations (PDEs). This notion allows a very terse formulation of general PDEs as compared to the old-fashioned but much more concrete formulation that I have used. The formulation is rather formula rich and reader might lose easily his/her patience since one must do hard work to learn which formulas follow trivially from the basic definitions.

I will describe this formulation in TGD framework briefly but without explicit field equations, which can be found at [K2]. To my view a representation by using a concrete example is always more reader friendly than the general formulas derived in some reference. I explain my view about the general ideas behind jet bundle formulation with minimal number amount of formulas. The reader can find explicit formulas from the Wikipedia link above.

The basic goal is to have a geometric description of PDE. In TGD framework the geometric picture is of course present from beginning: field patterns as 4-surfaces in field space - somewhat formal geometric objects - are replaced with genuine 4-surfaces in $M^4 \times CP_2$.

7.6.1 Field equations as conservation laws, Frobenius integrability conditions, and a connection with quaternion analyticity

The following represents qualitative picture of field equations of TGD trying to emphasize the physical aspects. Also the possibility that Frobenius integrability conditions are satisfied and correspond to quaternion analyticity is discussed.

1. Kähler action is Maxwell action for induced Kähler form and metric expressible in terms of imbedding space coordinates and their gradients. Field equations reduce to those for imbedding space coordinates defining the primary dynamical variables. By GCI only four of them are independent dynamical variables analogous to classical fields.
2. The solution of field equations can be interpreted as a section in fiber bundle. In TGD the fiber bundle is just the Cartesian product $X^4 \times CD \times CP_2$ of space-time surface X^4 and causal diamond $CD \times CP_2$. CD is the intersection of future and past directed light-cones having two light-like boundaries, which are cone-like pieces of light-boundary $\delta M_{\pm}^4 \times CP_2$. Space-time surface serves as base space and $CD \times CP_2$ as fiber. Bundle projection Π is the projection to the factor X^4 . Section corresponds to the map $x \rightarrow h^k(x)$ giving imbedding space coordinates as functions of space-time coordinates. Bundle structure is now trivial and rather formal.

By GCI one could also take suitably chosen 4 coordinates of $CD \times CP_2$ as space-time coordinates, and identify $CD \times CP_2$ as the fiber bundle. The choice of the base space depends on the character of space-time surface. For instance CD , CP_2 or $M^2 \times S^2$ (S^2 a geodesic sphere of CP_2), could define the base space. The bundle projection would be projection from $CD \times CP_2$ to the

base space. Now the fiber bundle structure can be non-trivial and make sense only in some space-time region with same base space.

3. The field equations derived from Kähler action must be satisfied. Even more: one must have a *preferred* extremal of Kähler action. One poses boundary conditions at the 3-D ends of space-time surfaces and at the light-like boundaries of $CD \times CP_2$.

One can fix the values of conserved Noether charges at the ends of CD (total charges are same) and require that the Noether charges associated with a sub-algebra of super-symplectic algebra isomorphic to it and having conformal weights coming as n -ples of those for the entire algebra, vanish. This would realize the effective 2-dimensionality required by SH. One must pose boundary conditions also at the light-like partonic orbits. So called weak form of electric-magnetic duality is at least part of these boundary conditions.

It seems that one must restrict the conformal weights of the entire algebra to be non-negative $r \geq 0$ and those of subalgebra to be positive: $mn > 0$. The condition that also the commutators of sub-algebra generators with those of the entire algebra give rise to vanishing Noether charges implies that all algebra generators with conformal weight $m \geq n$ vanish so the dynamical algebra becomes effectively finite-dimensional. This condition generalizes to the action of super-symplectic algebra generators to physical states.

M^4 time coordinate cannot have vanishing time derivative dm^0/dt so that four-momentum is non-vanishing for non-vacuum extremals. For CP_2 coordinates time derivatives ds^k/dt can vanish and for space-like Minkowski coordinates dm^i/dt can be assumed to be non-vanishing if M^4 projection is 4-dimensional. For CP_2 coordinates $ds^k/dt = 0$ implies the vanishing of electric parts of induced gauge fields. The non-vacuum extremals with the largest conformal gauge symmetry (very small n) would correspond to cosmic string solutions for which induced gauge fields have only magnetic parts. As n increases, also electric parts are generated. Situation becomes increasingly dynamical as conformal gauge symmetry is reduced and dynamical conformal symmetry increases.

4. The field equations involve besides imbedding space coordinates h^k also their partial derivatives up to second order. Induced Kähler form and metric involve first partial derivatives $\partial_\alpha h^k$ and second fundamental form appearing in field equations involves second order partial derivatives $\partial_\alpha \partial_\beta h^k$.

Field equations are hydrodynamical, in other worlds represent conservation laws for the Noether currents associated with the isometries of $M^4 \times CP_2$. By GCI there are only 4 independent dynamical variables so that the conservation of $m \leq 4$ isometry currents is enough if chosen to be independent. The dimension m of the tangent space spanned by the conserved currents can be smaller than 4. For vacuum extremals one has $m = 0$ and for massless extremals (MEs) $m = 1$! The conservation of these currents can be also interpreted as an existence of $m \leq 4$ closed 3-forms defined by the duals of these currents.

5. The hydrodynamical picture suggests that in some situations it might be possible to assign to the conserved currents flow lines of currents even globally. They would define $m \leq 4$ global coordinates for some subset of conserved currents (4+8 for four-momentum and color quantum numbers). Without additional conditions the individual flow lines are well-defined but do not organize to a coherent hydrodynamic flow but are more like orbits of randomly moving gas particles. To achieve global flow the flow lines must satisfy the condition $d\phi^A/dx^\mu = k_B^A J_\mu^B$ or $d\phi^A = k_B^A J^B$ so that one can special of 3-D family of flow lines parallel to $k_B^A J^B$ at each point - I have considered this kind of possibly in [K2] at detail but the treatment is not so general as in the recent case.

Frobenius integrability conditions ([https://en.wikipedia.org/wiki/Frobenius_theorem_\(differential_topology\)](https://en.wikipedia.org/wiki/Frobenius_theorem_(differential_topology))) follow from the condition $d^2\phi^A = 0 = dk_B^A \wedge J^B + k_B^A dJ^B = 0$ and implies that dJ^B is in the ideal of exterior algebra generated by the J^A appearing in $k_B^A J^B$. If Frobenius conditions are satisfied, the field equations can define coordinates for which the coordinate lines are along the basis elements for a sub-space of at most 4-D space defined by conserved currents. Of course, the possibility that for preferred extremals there exists $m \leq 4$ conserved currents satisfying integrability conditions is only a conjecture.

It is quite possible to have $m < 4$. For instance for vacuum extremals the currents vanish identically For MEs various currents are parallel and light-like so that only single light-like coordinate can be defined globally as flow lines. For cosmic strings (cartesian products of minimal surfaces X^2 in M^4 and geodesic spheres S^2 in CP_2 4 independent currents exist). This is expected to be true also for the deformations of cosmic strings defining magnetic flux tubes.

6. Cauchy-Riemann conditions in 2-D situation represent a special case of Frobenius conditions. Now the gradients of real and imaginary parts of complex function $w = w(z) = u + iv$ define two conserved currents by Laplace equations. In TGD isometry currents would be gradients apart from scalar function multipliers and one would have generalization of C-R conditions. In citeallbprefextremals,twistorstory I have considered the possibility that the generalization of Cauchy-Riemann-Fueter conditions [A13, A10] (<http://arxiv.org/pdf/hep-th/9306080v2.pdf>) could define quaternion analyticity - having many non-equivalent variants - as a defining property of preferred extremals. The integrability conditions for the isometry currents would be the natural physical formulation of CRF conditions. Different variants of CRF conditions would correspond to varying number of independent conserved isometry currents.
7. The problem caused by GCI is that there is infinite number of coordinate choices. How to pick a physically preferred coordinate system? One possible manner to do this is to use coordinates for the projection of space-time surface to some preferred sub-space of imbedding - geodesic manifold is an excellent choice. Only $M^1 \times X^3$ geodesic manifolds are not possible but these correspond

to vacuum extremals.

One could also consider a philosophical principle behind integrability. The variational principle itself could give rise to at least some preferred space-time coordinates in the same manner as TGD based quantum physics would realize finite measurement resolution in terms of inclusions of HFFs in terms of hierarchy of quantum criticalities and fermionic strings connecting partonic 2-surfaces. Frobenius integrability of the isometry currents would define some preferred coordinates. Their number need not be the maximal four however.

For instance, for massless extremals only light-like coordinate corresponding to the light-like momentum is obtained. To this one can however assign another local light-like coordinate uniquely to obtain integrable distribution of planes M^2 . The solution ansatz however defines directly an integrable choice of two pairs of coordinates at imbedding space level usable also as space-time coordinates - light-like local direction defining local plane M^2 and polarization direction defining a local plane E^2 . These choices define integrable distributions of orthogonal planes and local hypercomplex and complex coordinates. Pair of analogs of C-R equations is the outcome. I have called these coordinates Hamilton-Jacobi coordinates for M^4 .

8. This picture allows to consider a generalization of the notion of solution of field equation to that of integral manifold (https://en.wikipedia.org/wiki/Integrability_conditions_for_differential_systems). If the number of independent isometry currents is smaller than 4 (possibly locally) and the integrability conditions hold true, lower-dimensional sub-manifolds of space-time surface define integral manifolds as kind of lower-dimensional effective solutions. Genuinely lower-dimensional solutions would of course have vanishing $\sqrt{g_4}$ and vanishing Kähler action.

String world sheets can be regarded as 2-D integral surfaces. Charged (possibly all) weak boson gauge fields vanish at them since otherwise the electromagnetic charge for spinors would not be well-defined. These conditions force string world sheets to be 2-D in the generic case. In special case 4-D space-time region as a whole can satisfy these conditions. Well-definedness of Kähler-Dirac equation [K15, K21] demands that the isometry currents of Kähler action flow along these string world sheets so that one has integral manifold. The integrability conditions would allow $2 < m \leq n$ integrable flows outside the string world sheets, and at string world sheets one or two isometry currents would vanish so that the flows would give rise 2-D independent sub-flow.

9. The method of characteristics (https://en.wikipedia.org/wiki/Method_of_characteristics) is used to solve hyperbolic partial differential equations by reducing them to ordinary differential equations. The (say 4-D) surface representing the solution in the field space has a foliation using 1-D characteristics. The method is especially simple for linear equations but can work also in the non-linear case. For instance, the expansion of wave front can be described in terms of characteristics representing light rays. It can happen that two characteristics intersect and a singularity results. This gives rise to physical phenomena like caustics and shock waves.

In TGD framework the flow lines for a given isometry current in the case of an integrable flow would be analogous to characteristics, and one could also have purely geometric counterparts of shockwaves and caustics. The light-like orbits of partonic 2-surface at which the signature of the induced metric changes from Minkowskian to Euclidian might be seen as an example about the analog of wave front in induced geometry. These surfaces serve as carriers of fermion lines in generalized Feynman diagrams. Could one see the particle vertices at which the 4-D space-time surfaces intersect along their ends as analogs of intersections of characteristics - kind of caustics? At these 3-surfaces the isometry currents should be continuous although the space-time surface has “edge”.

10. The analogy with ordinary analyticity suggests that it might be possible to interpret string world sheets and partonic 2-surfaces appearing in strong form of holography (SH) as co-dimension 2 surfaces analogous to poles of analytic function in complex plane. Light-like 3-surfaces might be seen as analogs of cuts. The coding of analytic function by its singularities could be seen as analog of SH.

7.6.2 Jet bundle formalism

Jet bundle formalism (https://en.wikipedia.org/wiki/Jet_bundle) is a modern manner to formulate PDEs in a coordinate independent manner emphasizing the local algebraic character of field equations. In TGD framework GCI of course guarantees this automatically. Beside this integrability conditions formulated in terms of Cartan’s contact forms are needed.

1. The basic idea is to take the partial derivatives of imbedding space coordinates as functions of space-time coordinates as independent variables. This increases the number of independent variables. Their number depends on the degree of the jet defined and for partial differential equation of order r , for n dependent variables, and for N independent variables the number of new degrees of freedom is determined by r , n , and N -just by counting the total number of various partial derivatives from $k = 0$ to r . For $r = 1$ (first order PDE) it is $N \times (1 + n)$.
2. Jet at given space-time point is defined as a Taylor polynomial of the imbedding space coordinates as functions of space-time coordinates and is characterized by the partial derivatives at various points treated as independent coordinates analogous to imbedding space coordinate. Jet degree r is characterized by the degree of the Taylor polynomial. One can sum and multiply jets just like Taylor polynomials. Jet bundle assigns to the fiber bundle associated with the solutions of PDE corresponding jet bundle with fiber at each point consisting of jets for the independent variables ($CD \times CP_2$ coordinates) as functions of the dependent variables (space-time coordinates).
3. The field equations from the variation of Kähler action are second order partial differential equations and in terms of jet coefficients they reduce to local

algebraic equations plus integrability conditions. Since TGD is very non-linear one obtains polynomial equations at each point - one for each imbedding space coordinate. Their number reduces to four by GCI. The minimum degree of jet bundle is $r = 2$ if one wants algebraic equations since field equations are second order PDEs.

4. The local algebraic conditions are not enough. One must have also conditions stating that the new independent variables associated with partial derivatives of various order reduces to appropriate multiple partial derivatives of imbedding space coordinates. These conditions can be formulated in terms of Cartan's contact forms, whose vanishing states these conditions. For instance, if dh^k is replaced by independent variable u^k , the condition $dh^k - u^k = 0$ is true for the solution surfaces.
5. In TGD framework there are good motivations to break the non-orthodoxy and use 1-jets so that algebraic equations replaced by first order PDEs plus conditions requiring vanishing of contact forms. These equations state the conservation of isometry currents implying that the 3-forms defined by the duals of isometry currents are closed. As found, this formulation reveals in TGD framework the hydrodynamic picture and suggests conditions making the system integrable in Frobenius sense.

REFERENCES

Mathematics

- [A1] Homogenous space. Available at: http://en.wikipedia.org/wiki/Homogenous_space.
- [A2] Kac-Moody algebra. Available at: http://en.wikipedia.org/wiki/KacMoody_algebra.
- [A3] Super Virasoro algebra. Available at: http://en.wikipedia.org/wiki/Super_Virasoro_algebra.
- [A4] Symmetric space. Available at: http://en.wikipedia.org/wiki/Symmetric_space.
- [A5] Yangian symmetry. Available at: <http://en.wikipedia.org/wiki/Yangian>.
- [A6] Prastaro A. Geometry of PDEs: I: Integral bordism groups in PDEs. *J Math Anal & Appl*. Available at: <http://www.sciencedirect.com/science/article/pii/S0022247X05005998>, 319(2):547–566, 2006.
- [A7] Prastaro A. Geometry of PDEs: II: Integral bordism groups in PDEs. *J Math Anal & Appl*. Available at: <http://www.sciencedirect.com/science/article/pii/S0022247X05008115>, 321(2):930–948, 2006.

- [A8] Freed DS. The Geometry of Loop Groups, 1985.
- [A9] Baez JC. The Octonions. *Bull Amer Math Soc*. Available at: <http://math.ucr.edu/home/baez/Octonions/octonions.html>, 39(2002), 2001.
- [A10] Rotelli P Leo de S. A New Definition of Hypercomplex Analyticity. Available at: <http://arxiv.org/pdf/funct-an/9701004.pdf>, 1997.
- [A11] Thom R. *Comm Math Helvet*, 28, 1954.
- [A12] Geroch RP. Topology in general relativity. *J Math Phys*, 8:782–786, 1967.
- [A13] Vandoren S Wit de B, Rocek M. Hypermultiplets, Hyperkähler Cones and Quaternion-Kähler Geometry. Available at: <http://arxiv.org/pdf/hep-th/0101161.pdf>, 2001.

Books related to TGD

- [K1] Pitkänen M. About Nature of Time. In *TGD Inspired Theory of Consciousness*. In online book. Available at: http://tgdtheory.fi/public_html/tgdconsc/tgdconsc.html#timenature, 2006.
- [K2] Pitkänen M. Basic Extremals of Kähler Action. In *Physics in Many-Sheeted Space-Time*. In online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#class, 2006.
- [K3] Pitkänen M. Construction of WCW Kähler Geometry from Symmetry Principles. In *Quantum Physics as Infinite-Dimensional Geometry*. In online book. Available at: http://tgdtheory.fi/public_html/tgdgeom/tgdgeom.html#compl1, 2006.
- [K4] Pitkänen M. Does TGD Predict the Spectrum of Planck Constants? In *Hyper-finite Factors and Dark Matter Hierarchy*. In online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#Planck, 2006.
- [K5] Pitkänen M. Negentropy Maximization Principle. In *TGD Inspired Theory of Consciousness*. In online book. Available at: http://tgdtheory.fi/public_html/tgdconsc/tgdconsc.html#nmpc, 2006.
- [K6] Pitkänen M. Nuclear String Hypothesis. In *Hyper-finite Factors and Dark Matter Hierarchy*. In online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#nuclstring, 2006.
- [K7] Pitkänen M. p-Adic Physics: Physical Ideas. In *TGD as a Generalized Number Theory*. In online book. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#phblocks, 2006.
- [K8] Pitkänen M. Quantum Astrophysics. In *Physics in Many-Sheeted Space-Time*. In online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#qastro, 2006.

- [K9] Pitkänen M. TGD and Astrophysics. In *Physics in Many-Sheeted Space-Time*. In online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#astro, 2006.
- [K10] Pitkänen M. TGD and Cosmology. In *Physics in Many-Sheeted Space-Time*. In online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#cosmo, 2006.
- [K11] Pitkänen M. TGD as a Generalized Number Theory: Quaternions, Octonions, and their Hyper Counterparts. In *TGD as a Generalized Number Theory*. In online book. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#visionb, 2006.
- [K12] Pitkänen M. The classical part of the twistor story. In *Towards M-Matrix*. In online book. Available at: http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#twistorstory, 2006.
- [K13] Pitkänen M. The Relationship Between TGD and GRT. In *Physics in Many-Sheeted Space-Time*. In online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#tgdgrt, 2006.
- [K14] Pitkänen M. Was von Neumann Right After All? In *Hyper-finite Factors and Dark Matter Hierarchy*. In online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#vNeumann, 2006.
- [K15] Pitkänen M. WCW Spinor Structure. In *Quantum Physics as Infinite-Dimensional Geometry*. In online book. Available at: http://tgdtheory.fi/public_html/tgdgeom/tgdgeom.html#cspin, 2006.
- [K16] Pitkänen M. Topological Geometroynamics: Basic Visions. In *TGD based view about living matter and remote mental interactions*. In online book. Available at: http://tgdtheory.fi/public_html/tgdlian/tgdlian.html#lianPTGD, 2012.
- [K17] Pitkänen M. *p-Adic length Scale Hypothesis*. Online book. Available at: http://tgdtheory.fi/public_html/padphys/padphys.html, 2013.
- [K18] Pitkänen M. What are the counterparts of Einstein's equations in TGD? In *Physics in Many-Sheeted Space-Time*. In online book. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#EinsteinTGD, 2013.
- [K19] Pitkänen M. Criticality and dark matter. In *Hyper-finite Factors and Dark Matter Hierarchy*. In online book. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#qcritdark, 2014.
- [K20] Pitkänen M. Quantum gravity, dark matter, and prebiotic evolution. In *Genes and Memes*. In online book. Available at: http://tgdtheory.fi/public_html/genememe/genememe.html#hgrprebio, 2014.

- [K21] Pitkänen M. Recent View about Kähler Geometry and Spin Structure of WCW . In *Quantum Physics as Infinite-Dimensional Geometry*. In online book. Available at: http://tgdtheory.fi/public_html/tgdgeom/tgdgeom.html#wcwnew, 2014.
- [K22] Pitkänen M. Unified Number Theoretical Vision. In *TGD as a Generalized Number Theory*. In online book. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#numbervision, 2014.
- [K23] Pitkänen M. From Principles to Diagrams. In online book. Available at: http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#diagrams, 2016.
- [K24] Pitkänen M. Questions related to the twistor lift of TGD. Available at: http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#twistquestions, 2016.