

# The realization of genetic code in terms of dark nucleon and dark photon triplets

M. Pitkänen,

August 12, 2022

Email: matpitka6@gmail.com.

[http://tgdtheory.com/public\\_html/](http://tgdtheory.com/public_html/).

Recent postal address: Rinnekatu 2-4 A 8, 03620, Karkkila, Finland.

## Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Basic notions and ideas . . . . .	3
1.2	Two models of the genetic code based on dark particles . . . . .	3
1.2.1	Bioharmony model . . . . .	4
1.2.2	Codons as dark 3-nucleons . . . . .	4
1.3	The relationship between the two models of genetic code? . . . . .	4
<b>2</b>	<b>Dark nucleon realization of the genetic code and basic information molecules</b>	<b>5</b>
2.1	The basic vision and the first guess . . . . .	5
2.2	The charge DNA as a guideline . . . . .	6
2.3	Dark nucleons or dark protons? . . . . .	6
2.4	Dark variants of information molecules as analogs of nucleon and $\Delta$ obtained by replacing quarks with dark nucleons . . . . .	7
2.5	Angular momentum of the nuclear string as origin of $5 \oplus 3 \oplus 1$ . . . . .	7
2.6	Various pairings of the information molecules . . . . .	8
2.7	A model for the symmetry breaking of the genetic code . . . . .	9
2.8	Chemical bonds as flux tube links and a realization of dark codons using only dark protons . . . . .	11
2.8.1	The sign of the magnetic flux as bit? . . . . .	11
2.8.2	Dark codons as triplets of dark protons at linked closed flux tubes? . . . . .	13
2.9	Could dark genes be dynamical? . . . . .	13
<b>3</b>	<b>Connection between dark nucleon code and bioharmony</b>	<b>14</b>
3.1	Bioharmony and resonance mechanism for dark photon communications . . . . .	14
3.2	Details of the bioharmony model . . . . .	15
3.3	Bioharmony, dark nucleon code, and icoso-tetrahedral code as a tessellation of $H^3$ . . . . .	16

---

<b>4</b>	<b>Pythagorean number mysticism, music harmony, and genetic code</b>	<b>19</b>
4.1	Rational Platonia . . . . .	19
4.2	Special role of primes 2, 3 and 5 . . . . .	19
4.3	Pythagorean scale . . . . .	20
4.4	Pythagorean scale and genetic code . . . . .	20
4.5	What about dodecahedral harmony and analog of genetic code? . . . . .	20

### Abstract

I have worked for more than 10 years with a proposal for two kinds of realizations of the genetic code. The first realization, bioharmony model, represents genetics as light 3-chords consisting of dark photons. The second realization is in terms of dark proton or nucleon triplets forming closed or open strings. I have considered several variants of both realizations but the details have remained poorly understood and I have spent a considerable time on wrong tracks.

It however seems that the dust is finally settling (I am writing this in the beginning of 2022). One can see the dark nucleon model as a generalization of the quark model of nucleon and  $\Delta$  baryons obtained by replacing u and d quarks with dark nucleons. Galois confinement solves the statistics problem. The nucleons are connected by pionic flux tubes to form a closed string-like entity. The dark variants of DNA, RNA, tRNA, and amino-acids (AAs) follow as a prediction. In the sequel, the notation DDNA, DRNA, DtRNA, DAA will be used for the dark variants of the basic information molecules. One can also understand the small symmetry breaking associated with the genetic code.

A concrete realization of bioharmony in terms of the dark nucleon model for codons emerges. The small symmetry breaking effects - the members of doublet that should code for the same amino acid (or act as stop codons), code for different amino acid (or amino acid and stop), are understood. Also the differences between vertebrate and bacterial codes are understood.

## 1 Introduction

I have worked for more than 10 years with a proposal for a realization of the genetic code in terms of dark proton or nucleon triplets forming closed or open strings. I have considered several variants of the code but the details have remained poorly understood and I have spent a considerable time on wrong tracks. Also the contents of this chapter reflect this wandering.

It however seems that the dust is finally settling (I am writing this in the beginning of 2022). One can see the model as a generalization of the quark model of nucleon and  $\Delta$  baryons obtained by replacing u and d quarks with dark nucleons. The color group solving the statistic problem for  $\Delta$  baryon is in the recent case solved by Galois confinement involving Galois group  $Z_3$  assignable to the codons.

### 1.1 Basic notions and ideas

The basic notions behind the models of genetic code and of biomolecules rely on the notion of dark matter as  $h_{eff} = nh_0$  phases of ordinary matter predicted by number theoretic vision.  $n = h_{eff}/h_0$  serves as a measure for algebraic complexity and as a kind of universal IQ.

Dark matter at the magnetic body (MB) has large  $h_{eff}$  so that it is quantum coherent in long scales and acts as a master controlling ordinary biomatter. The control dynamics at dark level is very simple as compared to the biochemical dynamics, which is a kind of shadow dynamics.

Galois confinement is an essential element of the picture. Physical states are singlets under the action of the Galois group associated with the real polynomial with rational coefficients permuting the roots of the polynomial defined a 4-surface in  $M_c^8$  and mapped by  $M^8 - H$  duality to  $M^4 \times CP_2$ . Among other things, this implies that the quark momenta, which are algebraic integers in an extension of rationals defined by the polynomial, sum up to an ordinary integer when the momentum scale corresponds to the largest ramified prime assignable to the extension.

Galois confinement provides a universal mechanism for the formation of bound states. Dark codons as dark nucleon- and dark photon triplets are Galois confined states behaving like a single quantum unit. Dark 3N-nucleons and dark 3N-photons define dark genes.

### 1.2 Two models of the genetic code based on dark particles

Both models are based on Galois confinement providing a universal mechanism for the formation of bound states in TGD Universe.

### 1.2.1 Bioharmony model

The faces of icosahedron and tetrahedron (and octahedron) are triangles. They would correspond to 3-chords made of dark photons, which in turn represent genetic codons.  $m$

Communications are by dark 3N-photons representing genes and are based cyclotron 3N-resonance. Information coded in the frequency modulation of cyclotron frequencies. The chords serve as address and the message is coded to the frequency modulation. The outcome is sequence of resonances giving rise to pulses. Nerve pulse patterns could emerge by this mechanism.

Biophotons are ordinary photons resulting from the decay of dark N-photons to ordinary photons.

### 1.2.2 Codons as dark 3-nucleons

This work led to a more detailed model of the realization of the genetic code in terms of dark nucleon triplets forming a linear structure as the dark counterpart of linear biomolecule pairing and parallel with it.

The nucleons are connected by pionic flux tubes carrying charge  $0, \pm 1$  to form a closed string-like entity carrying angular momentum  $0, 1, \text{ or } 2$ . The dark variants DDNA, DRNA, DtRNA, DAA of DNA, RNA, tRNA, and amino acids (AA) follow as a prediction. AAs correspond to non-rotating analogs of  $N$  (p,n) and  $\Delta$ , DNA and RNA to rotating analog of  $\Delta$ , and tRNA to rotating analog of  $N$ .

Also the pairings between dark information molecules can be understood to a high degree, and the chemical and functional differences between DNA and RNA could reflect the differences between DDNA and DRNA. The almost exact T-C and A-G symmetries of the third letter of the genetic codon could be seen as reflection of almost exact spin or isospin symmetry. The latter option was considered in [K1] but this work strongly favors spin symmetry. One can understand the numbers of DDNAs coding for given DAA and also the the breaking of spin symmetry. The number of DtRNAs is the minimal 32 and this predicts 1-to-many character of DtRNA-tRNA pairing which would induce wobble base pairing.

## 1.3 The relationship between the two models of genetic code?

The precise relation between the two models of genetic code remained poorly understood for a long time. The connection came from the realization of bioharmony model as so-called icosatetrahedral tessellation of hyperbolic 3-space  $H^3$ , which corresponds to either mass shell in momentum space or light-cone proper time constant hyperboloid [L11].  $H^3$  allows an infinite number of tessellations as analogs of 3-D lattices in the Euclidean 3-space  $E^3$ .

1. Basic biomolecules would correspond to linear sub-lattice-like structures of the icosahedral tessellation formed from triplets of icosahedron, tetrahedron and octahedron. One can say that DDNA codon  $X$  is associated with icosahedron-pair and corresponds to a face. This face represents  $X$  in the bioharmony model and also the entangled dark nucleons at its vertices represent  $X$ .
2. The cyclotron frequencies for the nucleons of  $X$  correspond to the frequencies of a dark 3-photon emitted by the dark 3-nucleon. This picture generalizes to genes represented as dark 3N-codons and dark 3N photons emitted in their communications involving 3N-resonance and frequency modulation yielding a series of resonance peaks at the end of the receiver as an analog of nerve pulse pattern.
3. Hamiltonian cycle must be physically realized at the icosatetrahedron which would serve as the basic structure for all dark counterparts of the information molecules. The simplest option is that the Hamiltonian cycle corresponds to a closed flux tube going through all vertices of the tetra-icosahedron. If the cyclotron frequencies, that is magnetic field strengths, are scaled by factor  $3/2$  (and scaled down to the same octave by octave equivalence) at each step along the cycle, the model of bioharmony is realized in terms of cyclotron frequencies. The codon realized as 3 dark nucleons associated with corresponds to the codon realized as dark photon triplet.

4. This picture generalizes to DtRNAs and DAAs. DtRNAs would have as active faces those DtRNA codons, which pair with DmRNA codons. DAAs would have at active faces those DtRNA codons which pair with them. Common dark codon would make pairing by dark 3-photon resonance possible. DtRNA could attach to correct DmRNA during translation and for DAA to correct DtRNA. Ordinary biomolecules would be paired with their dark variants so that dark variants of basic processes would induce their biochemical variants as a kind of shadow dynamics. The pairing by 3N-resonance could be a completely general mechanism involved with biocatalysis.

A concrete realization of bioharmony [?] in terms of the dark nucleon model for codons emerges. The small symmetry breaking effects - the members of doublet that should code for the same amino acid (or act as stop codons), code for different amino acid (or amino acid and stop), are understood. A crucial piece of the puzzle is that one particular chord  $CEG\sharp$  has identical intervals between the notes in even tempered scale. Also the failure of the Pythagorean quint cycle (notes are obtained by scaling the basic frequency by power powers of  $(3/2)$  and using octave equivalence) to close, which bothered Pythagoras, is in an essential role. Also the differences between vertebrate and bacterial codes are understood.

## 2 Dark nucleon realization of the genetic code and basic information molecules

In this section I will represent the arguments leading to the recent (2022) view about dark nucleon realization of the genetic code and dark counterparts of basic information molecules DNA, RNA, tRNA, and amino acids.

### 2.1 The basic vision and the first guess

The basic vision is that dark matter and magnetic body (MB) serves as a master controlling the dynamics of the ordinary biomatter so that its dynamics is shadow dynamics, and the huge complexity of living matter could reduce to relatively simple control dynamics at the level of MB.

Biomolecules are information molecules and dark matter has a higher "IQ" than ordinary matter measured as the dimension  $n = h_{eff}/h_0$  of the extension of rationals associated with the space-time regions characterizing the system measuring also the scale of quantum coherence. Therefore the natural expectation is that basic information molecules have dark counterparts and genetic code is realized at a darl level so that the chemical realization would be a secondary realization.

1. I started with a proposal [K5, K7, K4, K8] that dark codons could correspond to dark nucleons, that is dark quark triplets, assignable to open or closed string like objects (flux tubes). This led to the proposal for the basic group theoretical decomposition DDNA and DRNA codons in terms of group representations of  $SU(2)_I \times SU(2)_R$  as  $(4_I \otimes 4_s \oplus 2_I \otimes 2_s) \otimes (5_s \oplus 3_s) = [(3/2, -3/2)_I + (1/2, -1/2)_I] \otimes (4_s \oplus 2 \otimes 2) \otimes (5_s \oplus 3_s) = 64_{DNA} \oplus 64_{RNA}$ . In the quark model, this corresponds in fermionic degrees of freedom to nucleon  $N$  and  $\Delta$  assuming color degrees of freedom to get statistics right.  $5 \oplus 3$  could be assigned to 2 *rho*-meson-like bonds for an open string-like object.
2. The realization of  $4 \times 4 \oplus 2 \oplus 2$  in terms of quark color triplets is not physically plausible and the challenge is to realize  $4 \times 4 \oplus 2 \oplus 2$  and  $5 \oplus 3$  physically in terms of more plausible dark states. The natural guess is that this realization involves dark protons and neutrons. u and d would be replaced with p and n.
3. Color would be absent and since the  $\Delta$  is completely symmetric, antisymmetry required by Fermi statistics must be realized by bringing in some new degrees of freedom. Galois confinement is suggestive in the TGD framework [L11, L12, L13, L14, L15].  $Z_3$  is a natural guess for the Galois group in the case of codons and one has 3 states in geometric degrees of freedom and 3-nucleon state, which is  $Z_3$  singlet would be antisymmetric.

Since the induced spinors do not have color as spin-like quantum numbers, one must leave open the possibility that even ordinary color confinement has this kind of description as

a description at space-time level (as opposed to the descriptions at embedding space and "world of classical worlds" (WCW) level).

## 2.2 The charge DNA as a guideline

A strong constraint comes from the observation that DDNA and DRNA codons have charge -3.

1. If DDNA and DNA form parallel string-like structures, this strongly suggests that there is a neutralizing charge +3 associated with the dark codon paired with the ordinary codon. This charge could be assigned either to 3 protons or 3 nucleons if there is some additional charge allowing to take care that the total charge is 3 units in the case of DDNA and DRNA.

If also the notions of DAA and DtRNA make sense, charge neutrality for them is a plausible option. Of course, the additional charges could be dynamical in the same way as the charges of corresponding bio-molecules.

2. The compensating charge could be assigned to meson-like states with charges  $0, \pm 1$ . They could be meson-like bonds connecting the dark nucleons to a string-like object.  $\rho$  mesons with spin 1 and charges  $(0, \pm 1)$  and pions with spin 0 and charges  $(0, \pm 1)$  are the natural candidates. For closed string-like objects one would have 3 bonds and for open string-like objects 2 bonds.

For  $\rho$  mesons associated with two bonds one would have spin representations  $3 \otimes 3 = 5 \oplus 3 \oplus 1$ . One should somehow get rid of the spin singlet if one assumes the proposal considered above. Spin-statistics represents the second problem: Bose-Einstein statistics does not allow 3.

For  $3 \otimes 3 \otimes 3 = (7 \oplus 5 \oplus 3) \oplus (5 \oplus 3 \oplus 1) \oplus 3$  associated with closed string-like objects, the number of states is quite too high. The only completely symmetric representation is 7 whereas 1 is antisymmetric. Thus it seems that the  $\rho$ -meson option is not realistic so that only pionic realization remains.

This leaves open only the possibility that  $5 \oplus 3(\oplus 1)$  corresponds to the rotational degrees of freedom of a closed or open string-like object.

## 2.3 Dark nucleons or dark protons?

Are both dark  $p$  and  $n$  needed or are dark protons enough as mildly suggested by the model [L1] of Pollack effect [I3, L1, I7, I6]? Could one build the needed states using only dark protons and suitably chosen bonds?

1. Could number theoretic Galois degrees of freedom [L10, L9, L5, L6, L12, L14, L15] come to rescue? The analog of isospin could be assigned with Galois degrees of freedom. For 3-D algebraic extension of rationals replacing color, the extension increases the algebraic dimension of the 3-space consisting of rational points by factor  $n = 3$  to  $nd$ ,  $d = 3$ . The dimension of number theoretic spinors would be  $2^{n(d-1)}/2 = 2^3$ , which is much larger than the dimension  $d = 2$  of isospin spinors in 3-D space. One could speak of Galois-spin or G-spin. Nucleon isospin is therefore a more reasonable candidate.
2. The objection against the dark nucleon-triplet picture is that, in the standard nuclear physics, neutrons are not thought to be important in living matter. Note however that the dark electroweak length scale scaled by  $h_{eff}/h_0$  could be much longer than the ordinary weak scale, even of the order of cell length scale.

Weak gauge bosons would be effectively massless below the dark weak scale and weak interactions associated with the dark pion-like bonds would be as strong as electromagnetic interactions. This could explain the mysterious large chirality breaking effects in living matter.

3. One can also ask whether the dark neutron is effectively a dark proton plus pionic bond so that dark protons would be the basic building bricks after all. One cannot exclude the possibility that this applies also to the neutrons of ordinary nuclei [K6, K5]. This does not however conform with charge +3 for DDNA codons. This would leave the option that the dark neutron is an ordinary proton plus dark  $\pi^-$  bond.

## 2.4 Dark variants of information molecules as analogs of nucleon and $\Delta$ obtained by replacing quarks with dark nucleons

Could the dark analogs of  $N$  and  $\Delta$  with quarks replaced by nucleons give rise to the genetic code and dark analogs of the basic information molecules?

1. The analogs of  $N$  and  $\Delta$  would give 2 spin doublets (counterparts of p and n as ppn and pnn) and 4 spin 4-plets as a counterpart of  $\Delta$  (ppp,ppn,pnn,nnn): altogether 20 states, which brings into mind AAs. Note that pion bonds could modify the charges for ordinary nucleons.
2. The analogs of  $N$  and  $\Delta$  can be tensored with  $A = 5_s \oplus 3_s \oplus 1_s$  or  $B = 5_s \oplus 3_s$ .  $\Delta \otimes B$  would give  $[(3/2, -3/2)_I \oplus (1/2, -1/2)_I] \otimes (5_s \oplus 3_s)$ : these two 64-plets could be identified as DDNA and DRNA.  
 $N \otimes 1$  could give 20 AAs. For both options,  $N \otimes B = (2_I \oplus 2_s) \otimes (5_s \oplus 3_s)$  would give  $20+12=32$  states, which is the minimal number of tRNA codons. The number of chemical tRNA codons is larger than 40 so that DtRNA-tRNA pairing would be 1-to-many. This could induce the wobble base pairing [I1].
3. DDNA and DRNA would correspond to the analogs of  $\Delta$  nucleons with rotation. For instance, ppp and ppn as counterparts of  $\Delta^{++}$  and  $\Delta^+$  could correspond to DDNA and pnn and nnn as counterparts of  $\Delta^-$  and  $\Delta^0$  could correspond to DRNA. This implies charge asymmetry which should relate to the differences between DNA and RNA. DtRNA would correspond to  $N$  with rotation. DAA would correspond to  $\Delta$  and  $N$  without rotation, which should relate to the different functions between AAs and molecules DNA, RNA, and tRNA.

**Remark:** Note that pionic bonds would guarantee that the total charge of DDNA and DRNA codons is 3 units.

## 2.5 Angular momentum of the nuclear string as origin of $5 \oplus 3 \oplus 1$

One must understand the origin of  $5 \oplus 3 \oplus 1$  or of  $5 \oplus 3$ . There are 3 scenarios to consider. Closed string scenario favors  $5 \oplus 3$  but does not predict AAs. Open string scenario favors  $5 \oplus 3 \oplus 1$  and allows the identification of dark counterparts of all basic biomolecules. If  $\rho$  mesons do *not* give rise to  $5 \oplus 3 \oplus 1$ , both closed and open strings can be considered.

1. What came first into mind was that the bonds between protons are analogous to  $\rho$  mesons. This would allow only an open string option. There is however a problem with statistics: 3 is antisymmetric (analogous to the cross product of 3-vectors).  
This suggests that the bonds are pionic and do not contribute to the spin but allow to obtain desired total charges for 3 proton states. Charge neutrality is attractive for AAs and would require maximum neutralizing charge -3 so that only the closed string option with ordinary nucleons plus dark pionic bonds remains. Even dynamical charges would be possible also at the level of dark bio-molecules and one can consider the possibility that the MB controls the charge state of the basic biomolecules.
2. Concerning the identification of  $5 \oplus 3(\oplus 1)$ , the rotational degrees of freedom of string seem to be the only reasonable option. 1 and 5 could correspond to spin 0 and spin 2 states of the Regge trajectory and 3 to spin 1 state of a possibly exchange degenerate trajectory. What is encouraging is that only the bosonic spins 0, 1, and 2 and fermionic spins 1/2 and 3/2, which are in a very special role physically would be needed.

To sum up, closed strings with dark nucleon triplets and stringy rotational degrees of freedom allow us to predict the dark counterparts of all basic information molecules as analogs of nucleon and  $\Delta$  states with pionic bonds. The number of dark tRNA codons is predicted to be minimal and equal to 32, and the considerably larger number of the chemical tRNA codons implying that dark tRNA-tRNA pairing is 1-to-many. This would explain wobble base pairing.

## 2.6 Various pairings of the information molecules

The basic vision is that the dynamics of the MB induces the dynamics of the biological body and the observed chemical pairings are induced by dark pairings. One should therefore understand the DDNA-DRNA, DRNA-DtRNA, and DtRNA-DAA pairings.

1. DDNA-DRNA pairing is obtained trivially. DDNA-DAA pairing is induced by DRNA-DtrNA pairing and DtRNA-AA pairing.
2. The decomposition  $2 \times (20 \oplus 12)$  for DRNA suggests a natural pairing with tRNA identified as  $20 \oplus 12$ . The spin contents of the codons are however different and the situation is more complex. This leads to a model for the breaking of the (A,G) symmetry of the third codon in RNA-AA pairing. The pairing of DtRNA with DAA requires pairing of  $20 \oplus 12$  with 20.  $20 \rightarrow 20$  is a natural pairing but how to realize  $12 \rightarrow 20$ ?
3. Could icoso-tetrahedral realization of the genetic code in terms of dark photon triplets (bioharmony) [K3] [L8, L11] help here? In this realization the faces of icosahedron and tetrahedron identifiable as 3-chords correspond to codons and 3 isohedral Hamiltonian cycles providing a model for 12-note scale and the unique tetrahedral Hamiltonian cycle are needed for the realization of bioharmony as genetic code.

Icosahedron has 20 faces and 12 vertices defining Hamiltonian cycles essential for the realization in terms of bioharmony. Hamiltonian cycles have  $Z_6$ ,  $Z^4$  or  $Z^2$  as a symmetry group:  $Z_2$  can correspond to reflection of rotation by  $\pi$ .  $Z_6$  has 3 orbits with 6 faces and 1 orbit with 2 faces.  $Z_4$  has 5 orbits with 4 faces.  $Z_2$  has 10 orbits with 2 faces.

Could the missing faces correspond to the missing orbits for one of these symmetry groups: a) to 6-orbit and 2-orbit for  $Z_6$  or b) 2 4-orbits for  $Z_4$  or c) 4 2-orbits for  $Z_2$ .  $Z_6$  and  $Z_4$  Hamiltonian cycles are unique and part of any realization. Option b) is more symmetric than option a) and is a more promising candidate.

Consider now the pairings of type DX-X.

1. All dark pairings as a bound state formation by Galois confinement [L14, L15] would involve formation of a composite  $P_n \circ P_{n-1} \circ \dots \circ P_1$  of the polynomials  $P_i$  determining at  $M^8$  level the 4-surfaces of systems participating in the interaction. This implies that Galois groups extend to a larger group having the Galois groups of composites as normal subgroups.
2. The already mentioned charge asymmetry reflecting the violation of the weak isospin symmetry between DDNA and DRNA could explain why DNA *resp.* RNA involves deoxyribose *resp.* ribose molecule and the nucleotide T is replaced with U in RNA. The instability of RNA molecules and the rarity and short life-time of double RNA strands could derive from the properties of DRNA .

The almost exact T-C and A-G symmetries of the third letter of genetic codon could also reflect almost exact isospin symmetry as proposed in [K1]. The number of DtRNAs is the minimal 32 and this predicts 1-to-many character of DtRNA-tRNA pairing which would induce wobble base pairing.

Dark base pairing could involve an extension of Galois group  $Z_3$  of codon to  $Z_6$  of codon pair. This could make DDNA double strand stable and perhaps induce the stability of DNA double strand. DDNA double strand would permanently be in the bound state with  $Z_6$  as the Galois group of the dark base pair. This would support the view that DDNA is above DRNA in the dark master-slave hierarchy.

Also the functional differences between DNA and RNA could relate to the differences of their dark counterparts. The DDNA double strand with larger  $h_{eff}$  would represent a higher evolutionary level than the DRNA strand.

3. The number of DtRNA molecules is 32, and minimal one, so that DtRNA-tRNA pairing is not unique. This explains the wobbling of RNA-tRNA pairing [I1]. Does the wobble phenomenon have some biological function or does it signal that dark tRNA-tRNA pairing has not yet evolved to its final form?

That tRNA as such does not represent information storage but plays a role of servant in the translation process, supports the first view. The basic function of dark tRNA in the translation is unique but it is less risky to have several manners to perform this function: hence the large number of ordinary tRNAs.

## 2.7 A model for the symmetry breaking of the genetic code

The model predicts that the numbers of DRNA and DtRNAs are 64 and 32 respectively. This condition does not force DRNA-DtRNA correspondence to be 2-1 in a codon-wise way. This is however true in an excellent approximation as becomes clear by looking at the code table.

For the third letter, RNA-AA correspondence has an exact U-C symmetry and almost exact A-G symmetry. There are only 2 exceptions. TTX 4-plet decomposes to  $3 \times \text{ile} + 1 \text{ met}$ : (A,G) doublet for the third codon splits to (ile,met). The (A,G) doublet in TGX splits to (stop,trp). Both stop codons and met as a start codon are therefore very special.

In bacterial genetic codes also the (A,G) doublet in TGX, which usually corresponds to (stop,stop), corresponds sometimes to (stop,pyl) doublet so that CG symmetry is broken. Also the (A,G) doublet usually mapped to (stop,trp) can be mapped to (sec,trp). The interpretation would be that a stop codon is obtained if DtRNA corresponding to UAG or UGA does not pair at all with tRNA. If it pairs, UAG codes for pyl and UGA codes for sec.

One should understand this symmetry breaking.

1. Since iso-spin and spin are involved, either isospin or spin symmetry breaking is suggestive. In the nucleon sector the situation is completely symmetric between spin and isospin. In the string sector, the situation is different for DDNA, DRNA and DtRNA.
2. The earlier interpretation for (U,C) and (A,G) doublets was as isospin doublets and isospin symmetry breaking. The conjugations  $G \leftrightarrow C$  and  $U \leftrightarrow A$  were interpreted as an analog of particle-antiparticle conjugation.

The following model leads to the proposal that dark (T,C) doublet corresponds to spin (rather than isospin-) doublet  $2_s = (1/2, -1/2)$  and (A,G) doublet to pseudo-spin doublet  $(3/2, -3/2)$ . As if rotational symmetry would have reduced to axial symmetry (this would conform with the linear structure of DNA). Letter and its conjugate would correspond to different spin doublets. Interestingly, the repetitive purine (A, and G) sequences for intronic portions of active DNA strand would correspond to dark  $(3/2, -3/2)$  doublet for which the breaking of rotational symmetry is larger for the active strands in the transcribed portion of DNA.

At the level of DNA, DRNA and DtRNA, the natural possibility is that (T,C) doublet corresponds to  $2_s$  and (A,G) doublet to the spin symmetry violating  $(3/2, -3/2)_s$ . (T,C) and (A,G) could form an isospin doublet.

3. An important point to note is that  $(3 \times \text{ile,met})$  and (stop,stop), (stop,trp) dot correspond to identical situations since 2 ile correspond to (T,C) for which there is no symmetry breaking. Actually one has 3 (A,G) symmetry breakings.

Consider now the identification of spin- and isospin contents of various dark information molecules.

1. Suppose that the spin symmetry is not broken at DDNA and DRNA level but isospin 4-plet  $4_I = (3/2, 1/2, -1/2, -3/2)$  splits into pseudo-doublets  $2_{I_1} = (1/2, -1/2)$  and  $2_{I_2} = (3/2, -3/2)$ . If DDNA were maximally symmetric it would correspond to  $2_{I_1} = (1/2, -1/2)$ . Which option one chooses, does not matter in the sequel so that this option is selected.

This would give

$$\begin{aligned} DDNA &= 2_{I_1} \otimes [4_s \otimes (5_s \oplus 3_s)] , \\ DRNA &= 2_{I_2} \otimes [4_s \otimes (5_s \oplus 3_s)] . \end{aligned}$$

2. DtrRNA and DAA correspond to

$$\begin{aligned} DtrRNA &= 2_I \otimes [2_s \otimes (5_s \oplus 3_s)] , \\ DAA &= 4_I \otimes 4_s \oplus 2_I \otimes 2_s . \end{aligned}$$

3. One would expect that the pairing minimizes the breaking of rotational symmetry meaning that spins are the same for paired dark molecules if possible and the spin difference is minimized otherwise. To get some idea about the symmetry breaking, one can decompose the tensor products for the spin representations

$$\begin{aligned} DRNA &= 2_{I_2} \otimes (8_s \oplus 2 \times 6_s \oplus 2 \times 4_s \oplus 2 \times 2_s) , \\ DtrRNA &= 2_I \otimes (6_s \oplus 2 \times 4_s \oplus 2_s) . \end{aligned}$$

The representation contents are different and the number of spin states for DRNA is twice that for DtrRNA so that the symmetry breaking relates to spin pairing rather than isospin pairing.

The first thing to notice is that  $2 \times 2_s$  for DRNA naturally projects to  $2_s$  for DtrRNA. Also  $2 \times 6_s$  projects to  $6_s$ . Both decompositions however have  $2 \times 4_s$ 's so that  $8_s$  must pair with  $2 \times 4_s$ . Symmetry breaking must localize to this pairing.

$8_s$  is not present in DtrRNA and forces a pairing between different spins. This should cause the violation of spin symmetry for dark (A,G) doublets in the sense that they couple to different DAAs, which in turn requires that they couple to different DtrRNAs.

1. One can decompose  $8_s$  as

$$8_s = (7/2, -7/2)_s \oplus (5/2, -5/2)_s \oplus 4_{s_1} .$$

$8_s$  should correspond to  $2 \times 4_s$  in DtrRNA. The pseudo 4-plet  $4_{s_1}$  pairs with  $4_s$  in a spin conserving manner.

2. What is left is  $2_{I_2} \otimes [(7/2, -7/2)_s \oplus (5/2, -5/2)_s]$ , which should pair  $2_I \times 4_s$ . This pairing cannot conserve spin and the 2-1 symmetry must be violated in the sense that the DRNAs paired with  $(3/2)_s$  and  $-(3/2)_s$  are different. One can ask whether the change of the magnitude of the spin component is minimal in the DRNA-DtrRNA pairing.
3. At the level of DAA and DtrRNA, ile could correspond to the first DtrRNA doublet  $((3/2)_I \otimes (1/2, -1/2)_s$  and  $(3/2)_I \otimes (3/2)_s$  as a singlet and met to  $(3/2)_I \otimes (-3/2)_s$  as a singlet.  $(stop, stop)$  and  $(stop, trp)$  could correspond to  $(3/2)_I \otimes (3/2, -3/2)_s$  and  $(-3/2)_I \otimes (3/2, -3/2)_s$ . Spin symmetry breaking would therefore mean that different DRNAs pair with DtrRNAs in the doublet  $(3/2, -3/2)_s$ .

How do DRNAs and DtrRNAs correspond to each other in the (A,G) symmetry violating sector? In the absence of symmetry breaking DRNA-DtrRNA pairing is 2-1 in a codon-wise way if DRNA with opposite values  $3/2$  and  $-3/2$  of isospin pair with the same DtrRNA. Symmetry breaking would mean that some DRNAs with spins  $3/2$  and  $-3/2$  pair with different DtrRNAs and therefore with different DAAs for some (A,G) doublets  $(3/2, -3/2)_s$ . For the (T,C) doublets  $(1/2, -1/2)_s$  this would not occur.

1. There are 8 spin symmetry violating DRNAs and 8 DtrRNAs corresponding to  $UA(A, G)$  and  $UG(A, G)$  and  $UA(T, C)$  and  $UA(A, G)$  ( $3 \times$  ile+ met).  $UA(T, C)$  is strictly speaking not spin symmetry violating but ile corresponds to DRNA triplet instead of doublet. As if the DRNA doublet paired with 2 DtrRNAs pairing with met would pair with DtrRNAs coding for ile and met.

There are 2  $[(7/2, -7/2)_s \oplus (5/2, -5/2)_s]$  multiplets at the DRNA side. At the DtrRNA side one has pseudo doublets  $(3/2, -3/2) \oplus 2_{s_1}$ ,  $s_1 = (1/2, -1/2)$ . There are two of these corresponding isospin doublet  $2_{I_1}$ .

2. Since there are 3 UCAG 4-plets with symmetry breaking, the symmetry violation is not independent of the value of isospin for  $(3/2, -3/2)_I$  for DRNA and  $(1/2, -1/2)_I$  for DtrRNA. Symmetry breaking for isospin should localize to the DRNA side. There are several options  $(3,0)$ ,  $(0,3)$ ,  $(2,1)$   $(1,2)$  for the numbers of symmetry breakings for DRNA multiplets. One can restrict in the sequel to a single value of isospin, say  $3/2$ .
3. One should find symmetry violating pairing between these 8 DRNAs and 8 DtrRNAs.  $[(7/2, -7/2)_s \oplus (5/2, -5/2)_s]$  should be mapped to  $(3/2, -3/2)_s \oplus 2_{s_1}$ . Assume that the spin difference between paired DRNA and DtrRNA is as small as possible.
  - (a) (T,C) doublet without symmetry breaking would correspond to the pairing

$$(5/2, -5/2)_s \rightarrow (1/2, -1/2)_s .$$

(T,C) symmetry is not violated if the both doublets correspond to the same DAA and DtrRNA.

- (b) (A,G) doublet could correspond to the pairing

$$(7/2, -7/2)_s \rightarrow (3/2, -3/2)_s .$$

Now the members of the doublets would correspond to different DAA.

- (c) For instance, the DRNA corresponding to the second met in (met,met) in absence of symmetry breaking, would pair with DtrRNA, which pairs with ile. The symmetry present at the DRNA level would be broken by the pairing.

In the case of (stop,trp) doublet, the same would occur. This would also happen in the replacements (stop,stop)  $\rightarrow$  (stop,pyl) and (stop,trp) $\rightarrow$  (sec,trp). Now the DtrRNA in question would not pair at all with tRNA and AA or it would be with exotic tRNA pairing with an exotic AA.

## 2.8 Chemical bonds as flux tube links and a realization of dark codons using only dark protons

In the proposed model of dark DNA, one must assume that the dark codon is formed by a triplet of dark nucleons (proton and neutron). In the TGD framework one could justify the presence of neutrons by the large value of Planck constant increasing the weak scale to at least atomic length scale so that weak bosons would behave like massless particles in atomic scales at the MB. Therefore the dark protons could transform to dark neutrons easily. Neutron would be connected to either neighbor by a meson-like flux tube bond which is positively charged so that each codon would have a charge of 3 units neutralized by an opposite charge of 3 phosphates.

### 2.8.1 The sign of the magnetic flux as bit?

The introduction of neutrons brings in an additional bit. Therefore one could use only dark protons, if one could bring in this additional bit in some way. An obvious candidate would be the direction of a monopole magnetic flux assignable to the letter of the codon as a closed flux tube with respect to reference direction defined by the DNA sequence. If the letters of codon are closed linked flux tubes containing dark protons forming dark DNA as a chain, this kind of option might work.

Consider first the topology of the monopole flux tubes.

1. Magnetic monopole flux tubes correspond to closed 3-surfaces in the TGD framework. They are closed because the boundary conditions do not allow boundaries with a monopole charge nor boundaries at all. In dimension 3, these flux tubes can become knotted and closed flux tubes can get linked.
2. If one has a braiding of  $N$  flux tubes, one can connect the ends of the  $N$  flux tubes. There are many manners to connect the ends, and one obtains at most  $N$  linked closed flux tubes, which are knots. The simplest option is that the ends of each braid strand are connected so that one has  $N$  linked flux tubes. This corresponds to the "upper" ends as a trivial permutation of the "lower" ends.

- Any permutation in the permutation group  $S_N$  is possible. A given permutation can be expressed as a product of permutations such that each permutation leaves invariant a subset. Permutations are therefore characterized by a partition of  $N$  objects to subsets such that the given set consist of  $N_i$  objects with  $\sum N_i = N$  and that these sets do not decompose to smaller subsets. The allowed permutations for  $N_i$  objects correspond to elements of the cyclic group  $Z_{N_i}$ . These cyclic permutations give rise to a single closed tube when the ends of the braid ends and permuted braid ends are connected. The number of closed flux tubes is therefore the number of summands in  $\sum N_i = N$ .

These permutations are obtained by reconnections from the permutation corresponding to  $N$  closed loops so that there are two levels: the level of braiding and the level of reconnections behind the stages not visible in the properties of the braiding.

Linking is a metaphor for bonding. One speaks of the chain of generations, of a weak link in the chain, etc.

- Chemical bonds are classified into ionic bonds, valence bonds involving delocalization of electrons, and hydrogen bonds involving delocalization of protons. Chemical bonds are not well-understood in the framework of standard chemistry. TGD suggests that they involve space-time topology: monopole flux tube pairs would be associated with the bonds and the splitting of the bond would correspond to a reconnection splitting the pair to two U-shaped flux tubes. Flux tubes and connecting molecules as nodes are proposed to form a network.
- I have not considered in detail how the U-shaped flux tubes are associated with the nodes. Bonding=linking metaphor encourages a crazy question. The members of the flux tube pairs, which are proposed to connect molecules, which serve as nodes of a network. These flux tubes must close and could be linked with shorter closed flux tubes assignable to molecules.
- Could this linking bind the molecules and atoms to a single topological structure. If so, both chemistry and topological quantum computation (TQC) in the TGD framework would involve linking, braiding, and reconnections as new topological elements. Biomatter at molecular level would consist of chains of closed flux tubes which can be also stretched and give rise to braids.

Note that 2 U-shaped flux tubes can reconnect and this transition can lead to a pair of flux tubes or to a linked pair of U-shaped flux tubes so that 3 different states are possible.

- I have proposed that the pairing of molecules by a pair of monopole flux tubes serves as a correlate for entanglement. If dark protons are associated with closed flux tubes, they must entangle. Could also the linking of the U-shaped flux tubes give rise to entanglement? Stable linking correlates the positions of the flux tubes but this need not mean entanglement since wave function can be a product of wave functions in cm coordinates and relative coordinates.

Linking as an additional topological element inspires some quantum chemical and -biological speculations.

- Could the presence of valence-/hydrogen bonds involve a closed flux tube at which the electron (pair)/proton is delocalized and that this flux tube is linked with another such flux tube. This picture is consistent with the proposed role of quantum gravitation in metabolism [L16] and generation of the predecessor of the nervous system [L17] based on very long variants of hydrogen bonds characterized by gravitational Planck constant. In this view, living matter would be an extremely highly organized structure whereas in the standard chemistry organism would be a soup of biomolecules.
- What comes to mind as an example, is the secondary structure of proteins (<https://cutt.ly/sZ5rRiQ>) involving  $\alpha$ - helices,  $\beta$ -strands and  $\beta$ -sheets. Tertiary structure refers to 3-D structure created by a single protein molecule. It can have several domains. There are also quaternary structures formed by several polypeptide chains. Proteins consist of relatively few substructures known as domains, motives and folds. Could these structures involve braided and linked flux tube structures with dynamical reconnections?

### 2.8.2 Dark codons as triplets of dark protons at linked closed flux tubes?

Consider now a possible model of dark DNA involving only dark protons.

1. One can imagine that dark protons are associated with closed flux tubes acting as hydrogen bonds, such that 3 closed flux tubes as letters are linked to form a dark codon. The dark codons could in turn be linked to form genes as sequences of codons. The direction of the magnetic flux can be opposite or parallel to that of the chain so that each closed flux tube represents a bit of topological information. The chains of links would define sequences of bits and even qubits. Could this define the predecessor of the genetic code for which letter represents a single bit?
2. If one has only dark protons, one obtains only 32 dark codons. An additional bit is required to get 64 codons. Could the direction of the closed flux tube in the chain provide the missing bit and thus represent strong isospin distinguishing between p and n?

What implications could this identification have?

1. It is known that the genetic code has a slightly broken symmetry with respect to the last letter of the codon. For almost all RNA codons U and C resp. A and G define code for the same amino-acid. A possible interpretation of the symmetry is that this symmetry is that U-C pair and A-G pair correspond to the bit defined by magnetic flux so that the sign of magnetic flux would not matter much at the level of proteins. For this interpretation, the additional bit would not mean much at the level of proteins.

Dark DNA and presumably also RNA codons are linked chains of 3 closed flux tubes serving as bits. Could this chain in the case of dark amino acids be replaced with a single closed flux tube obtained by two reconnections so that amino-acid becomes a basic unit?

2. Could the breaking of A-G symmetry (stop-trp pair and ile-met pair) have a topological meaning? Could the direction of the magnetic flux for the third flux tube of the dark codon coding for these pairs matter (this is not the only possibility that one can imagine)? Note that the 4 tetrahedral dark codons in the bioharmony model [L19, L8] contain at least one of these pairs.
3. What could topologically distinguish met as a starting codon and stop codons from the other codons? Could it be that met is not linked to the codons preceding it so that transcription would naturally start at it.

Could stop codons be unlinked to the codons following them so that the transcription would naturally stop at them? Or could the stop codons correspond to a single closed flux tube so that no RNA codon could be assigned to them?

4. Genes contain intronic parts and the splicing of RNA eliminates these parts after the transcription. Could the topology of DNA and RNA isolate intronic portions from those to be translated. Could the intronic portions correspond to a single flux tube linked to the rest of the gene both at the level of DNA and RNA. If so, the information about the decomposition of intron to RNA codons would be missing and the assignment of tRNA codons to the intronic portion would not be possible.

## 2.9 Could dark genes be dynamical?

In [L18] it was found that the earlier 1-1 correspondence between dark codons and ordinary genetic codons is unnecessarily strict and a modification of the earlier picture of the relation between dark and chemical genetic code and of the function of dark genetic code was considered.

1. Dark DNA (DDNA) strand is dynamical and has the ordinary DNA strand associated with it and dark gene state can be in resonant interaction with ordinary gene only when it corresponds to the ordinary gene. This applies also to DRNA, DtRNA and DAA (AA is for amino acids).

This would allow DDNA, DRNA, DtRNA and DAA to perform all kinds of information processing such as TQC by applying dark-dark resonance in quantum communications. The control of fundamental biomolecules by their dark counterparts by energy resonance would be only one particular function.

2. Most importantly, flux tubes magnetization direction could define qubit. If the additional qubit corresponds to nucleon isospin, it is not clear whether this is the case. One can also allow superpositions of the dark genes representing 6-qubit units. A generalization of quantum computation so that it would use 6-qubits units instead of a single qubit as a unit, is highly suggestive.
3. Genetic code could be also interpreted as an error code in which dark proteins correspond to logical 6-qubits and the DNA codons coding for the protein correspond to the physical qubits associated with the logical qubit.
4. The teleportation mechanism discussed in [L18] could make possible remote replication and remote transcription of DNA by sending the information about the ordinary DNA strand to the corresponding dark DNA strand by energy resonance. After that, the information would be teleported to a DNA strand in a ferromagnetic ground state at the receiver. After this, ordinary replication or transcription, which would also use the resonance mechanism, would take place.

### 3 Connection between dark nucleon code and bioharmony

The model of genetic code based on bioharmony has evolved through many sidetracks [K3] but the the version discussed in [L3, L8, L11] is roughly consistent with the original model and also gives a connection with the model of dark nucleon code.

#### 3.1 Bioharmony and resonance mechanism for dark photon communications

The faces of icosahedron and tetrahedron (and also octahedron appearing in the model of genetic code as icoso-tetrahedral tessellation of hyperbolic space  $H^3$  [L11]) are triangles. The proposal is that they somehow correspond to 3-chords made of dark photons, which in turn represent genetic codons.

Communications by dark 3-photons represent codons. 3N-photons represent in turn genes. The communications rely on cyclotron 3N-resonance so that the vertices of the faces of icoso-tetrahedron must contain charged particles coupling to a magnetic field. The magnetic field strengths at flux tubes associated with charged particles would determine the cyclotron frequencies.

Information is encoded to the frequency modulation of cyclotron frequencies. The chords serve as addresses much like in computer language LISP. If the modulations of 3N frequencies are identical and in synchrony, the outcome of the receiver consisting of 3N charged particles is a sequence of 3N-resonances giving rise to an 3N-pulse sequence. Nerve pulse patterns could emerge by this mechanism.

One can also consider 3N-signals for which only  $M < 3N$  modulations are identical and in synchrony. In this manner communications to subsets of the receiver are possible. For instance, some subset of codons of dark gene or dark protein can be selected as a receiver, possibly controlled. This selection could de-entangled the receiver to de-entangled coherent pieces.

There is a direct connection with empiria. Biophotons, whose origin remains poorly understood, can be identified as ordinary photons resulting from the decay of dark 3N-photons to ordinary photons.

The realization in terms of dark nucleons looks more plausible if also DtRNA and DAAs are realized in terms of icoso-tetrahedral picture. This is because the amino acids (AAs) are often neutral unlike DNA nucleotides, which are negatively charged. The dark charge assignable to the dark codon can be controlled by pion-like bonds with charges  $0, \pm 1$  so that it can be 3 units for DDNA and vanish for AAs. A natural proposal is that the pionic charge of the codon compensates

the charge of the AA and tRNA but the dark charge could be also dynamical and control the ionization of the biomolecule.

Large value of  $h_{eff}$  would make possible dark nuclear interactions in the scale of the dark codons (about codon size) allowing the transformation of dark protons to neutrons by dark variant of strong interactions (dark nuclear interactions are central in the TGD based model of "cold fusion" [L2, L7] having implications also for the stellar evolution [L4]). Dark codons would be analogous to dark nuclei tritium (pnn),  $^3\text{He}$  (ppn), and also ppp and nnn not realized as ordinary nuclei).

There are pairings of type DX-DY. The pairings DDNA-DRNA, DRNA- DtRNA and DtRNA-DAA induce the biochemical dynamics of transcription and translation. There are also pairings DX-X involving the transformation of dark 3-photon to ordinary 3-photon and occurring via energy-resonance but involving downwards scaling of wave-length. DDNA-DNA and DRNA-RNA unique DtRNA-tRNA pairing is 1-to-many and relates to the wobble phenomenon. The pairings between dark nucleon variants of biomolecules and corresponding dark 3N-photons make possible biocommunications and control.

### 3.2 Details of the bioharmony model

Consider now a more detailed bioharmony model of the genetic code based on the geometries of icosahedron and tetrahedron.

1. Icosahedron has 12 vertices and 20 faces, which are triangles. The idea is that the 12 vertices correspond to the notes of 12-note scale. Tetrahedron has 4 vertices and 4 faces and is self-dual whereas the dual of icosahedron is dodecahedron having 20 faces and 12 faces.

2. 12-note scale can be represented as a Hamiltonian cycle at an icosahedron going once through all vertices. The frequencies at the neighboring points as edges of a face in cycle relate by a frequency scaling of  $3/2$ : this gives rise to the Pythagorean variant of quint cycle.

Octave equivalence means the identification of frequencies differing by a multiple of octaves. Octave equivalence can be used to reduce all frequencies to a single octave. If the scaling is exactly  $3/2$  at all steps there is a slight-breaking of octave equivalence since  $(3/2)^{12}$  does not quite correspond to an integer number (7) of octaves. Pythagoras was well aware of this.

Given cycle assigns to faces 3-chords defining a harmony with 20 chords assignable to the faces of the icosahedron. For dodecahedron there is only single harmony with 12 chords and 20-note scale which could correspond to Eastern scales. For the tetrahedron the Hamiltonian cycle is unique.

3. Icosahedral Hamiltonian cycles can be classified by symmetries. The group  $Z_6$ ,  $Z_4$ , or  $Z_2$  (rotation by  $\pi$  or reflection) as a group of symmetries

The connection with the genetic code emerges in the following manner.

1. The natural idea is that the faces of the ico-tetrahedron correspond to both 3-chords and genetic DNA/RNA codons. If the orbits of faces could correspond to AAs (AAs), the DNA codon would code for AA AA if the corresponding face is at the orbit corresponding to AA.
2. One wants 64 DNAs:  $Z_6, Z_4$  ja  $Z_2$  cycle give rise to  $20+20+20=60$  DNAs codons. Tetrahedron gives the remaining 4 codons.
3. Does one obtain a correct number of AAs? Do the numbers of faces at the orbits correspond to numbers of DNAs coding for the corresponding AA?
  - (a)  $Z_6$  decomposes to 3 6-orbits and 1 2-orbit ( $3 \times 6 + 2 = 20$ ). There are 3 AAs coded by 6 DNAs. 2-orbit corresponds to AA coded by two DNAs.
  - (b)  $Z_4$  decomposes to 5 4-orbits. There are 5 AAs coded by 4 codons.

- (c)  $Z_2$  corresponds to 10 2-orbits predicting 10 AAs coded by 2 codons. There would be 11 2-orbits altogether. There are 9 AAs coded by 2 codons.

Some kind of symmetry breaking is present as in the case of dark nucleon code. 2 AA doublets must split to singlets. If (ile,ile,ile,met) coded by UAX could correspond to (ile,ile) and (met,met) such that (met,met) is split to (ile,met). In absence of symmetry breaking one would have 11 doublets as predicted.

4. There are also 4 tetrahedral codons.

There is (stop,stop) doublet (UAA, UAG) and (stop,trp) doublet (UGA,UGG). These doublets could correspond to the faces of the tetrahedron. Only one face would code for AA in the vertebrate code. Other faces would not have corresponding tRNA?

For bacterial codes, the situation can be different. Pyl and sec appear as exotic AAs. Could (UAA,UAG) code for (stop,pyl) and (UGA,UGG) for (sec,trp) instead of (stop,trp)? Orientation preserving rotations form a 12-element group having  $Z_2$  and  $Z_3$  as subgroups. For  $Z_2$  the orbits consist of 2 vertices and for  $Z_3$  of 3 vertices (face) and 1 vertex.  $Z_3$  symmetry could correspond to trp as singlet and vertebrate stop codons as triplet. For bacterial pyl and sec  $Z_2$  with symmetry breaking is suggestive.

### 3.3 Bioharmony, dark nucleon code, and icoso-tetrahedral code as a tessellation of $H^3$

Bioharmony model involves icosahedron and tetrahedron. This looks ugly unless there is some really deep reason for their emergence. One can also ask why not also octahedron having triangular faces.

Hyperbolic 3-space  $H^3$  has interpretations as a mass shell of Minkowski space  $M^4$  at the level of  $M^8$  and as light-cone proper-time constant surface at the level of  $H$ . The 4-surface  $X^4$  in  $M^8$  contains mass shells of  $M^4$  corresponding to the roots of the polynomial  $P$  defining  $X^4$ . Hence one expects that  $H^3$  plays a key role in quantum TGD both discretized momentum as defining a cognitive representation with momenta, which are algebraic integers associated with extension of rationals defined by  $P$ .  $H^3$  has infinite discrete subgroups of the Lorentz group analogous to discrete groups of translations in  $E^3$  as isometries and  $H^3$  allows an infinite number of tessellations (lattices). Perhaps the simplest tessellation is icoso-tetrahedral tessellation involving also octahedrons and thus all triangular Platonic solids. This tessellation could give rise to genetic code by induction of tessellation to 3-surfaces or lower-D objects such as linear biomolecules, and cell membranes [L11]. I do not however understand the mathematical details well enough but the following discussion is general.

Consider first the model for DDNA and DRNA allowing us to understand the connection between dark nucleon and dark photon realization of the genetic code physically.

1. The realization of DDNA/DRNA/DtRNA/DAA could correspond to a sequence of icosahedron-tetrahedron pairs at  $H^3$  contained by the 4-surface  $X^4 \subset M^8$  and its  $H$  images which is also  $H^3$ .
2. Each icoso-tetrahedron would contain a dark codon realized both as a face and dark nucleon triplet associated with it. The dark photon chord associated with the face must be the same as the codon defined by dark nucleon triplet. The dark nucleon triplets correspond to cyclotron frequency triplets, which in turn correspond to dark photon 3-chords associated with the Hamiltonian cycles.
3. The cyclotron frequencies are determined by magnetic fields at flux tubes so that Hamilton cycles must correspond to flux tube patterns. The simplest hypothesis is that the Hamilton cycle is a closed flux tube connecting all vertices of the icosahedron. Dark codon triplet corresponds to a face. It does have 1 or 2 flux tube edges if the corresponding chord contains 1 or two quints and otherwise no flux tube edges. Therefore cyclotron frequencies cannot be always associated with the edges of the triangle.

The simplest option is that the Hamiltonian flux tube following the vertex at the cycle defines the cyclotron frequency associated with the vertex. The harmony depends on the

orientation of the cycle and for 8-note scale roughly corresponds the transformation from major to minor. The variation of flux tube thickness implies frequency modulation crucial for communications.

The realization of the Hamilton cycle requires that the magnetic field strength along the cycle is scaled by factor  $3/2$  to give a Pythagorean quint cycle. For an evenly tempered quint scale the scaling is  $2^{5/12}$ .

4. An interesting question relates to the relation of DDNA strand and its conjugate. The change of the orientation of the Hamiltonian cycle changes the chord of the harmony. For the ordinary 8-note scale one can roughly say that major and minor chords are transformed to each other. The orientation reversal could correspond to time reversal. The fact that the orientations of two DNA strands are opposite suggests that DNA and conjugate DNA are related by the orientation reversal of the Hamiltonian cycle inducing the map  $G \rightarrow C$ ,  $U \rightarrow A$  at the level of DNA letters. The conjugation does not imply any obvious symmetry for the corresponding AAs as the inspection of the code table demonstrates.

How could the Hamiltonian cycle determine the DtRNA codons?

1. DRNA codons pair with 32 DtRNA codons and DtRNA codons pair with tRNA codons in 1-to-many manner. Therefore DRNA-DtRNA pairing could be universal and 2-1, although not in a codon-wise manner. This pairing should be the same for both bioharmony and dark nucleon triplets.
2. The pairing by 3-resonances requires that DtRNA icoso-tetrahedron contains the DRNA codons, which pair with DtRNA codon. There would be 2 DRNA codons in DtRNA icoso-tetrahedron for most DtRNA codons and 1 codon for DtRNA pairing with DAA corresponding to met and trp. The number 32 of DtRNA implies in the case of icoso-tetrahedral code that there are  $10+10-10=30$  icosahedral DtRNAs and only 2 tetrahedral DtRNAs so that two faces of tetrahedron cannot correspond to DtRNA codon so that corresponding DRNAs must serve as stop codons.

One of the DtRNAs could correspond to trp. The second one would correspond to a stop codon in the vertebrate code: either the DtRNA codon is not present at all or it does not pair with tRNA. TAG and TGA can code for pyl and sec in some bacterial versions of the code and in this case the corresponding dark DRNA codon would be represented at the DtRNA tetrahedron.

3. For bioharmony DDNA-DAA correspondence means that AAs correspond to orbits of the faces of icosahedron under the subgroup  $Z_6, Z_4$ , or  $Z_2$  which could correspond to reflection or to a rotation by  $\pi$ .

Since DRNA-DtRNA correspondence is 2-1 although not codon-wise, the natural first guess is that  $Z_2$  orbits of the faces define the DRNA codons at the DtRNA icosahedron so that it would contain 2 codons for most DtRNAs. At the DtRNA tetrahedron the only option is  $Z_1$  so there is a symmetry breaking.

If  $Z_2$  corresponds to a reflection, the orbit always contains 2 codons. If  $Z_2$  corresponds to a rotation by  $\pi$ , it might happen that the face invariant under  $\pi$  rotation and the orbit would consist of a single point. Could this explain why one has (ile,ile,ile,met) instead of (ile,ile) and (met,met)? The rotation axis should go through the invariant face and since the face is a triangle,  $\pi$  rotations lead out of the icosahedron. Therefore the answer is negative.

Ile-met problem deserves a separate discussion.

1. The pairing of  $Z_2$  related DDRNA faces with two different DtRNAs coding for ile and met rather than two mets means  $Z_2$  symmetry breaking at the level of bioharmony. Could the fact that AUG acts as a start codon relate to this? Could it be that both AUG and AUA cannot act as start codons? It is difficult to invent any reason for this.
2. The symmetry breaking could occur in DtRNA-DAA pairing and replace Dmet with Dile. Is it possible that the 3-chords for coding for ile and second met are nearly identical so

that the resonance mechanism selects ile instead of met? Could the situation be similar for the codons coding for (stop,stop) and (stop,trp) and cause the coding of pyl or sec in some situations? The scale for the quint cycle model with octave equivalence does not quite close. Could this have some role in the problem?

3. Since similar ambivalence occurs for stop codons assigned to the tetrahedral Hamiltonian cycle, one can look at the tetrahedral Hamiltonian cycle. In this case one has 3-quint cycle and a given edge of the cycle corresponds to a scaling by  $(3/2)^3$  so that 4 steps gives  $(3/2)^{12}$ , which is slightly more than 7 octaves. For the quint scale in Pythagorean sense, one obtains 4 notes in the same octave.

Exact octave equivalence corresponding to equally tempered scale in which half-note corresponds to frequency scaling  $2^{1/12}$ , implies that there is only one 3-chord  $CEG\sharp$ : this would explain why there are 3 stop codons in the vertebrate code!

The original guiding idea in the attempts to understand the fusion of icosahedral and tetrahedral codes was that the tetrahedron is effectively glued to the icosahedron along one face. This is consistent with the icosahedral quint cycle only if the common face contains no edges of the icosahedral cycle but contains tetrahedral flux edges with  $(3/2)^3$  scaling. This would give strong constraints on the common face.

If bacterial codes correspond to Pythagorean scale, there would be two different 3-chords since  $CEG\sharp$  and  $EG\sharp C$  are not quite the same. The reason is that the frequency ratios of chords are powers of  $3/2$ )<sup>12</sup>. This situation is completely exceptional.

In the quint scale there are small differences between the 4 chords. Could this explain why only one of these 3-chords codes for AA (trp) in vertebrate code and pyl or sec is coded instead of stop in bacterial codes? Amusingly, the chord  $CEG\sharp$  ends many finnish tangos and therefore acts like a stop codon!

Could bacteria have a perfect pitch and live in a Pythagorean world? Could the transition to multicellulars mean the emergence of an algebraic extension of rationals containing  $2^{1/12} \simeq 1.059$ , (which is considerably larger than to  $(3/2)^{12}/2^7 \simeq 1.0136$ )! Could people with perfect pitch have in their dark genome parts using Pythagorean scale or can they tune the magnetic flux tube radii to realize Pythagorean scale?

4. Could the ile-met problem have a similar solution? The chords associated with ile and met would differ by ascaling with  $(3/2)^3$  or  $(3/2)^6$  using octave equivalence. These chords are not quite the same: could it happen that the 3-chord associated with the second met is nearer to that for ile? These 3-chords do not contain quint scaling and should correspond to the special chords for which no edge belongs to a Hamiltonian cycle.

Also DtrRNA-DAA pairing is based on the 3-resonance.

1. DAA icosahedron must contain the DtrRNA codons pairing with DAA. This raises the question whether DDNAs could have a direct resonant coupling to DAAs. Could this pairing occur in DDNA-DAA occurring in transcription (<https://cutt.ly/QPP46St>) so that pieces of DDNA and DAA associated with an enzyme involved could pair with each other by 3N-resonance at DDA-DAA level? At the chemical level the base-AA interactions are extremely complex involving stereochemistry and formation of hydrogen bonds (<https://cutt.ly/RPP7p1M>) so that the reduction of these interactions to 3N-resonance would mean a huge simplification.
2. Could this resonance pairing serve as a universal mechanism of bio-catalysis and take place for various enzymes and ribozymes? One example are promoters and enhancers involved with the transcription. Enhancers and promoters induce a highly non-local process generating a chromosome loop in which two portions of DNA become parallel and near to each other and dark 3N-photons could explain the non-locality as an outcome of quantum coherence in long scales.
3. Why would DDNA-DAA pairing not occur? 3N-resonance relies on cyclotron frequencies and therefore on the magnetic field strength determined by the radii of the monopole flux

tubes. One explanation would be that the frequency scales of DAA and DDNA are slightly different. Could the attachment of DRNA to translation machinery scale the magnetic field strengths of the flux tubes and their cyclotron frequencies so that only dRNA-DtRNA and DtRNA-DAA couplings are possible.

## 4 Pythagorean number mysticism, music harmony, and genetic code

The discussion with Marko Manninen renewed my interest in the ideas of Pythagoras related to mystics and mathematics and its role in music.

### 4.1 Rational Platonica

Pythagoras believed that rationals are all that is needed for a Universe and for him the discovery of  $\sqrt{2}$  represented geometrically by the diagonal of a unit square was probably a shock.

It is interesting that in the TGD framework the rationals appear naturally. In its simplest form, Galois confinement [L14, L15] states that the total 4-momenta of physical states are Galois singlets invariant under Galois group permuting roots of a given polynomial (the notion generalizes if one considers functions in momentum space). This would allow only momenta with components, which are integers when a physical natural momentum unit is used. Platon would have been right in a certain sense!

However, Galois singlets would at fundamental level consist of quarks (in particular leptons and bosons would do so) having 4-momenta with components, which are algebraic integers in the extension of rationals defined by the polynomial defining the space-time region considered [L5, L6]. One could regard the algebraic integer valued momenta as virtual momenta characterizing the building bricks of physical states.

### 4.2 Special role of primes 2, 3 and 5

The number mysticism of Pythagoras involves the idea that the numbers 2 and 3 are very special. Using the language of modern number theory, one could say numbers 2 and 3 span a group with respect to multiplication consisting of numbers  $2^m 3^n$ , where  $m$  and  $n$  are integers. One could call this group  $B(2, 3)$ . If  $m$  and  $n$  are restricted to non-negative integers, the inverses do not exist and only a semigroup is obtained. This object could be called  $A(2, 3)$ .

If Pythagoras identified rational numbers as a kind of Platonica, this group might be said to define an important province of Platonica. A more general object would be ideal consisting of all integers proportional to, say,  $6 = 2 \times 3$  closed with respect to multiplication by any integer.

It should be noticed that any set  $(p_1, \dots, p_n)$  of primes and even integers defines a group with respect to multiplication as the group  $B(m_1, m_2, \dots, m_n)$  of integers. Especially interesting example is the group  $B(2, 3, 5)$  containing  $B(2, 3)$  and  $B(3/2)$ .

p-Adic length scale hypothesis states that powers of small primes near to prime define important p-adic length scales. Powers of 2 are of special importance in p-adic mass calculations [K2] but there exists also evidence for powers of 3 [I4, I5].

Decimal system is the decimal system used in everyday life and very often numerologists freely change the position of a decimal number and get results, which make sense only if the decimal system is in a special role. Could this be the case? If so, then the decimal system would not reflect only the fact that we have 10 fingers, and also the algebras  $B(2, 5)$  and  $B(10)$  could be special.

There are some indications that this might be the case.

1. The faces of icosahedron *resp.* dodecahedron are triangles *resp.* pentagons so that numbers of 3 and 5 are natural.
2. DNA is a helical structure with a twist angle  $2\pi/10$  between to codons so that 10 codons make a  $6\pi$  twist and define length scale 10 nm which is the p-adic length scale associated with Gaussian Mersenne prime  $M_{G,151} = (1 + i)^{151} - 1$ , one of the 4 Gaussian Mersennes

defining p-adic length scales in the range 10 nm, 2.5  $\mu\text{m}$ . These scales are a number theoretic miracle. Numbers 2,3, and 5 relate to the geometry of DNA.

### 4.3 Pythagorean scale

Pythagoras also studied music and introduced the notion of Pythagorean scale for which the frequencies of notes are in rational ratios. A standard manner to realized this scale is by quint cycle, which means that one forms the multiples  $(3/2)^n f_0$  of fundamental frequency  $f_0$  and identifies them by octave equivalence with a frequency in the basic octave  $[f_0, 2f_0]$ . The quint cycle appears very often in jazz.

For  $n = 12$  the frequency obtained is almost a full number of octaves but quite not. This imperfectness of Platonica troubled Pythagoras a lot. In an equal tempered scale one introduces powers  $2^{m/2^n} f_0$  and avoids this problem. This means replacement of rationals by its algebraic extension generated by  $2^{1/12}$ .

Obviously, the Pythagorean scale is very natural in the framework of group  $B(3/2)$ . Pythagoras also had ideas about the relationship of music scale and Platonic solids.

### 4.4 Pythagorean scale and genetic code

In the TGD framework, the idea about a possible connection between music and Platonic solids inspired the proposal about realization of the 12-note scale as a Hamiltonian cycle at icosahedron. The Hamiltonian cycle is a closed curve connecting only neighboring points of the icosahedron and going through all its 12 vertices. There are quite a large number of icosahedral cycles and they assign to the 20 triangles of icosahedron 3-chords proposed to define icosahedral harmony with 20 chords. The non-chaotic icosahedral cycles have symmetry groups  $Z_6, Z_4$ , and  $Z_2$ , which can act as a rotation or reflection.

The big surprise was that the model of icosahedral harmony leads to a model of genetic code. The code would involve a fusion of 3 different icosahedral harmonies with symmetry groups  $Z_6, Z_4$ , and  $Z_2$  giving 60 codons plus tetrahedral code giving 4 codons. The counterparts of amino-acids would correspond to the orbits of these symmetry groups: 3 orbits with 6 triangles and 1 with 2 triangles as orbits for  $Z_6$ , 4 orbits with 4 triangles for  $Z_4$  and 10 orbits with 2 triangles for  $Z_2$ . The number of triangles at the orbit is the number of DNA codons. Tetrahedron would give the missing 4 codons and stop codons and one missing amino acid.

For a given choice of the 3 Hamiltonian cycles, the realization would be in terms of 3-chords of light defining harmony for a music of light (and possibly also sound). Since music expresses and generates emotions, the proposal was that this realization of the genetic code expresses emotions already at the molecular level and that emotional intelligence corresponds to this realization whereas bit intelligence would correspond to the interpretation of codons as 6-bit sequences.

It should be mentioned that Hamiltonian cycles are solutions to the travelling salesman problem at the icosahedron: cities would correspond to the vertices. In the case of dodecahedron, which is dual of icosahedron, there is only one Hamiltonian cycle so that the harmony is now unique. If this corresponds to harmony, the first guess is that there would be a 20-note scale and 12 5-chords.

### 4.5 What about dodecahedral harmony and analog of genetic code?

Could also dodecahedron define a bioharmony and an analog of genetic code?

1. The first guess is that dodecahedral harmony has 20 notes per octave and perhaps corresponds to the scale defined by micro-octaves used in Eastern music. There would be 12 5-chords and the harmony would be unique. There would exist only a single emotional mood, a kind of enlightened state.
2. Since the harmony is unique, and there are no other Platonic solids with pentagons as faces. The analog of genetic code should correspond to dodecahedral harmony. The 5-chords would define 12 analogs of DNA codons.

The dodecahedral cycle  $Z_3$  acts as a symmetry group (<https://jrh794.wordpress.com/2021/04/01/the-original-hamiltonian-cycle-continued/>). This means that there are

4 orbits of  $Z_3$  with 3 codons at each and they would correspond to 4 different analogs of amino-acids.

3. Could one consider instead of an icosahedral quint cycle with scaling  $3/2$  replaced with scaling  $5/2$ ? The tempered system would use powers of  $2^{1/20}$  to generate a 20-note scale. A single step along Hamilton's cycle connecting neighboring vertices of dodecahedron would correspond to a scaling  $f \rightarrow 5/2f$  plus octave equivalence.

By octave equivalence the scaling by  $5/2$  would correspond to a transition from say C to a note between  $E_b$  and  $E$ . The microintervals between this note and either  $E_b$  or  $E$  appears in blues, jazz etc as a blue note. This interval is between minor and major.

4. One can test this. The cold shower is that  $(5/2)^{20}$  is not near to a power of 2. However, one has  $(5/2)^{19}/2^{25} = 1.084$  (for the quint cycle in the icosahedral case the deviation that Platon was worried of, is about 1 percent). As if one had a 19-note scale. A completely analogous situation is encountered with bio-harmony. The scales assigned to  $Z_6, Z_4$ , and  $Z_2$  give rise to 19 amino-acids as orbits of these groups. One amino-acid is missing and the tetrahedral code gives this amino-acid plus 3 stop codons [L19].

The icoso-dodecahedral duality suggests that the scale should consist of 19 notes only. Note however that for an equal tempered variant of the scale one does not have this problem.

5. Dodecahedral code predicts 4 analogs of amino acids. Could these "amino acids" correspond to the 4 DNA codons? 3 dodecahedral codons would be needed to code for a single genetic codon.

Could the dodecahedral codons, which correspond to 5-chords, be realized as dark 5-photons and sequences of dark 5-protons. One should check whether the states of 5 dark protons could give rise to 12 dark dodecahedral codons and whether something analogous to 12 dark RNA codons, dark tRNA codons, and 4 dark amino acids could emerge.

For the dodecahedral bioharmony, 5-chords would label the codons and they would serve as addresses based on communications relying on cyclotron resonance. Icosahedral harmony would control codons and dodecahedral harmony would code for their letters so that the codes would appear in different scales.

This speculation raises some questions. One can argue that in the transcription and replication the control of both codons and individual letters is important. This suggests that both realizations are needed for both DNA strands and correspond to different control scales. This would be true for the transcribed DNA, at least.

One can however consider alternatives. For instance, could the passive DNA strand correspond to a dodecahedral realization at the level of letters and the active strand to the icosahedral realization at the level of codons. Or could "junk" DNA and introns in promoter regions correspond to the dodecahedral realization with dark dodecahedral DNA controlling single letters.

## REFERENCES

### Biology

- [I1] Wobble base pair. Available at: [https://en.wikipedia.org/wiki/Wobble\\_base\\_pair](https://en.wikipedia.org/wiki/Wobble_base_pair).
- [I2] The Fourth Phase of Water: Dr. Gerald Pollack at TEDxGuelphU, 2014. Available at: <https://www.youtube.com/watch?v=i-T7tCMUDXU>.
- [I3] Pollack G. *Cells, Gels and the Engines of Life*. Ebner and Sons, 2000. Available at: <https://www.cellsandgels.com/>.
- [I4] Fiaxat JD. A hypothesis on the rhythm of becoming. *World Futures*, 36:31–36, 1993.
- [I5] Fiaxat JD. The hidden rhythm of evolution, 2014. Available at: [https://byebyedarwin.blogspot.fi/p/english-version\\_01.html](https://byebyedarwin.blogspot.fi/p/english-version_01.html).

- [I6] Zhao Q Pollack GH, Figueroa X. Molecules, water, and radiant energy: new clues for the origin of life. *Int J Mol Sci*, 10:1419–1429, 2009. Available at: <https://tinyurl.com/ntkfhlc>.
- [I7] Pollack GH Zheng J-M. Long-range forces extending from polymer-gel surfaces. *Phys Rev E*, 68:031408–, 2003. Available at: <https://tinyurl.com/ntkfhlc>.

## Books related to TGD

- [K1] Pitkänen M. Genes and Memes. In *Genes and Memes: Part I*. Available at: <https://tgdtheory.fi/pdfpool/genememec.pdf>, 2006.
- [K2] Pitkänen M. *p-Adic length Scale Hypothesis*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/padphys.html>., 2013.
- [K3] Pitkänen M. Geometric Theory of Bio-Harmony. In *TGD based view about living matter and remote mental interactions: Part II*. Available at: <https://tgdtheory.fi/pdfpool/harmonytheory.pdf>, 2018.
- [K4] Pitkänen M. About the Correspondence of Dark Nuclear Genetic Code and Ordinary Genetic Code. In *Genes and Memes: Part II*. Available at: <https://tgdtheory.fi/pdfpool/codedarkcode.pdf>, 2019.
- [K5] Pitkänen M. Nuclear String Hypothesis. In *Hyper-finite Factors and Dark Matter Hierarchy: Part II*. Available at: <https://tgdtheory.fi/pdfpool/nuclstring.pdf>, 2019.
- [K6] Pitkänen M. TGD and Nuclear Physics. In *Hyper-finite Factors and Dark Matter Hierarchy: Part II*. Available at: <https://tgdtheory.fi/pdfpool/padnucl.pdf>, 2019.
- [K7] Pitkänen M. Three new physics realizations of the genetic code and the role of dark matter in bio-systems. In *Genes and Memes: Part II*. Available at: <https://tgdtheory.fi/pdfpool/dnatqccodes.pdf>, 2019.
- [K8] Pitkänen M. and Rastmanesh R. ZEO, adelic physics, and genes. In *Genes and Memes: Part II*. Available at: <https://tgdtheory.fi/pdfpool/zeogenes.pdf>, 2020.

## Articles about TGD

- [L1] Pitkänen M. Pollack's Findings about Fourth phase of Water : TGD View. Available at: [https://tgdtheory.fi/public\\_html/articles/PollackYoutube.pdf](https://tgdtheory.fi/public_html/articles/PollackYoutube.pdf)., 2014.
- [L2] Pitkänen M. Cold fusion, low energy nuclear reactions, or dark nuclear synthesis? Available at: [https://tgdtheory.fi/public\\_html/articles/krivit.pdf](https://tgdtheory.fi/public_html/articles/krivit.pdf)., 2017.
- [L3] Pitkänen M. An overall view about models of genetic code and bio-harmony. Available at: [https://tgdtheory.fi/public\\_html/articles/gcharm.pdf](https://tgdtheory.fi/public_html/articles/gcharm.pdf)., 2019.
- [L4] Pitkänen M. Solar Metallicity Problem from TGD Perspective. Available at: [https://tgdtheory.fi/public\\_html/articles/darkcore.pdf](https://tgdtheory.fi/public_html/articles/darkcore.pdf)., 2019.
- [L5] Pitkänen M. A critical re-examination of  $M^8 - H$  duality hypothesis: part I. Available at: [https://tgdtheory.fi/public\\_html/articles/M8H1.pdf](https://tgdtheory.fi/public_html/articles/M8H1.pdf)., 2020.
- [L6] Pitkänen M. A critical re-examination of  $M^8 - H$  duality hypothesis: part II. Available at: [https://tgdtheory.fi/public\\_html/articles/M8H2.pdf](https://tgdtheory.fi/public_html/articles/M8H2.pdf)., 2020.
- [L7] Pitkänen M. Could TGD provide new solutions to the energy problem? Available at: [https://tgdtheory.fi/public\\_html/articles/proposal.pdf](https://tgdtheory.fi/public_html/articles/proposal.pdf)., 2020.
- [L8] Pitkänen M. How to compose beautiful music of light in bio-harmony? [https://tgdtheory.fi/public\\_html/articles/bioharmony2020.pdf](https://tgdtheory.fi/public_html/articles/bioharmony2020.pdf)., 2020.

- [L9] Pitkänen M. The dynamics of SSFRs as quantum measurement cascades in the group algebra of Galois group. Available at: [https://tgdtheory.fi/public\\_html/articles/SSFRGalois.pdf](https://tgdtheory.fi/public_html/articles/SSFRGalois.pdf), 2020.
- [L10] Pitkänen M. Galois code and genes. [https://tgdtheory.fi/public\\_html/articles/Galoiscode.pdf](https://tgdtheory.fi/public_html/articles/Galoiscode.pdf), 2021.
- [L11] Pitkänen M. Is genetic code part of fundamental physics in TGD framework? Available at: [https://tgdtheory.fi/public\\_html/articles/TIH.pdf](https://tgdtheory.fi/public_html/articles/TIH.pdf), 2021.
- [L12] Pitkänen M. TGD as it is towards the end of 2021. [https://tgdtheory.fi/public\\_html/articles/TGD2021.pdf](https://tgdtheory.fi/public_html/articles/TGD2021.pdf), 2021.
- [L13] Pitkänen M. The recent view about TGD and applications to condensed matter. [https://tgdtheory.fi/public\\_html/articles/TGDcondmat.pdf](https://tgdtheory.fi/public_html/articles/TGDcondmat.pdf), 2021.
- [L14] Pitkänen M. About TGD counterparts of twistor amplitudes: part I. [https://tgdtheory.fi/public\\_html/articles/twisttgd1.pdf](https://tgdtheory.fi/public_html/articles/twisttgd1.pdf), 2022.
- [L15] Pitkänen M. About TGD counterparts of twistor amplitudes: part II. [https://tgdtheory.fi/public\\_html/articles/twisttgd2.pdf](https://tgdtheory.fi/public_html/articles/twisttgd2.pdf), 2022.
- [L16] Pitkänen M. Comparison of Orch-OR hypothesis with the TGD point of view. [https://tgdtheory.fi/public\\_html/articles/penrose.pdf](https://tgdtheory.fi/public_html/articles/penrose.pdf), 2022.
- [L17] Pitkänen M. How animals without brain can behave as if they had brain. [https://tgdtheory.fi/public\\_html/articles/precns.pdf](https://tgdtheory.fi/public_html/articles/precns.pdf), 2022.
- [L18] Pitkänen M. Quantum biological teleportation using multiple 6-qubits. [https://tgdtheory.fi/public\\_html/articles/fluteteleport.pdf](https://tgdtheory.fi/public_html/articles/fluteteleport.pdf), 2022.
- [L19] Pitkänen M. The realization of genetic code in terms of dark nucleon and dark photon triplets. [https://tgdtheory.fi/public\\_html/articles/darkcode.pdf](https://tgdtheory.fi/public_html/articles/darkcode.pdf), 2022.