

# Introduction to "Physics in Many-Sheeted Space-Time: Part II"

M. Pitkänen,

July 22, 2024

Email: [matpitka6@gmail.com](mailto:matpitka6@gmail.com).

[http://tgdtheory.com/public\\_html/](http://tgdtheory.com/public_html/).

Postal address: Rinnekatu 2-4 A 8, 03620, Karkkila, Finland. ORCID: 0000-0002-8051-4364.

## Contents

<b>1</b>	<b>Basic Ideas of Topological Geometrodynamics (TGD)</b>	<b>3</b>
1.1	Geometric Vision Very Briefly . . . . .	3
1.2	Two Visions About TGD as Geometrization of Physics and Their Fusion . . . . .	6
1.2.1	TGD as a Poincare Invariant Theory of Gravitation . . . . .	6
1.2.2	TGD as a Generalization of the Hadronic String Model . . . . .	7
1.2.3	Fusion of the Two Approaches via a Generalization of the Space-Time Concept . . . . .	7
1.3	Basic Objections . . . . .	8
1.3.1	Topological Field Quantization . . . . .	9
1.4	Quantum TGD as Spinor Geometry of World of Classical Worlds . . . . .	9
1.4.1	World of Classical Worlds . . . . .	9
1.4.2	Identification of Kähler function . . . . .	10
1.4.3	WCW spinor fields . . . . .	10
1.4.4	The role of modified Dirac action . . . . .	11
1.5	Construction of scattering amplitudes . . . . .	11
1.5.1	Reduction of particle reactions to space-time topology . . . . .	11
1.5.2	Construction of the counterparts of S-matrices . . . . .	12
1.5.3	The notion of M-matrix . . . . .	12
1.6	TGD as a generalized number theory . . . . .	13
1.6.1	The Threads in the Development of Quantum TGD . . . . .	13
1.6.2	Number theoretic vision very briefly . . . . .	14
1.6.3	p-Adic TGD and fusion of real and p-adic physics to single coherent whole . . . . .	15
1.6.4	Infinite primes . . . . .	17
1.7	An explicit formula for $M^8 - H$ duality . . . . .	17
1.7.1	Holography in $H$ . . . . .	17
1.7.2	Number theoretic holography in $M_c^8$ . . . . .	17

1.7.3	Can one find an explicit formula for $M^8 - H$ duality? . . . . .	18
1.7.4	What could the number theoretic holography mean physically? . . . . .	19
1.7.5	Twistor lift of the holography . . . . .	19
1.8	Hierarchy of Planck Constants and Dark Matter Hierarchy . . . . .	20
1.8.1	Dark Matter as Large $\hbar$ Phases . . . . .	20
1.8.2	Hierarchy of Planck Constants from the Anomalies of Neuroscience and Biology . . . . .	21
1.8.3	Dark Matter as a Source of Long Ranged Weak and Color Fields . . . . .	22
1.9	Twistors in TGD and connection with Veneziano duality . . . . .	22
1.9.1	Twistor lift at space-time level . . . . .	22
1.9.2	Twistor lift at the level of scattering amplitudes and connection with Veneziano duality . . . . .	23
<b>2</b>	<b>Bird's Eye of View about the Topics of "Physics in Many-sheeted Space-time: Part II"</b>	<b>26</b>
<b>3</b>	<b>Sources</b>	<b>27</b>
3.1	PART I: MANY-SHEETED COSMOLOGY . . . . .	27
3.1.1	Cosmic Strings . . . . .	27
3.1.2	TGD and Cosmology . . . . .	30
3.1.3	More about TGD and Cosmology . . . . .	32
3.1.4	Breaking of $CP$ , $P$ , and $T$ in cosmological scales in TGD Universe . . . . .	32
3.1.5	Comparing Electric Universe hypothesis and TGD . . . . .	33
3.2	PART II: MANY-SHEETED ASTROPHYSICS . . . . .	33
3.2.1	TGD and Astrophysics . . . . .	33
3.2.2	Quantum Astrophysics . . . . .	33
3.2.3	TGD View about Quasars . . . . .	35
3.2.4	Cosmic string model for the formation of galaxies and stars . . . . .	36
3.2.5	Could solar system be modelled as a miniature version of spiral galaxy? . . . . .	36
3.2.6	TGD View of the Engine Powering Jets from Active Galactic Nuclei . . . . .	37
3.2.7	Solar Metallicity Problem from TGD Perspective . . . . .	37
3.2.8	Solar Surprise . . . . .	39
3.2.9	LIGO and TGD . . . . .	39
3.2.10	About the Recent TGD Based View Concerning Cosmology and Astrophysics . . . . .	39

# 1 Basic Ideas of Topological Geometrodynamics (TGD)

Standard model describes rather successfully both electroweak and strong interactions but sees them as totally separate and contains a large number of parameters which it is not able to predict. For about four decades ago unified theories known as Grand Unified Theories (GUTs) trying to understand electroweak interactions and strong interactions as aspects of the same fundamental gauge interaction assignable to a larger symmetry group emerged. Later superstring models trying to unify even gravitation and strong and weak interactions emerged. The shortcomings of both GUTs and superstring models are now well-known. If TGD - whose basic idea emerged towards the end of 1977 - would emerge now it would be seen as an attempt to solve the difficulties of these approaches to unification.

The basic physical picture behind the geometric vision of TGD corresponds to a fusion of two rather disparate approaches: namely TGD as a Poincare invariant theory of gravitation and TGD as a generalization of the old-fashioned string model. After 1995 number theoretic vision started to develop and was initiated by the success of mass calculations based on p-adic thermodynamics. Number theoretic vision involves all number fields and is complementary to the geometric vision: one can say that this duality is analogous to momentum-position duality of wave mechanics. TGD can be also regarded as topological quantum theory in a very general sense as already the attribute "Topological" in "TGD" makes clear. Space-time surfaces as minimal surfaces can be regarded as representatives of homology equivalence classes and p-adic topologies generalize the notion of local topology and apply to the description of correlates of cognition.

## 1.1 Geometric Vision Very Briefly

*T(opological) G(eometro)D(ynamics)* is one of the many attempts to find a unified description of basic interactions. The development of the basic ideas of TGD to a relatively stable form took time of about half decade [K1].

The basic vision and its relationship to existing theories is now rather well understood.

1. Space-times are representable as 4-surfaces in the 8-dimensional embedding space  $H = M^4 \times CP_2$ , where  $M^4$  is 4-dimensional (4-D) Minkowski space and  $CP_2$  is 4-D complex projective space (see Appendix).
2. Induction procedure (a standard procedure in fiber bundle theory, see Appendix) allows to geometrize various fields. Space-time metric characterizing gravitational fields corresponds to the induced metric obtained by projecting the metric tensor of  $H$  to the space-time surface. Electroweak gauge potentials are identified as projections of the components of  $CP_2$  spinor connection to the space-time surface, and color gauge potentials as projections of  $CP_2$  Killing vector fields representing color symmetries. Also spinor structure can be induced: induced spinor gamma matrices are projections of gamma matrices of  $H$  and induced spinor fields just  $H$  spinor fields restricted to space-time surface. Spinor connection is also projected. The interpretation is that distances are measured in embedding space metric and parallel translation using spinor connection of embedding space.

Twistor lift of TGD means that one can lift space-time surfaces in  $H$  to 6-D surfaces a analogs of twistor space of space-time surface in the Cartesian product of the twistor spaces of  $M^4$  and  $CP_2$ , which are the only 4-manifolds allowing twistor space with Kähler structure [A7]. The twistor structure would be induced in some sense, and should coincide with that associated with the induced metric. Clearly, the 2-spheres defining the fibers of twistor spaces of  $M^4$  and  $CP_2$  must allow identification: this 2-sphere defines the  $S^2$  fiber of the twistor space of the space-time surface. This poses a constraint on the embedding of the twistor space of space-time surfaces as sub-manifold in the Cartesian product of twistor spaces. The existence of Kähler structure allows to lift 4-D Kähler action to its 6-D counterparts and the 6-D counterpart of twistor space is obtained by its dimensional reduction so that one obtains a sphere bundle. This makes possible twistorialization for all space-time surfaces: in general relativity the general metric does not allow this.

3. A geometrization of quantum numbers is achieved. The isometry group of the geometry of  $CP_2$  codes for the color gauge symmetries of strong interactions. Vierbein group codes

for electroweak symmetries, and explains their breaking in terms of  $CP_2$  geometry so that standard model gauge group results. There are also important deviations from the standard model: color quantum numbers are not spin-like but analogous to orbital angular momentum: this difference is expected to be seen only in  $CP_2$  scale. In contrast to GUTs, quark and lepton numbers are separately conserved and family replication has a topological explanation in terms of topology of the partonic 2-surface carrying fermionic quantum numbers.

$M^4$  and  $CP_2$  are unique choices for many other reasons. For instance, they are the unique 4-D space-times allowing twistor space with Kähler structure.  $M^4$  light-cone boundary allows a huge extension of 2-D conformal symmetries.  $M^4$  and  $CP_2$  allow quaternionic structures. Therefore standard model symmetries have number theoretic meaning.

4. Induced gauge potentials are expressible in terms of embedding space coordinates and their gradients and general coordinate invariance implies that there are only 4 field-like variables locally. Situation is thus extremely simple mathematically. The objection is that one loses linear superposition of fields. The resolution of the problem comes from the generalization of the concepts of particle and space-time.

Space-time surfaces can be also particle like having thus finite size. In particular, space-time regions with Euclidian signature of the induced metric (temporal and spatial dimensions in the same role) emerge and have interpretation as lines of generalized Feynman diagrams. Particles in space-time can be identified as a topological inhomogeneities in background space-time surface which looks like the space-time of general relativity in long length scales.

One ends up with a generalization of space-time surface to many-sheeted space-time with space-time sheets having extremely small distances of about  $10^4$  Planck lengths ( $CP_2$  size). As one adds a particle to this kind of structure, it touches various space-time sheets and thus interacts with the associated classical fields. Their effects superpose linearly in good approximation and linear superposition of fields is replaced with that for their effects.

This resolves the basic objection. It also leads to the understanding of how the space-time of general relativity and quantum field theories emerges from TGD space-time as effective space-time when the sheets of many-sheeted space-time are lumped together to form a region of Minkowski space with metric replaced with a metric identified as the sum of empty Minkowski metric and deviations of the metrics of sheets from empty Minkowski metric. Gauge potentials are identified as sums of the induced gauge potentials. TGD is therefore a microscopic theory from which the standard model and general relativity follow as a topological simplification, however forcing a dramatic increase of the number of fundamental field variables.

5. A further objection is that classical weak fields identified as induced gauge fields are long ranged and should cause large parity breaking effects due to weak interactions. These effects are indeed observed but only in living matter. The basic problem is that one has long ranged classical electroweak gauge fields. The resolution of the problem is that the quantum averages of induced weak and color gauge fields vanish due to the fact that color rotations affect both space-time surfaces and induced weak and color fields. Only the averages of electromagnetic fields are nonvanishing. The correlations functions for weak fields are nonvanishing below Compton lengths of weak bosons. In living matter large values of effective Planck constant labelling phases of ordinary matter identified as dark matter make possible long ranged weak fields and color fields.
6. General coordinate invariance requires holography so that space-time surfaces are analogous to Bohr orbits for particles identified as 3-surfaces. Bohr orbit property would be naturally realized by a 4-D generalization of holomorphy of string world sheets and implies that the space-time surfaces are minimal surfaces apart from singularities. This holds true for any action as long as it is general coordinate invariant and constructible in terms of the induced geometry. String world sheets and light-like orbits of partonic 2-surfaces correspond to singularities at which the minimal surface property of the space-time surfaces realizing the preferred extremal property fails. Preferred extremals are not completely deterministic, which implies what I call zero energy ontology (ZEO) meaning that the Bohr orbits are the fundamental objects. This leads to a solution of the basic paradox of quantum measurement

theory. Also the mathematically ill-defined path integral disappears and leaves only the well-defined functional integral over the Bohr orbits.

7. A string model-like picture emerges from TGD and one ends up with a rather concrete view about the topological counterpart of Feynman diagrammatics. The natural stringy action would be given by the string world sheet area, which is present only in the space-time regions with Minkowskian signature. Gravitational constant could be present as a fundamental constant in string action and the ratio  $\hbar/G/R^2$  would be determined by quantum criticality conditions. The hierarchy of Planck constants  $\hbar_{eff}/\hbar = n$  assigned to dark matter in TGD framework would allow to circumvent the objection that only objects of length of order Planck length are possible since string tension given by  $T = 1/\hbar_{eff}G$  apart from numerical factor could be arbitrary small. This would make possible gravitational bound states as partonic 2-surfaces as structures connected by strings and solve the basic problem of superstring theories. This option allows the natural interpretation of  $M^4$  type vacuum extremals with  $CP_2$  projection, which is Lagrange manifold as good approximations for space-time sheets at macroscopic length scales. String area does not contribute to the Kähler function at all.

Whether induced spinor fields associated with Kähler-Dirac action and de-localized inside the entire space-time surface should be allowed remains an open question: super-conformal symmetry strongly suggests their presence. A possible interpretation for the corresponding spinor modes could be in terms of dark matter, sparticles, and hierarchy of Planck constants.

It is perhaps useful to make clear what TGD is not and also what new TGD can give to physics.

1. TGD is *not* just General Relativity made concrete by using embeddings: the 4-surface property is absolutely essential for unifying standard model physics with gravitation and to circumvent the incurable conceptual problems of General Relativity. The many-sheeted space-time of TGD gives rise only at the macroscopic limit to GRT space-time as a slightly curved Minkowski space. TGD is *not* a Kaluza-Klein theory although color gauge potentials are analogous to gauge potentials in these theories.

TGD space-time is 4-D and its dimension is due to completely unique conformal properties of light-cone boundary and 3-D light-like surfaces implying enormous extension of the ordinary conformal symmetries. Light-like 3-surfaces represent orbits of partonic 2-surfaces and carry fundamental fermions at 1-D boundaries of string world sheets. TGD is *not* obtained by performing Poincare gauging of space-time to introduce gravitation and is plagued by profound conceptual problems.

2. TGD is *not* a particular string model although string world sheets emerge in TGD very naturally as loci for spinor modes: their 2-dimensionality makes among other things possible quantum deformation of quantization known to be physically realized in condensed matter, and conjectured in TGD framework to be crucial for understanding the notion of finite measurement resolution. Hierarchy of objects of dimension up to 4 emerge from TGD: this obviously means analogy with branes of super-string models.

TGD is *not* one more item in the collection of string models of quantum gravitation relying on Planck length mystics. Dark matter becomes an essential element of quantum gravitation and quantum coherence in astrophysical scales is predicted just from the assumption that strings connecting partonic 2-surfaces are responsible for gravitational bound states.

TGD is *not* a particular string model although AdS/CFT duality of super-string models generalizes due to the huge extension of conformal symmetries and by the identification of WCW gamma matrices as Noether super-charges of super-symplectic algebra having a natural conformal structure.

3. TGD is *not* a gauge theory. In TGD framework the counterparts of also ordinary gauge symmetries are assigned to super-symplectic algebra (and its Yangian [A1] [B4, B2, B3]), which is a generalization of Kac-Moody algebras rather than gauge algebra and suffers a fractal hierarchy of symmetry breakings defining hierarchy of criticalities. TGD is *not* one more quantum field theory like structure based on path integral formalism: path integral

is replaced with functional integral over 3-surfaces, and the notion of classical space-time becomes an exact part of the theory. Quantum theory becomes formally a purely classical theory of WCW spinor fields: only state function reduction is something genuinely quantal.

4. TGD view about spinor fields is *not* the standard one. Spinor fields appear at three levels. Spinor modes of the embedding space are analogs of spinor modes characterizing incoming and outgoing states in quantum field theories. Induced second quantized spinor fields at space-time level are analogs of stringy spinor fields. Their modes are localized by the well-definedness of electro-magnetic charge and by number theoretic arguments at string world sheets. Kähler-Dirac action is fixed by supersymmetry implying that ordinary gamma matrices are replaced by what I call Kähler-Dirac gamma matrices - this something new. WCW spinor fields, which are classical in the sense that they are not second quantized, serve as analogs of fields of string field theory and imply a geometrization of quantum theory.
5. TGD is in some sense an extremely conservative geometrization of entire quantum physics: *no* additional structures such as gauge fields as independent dynamical degrees of freedom are introduced: Kähler geometry and associated spinor structure are enough. "Topological" in TGD should not be understood as an attempt to reduce physics to torsion (see for instance [B1]) or something similar. Rather, TGD space-time is topologically non-trivial in all scales and even the visible structures of the everyday world represent non-trivial topology of space-time in the TGD Universe.
6. Twistor space - or rather, a generalization of twistor approach replacing masslessness in 4-D sense with masslessness in 8-D sense and thus allowing description of also massive particles - emerged originally as a technical tool, and its Kähler structure is possible only for  $H = M^4 \times CP_2$ . It however turned out that much more than a technical tool is in question. What is genuinely new is the infinite-dimensional character of the Kähler geometry making it highly unique, and its generalization to p-adic number fields to describe correlates of cognition. Also the hierarchy of Planck constants  $h_{eff} = n \times h$  reduces to the quantum criticality of the TGD Universe and p-adic length scales and Zero Energy Ontology represent something genuinely new.

The great challenge is to construct a mathematical theory around these physically very attractive ideas and I have devoted the last 45 years to the realization of this dream and this has resulted in 26 online books about TGD and nine online books about TGD inspired theory of consciousness and of quantum biology.

A collection of 30 online books is now (August 2023) under preparation. The goal is to minimize overlap between the topics of the books and make the focus of a given book sharper.

## 1.2 Two Visions About TGD as Geometrization of Physics and Their Fusion

As already mentioned, TGD as a geometrization of physics can be interpreted both as a modification of general relativity and generalization of string models.

### 1.2.1 TGD as a Poincare Invariant Theory of Gravitation

The first approach was born as an attempt to construct a Poincare invariant theory of gravitation. Space-time, rather than being an abstract manifold endowed with a pseudo-Riemannian structure, is regarded as a surface in the 8-dimensional space  $H = M^4 \times CP_2$ , where  $M^4$  denotes Minkowski space and  $CP_2 = SU(3)/U(2)$  is the complex projective space of two complex dimensions [A3, A6, A2, A5].

The identification of the space-time as a sub-manifold [A4, A9] of  $M^4 \times CP_2$  leads to an exact Poincare invariance and solves the conceptual difficulties related to the definition of the energy-momentum in General Relativity.

It soon however turned out that sub-manifold geometry, being considerably richer in structure than the abstract manifold geometry, leads to a geometrization of all basic interactions. First,

the geometrization of the elementary particle quantum numbers is achieved. The geometry of  $CP_2$  explains electro-weak and color quantum numbers. The different H-chiralities of  $H$ -spinors correspond to the conserved baryon and lepton numbers. Secondly, the geometrization of the field concept results. The projections of the  $CP_2$  spinor connection, Killing vector fields of  $CP_2$  and of  $H$ -metric to four-surface define classical electro-weak, color gauge fields and metric in  $X^4$ .

The choice of  $H$  is unique from the condition that TGD has standard model symmetries. Also number theoretical vision selects  $H = M^4 \times CP_2$  uniquely.  $M^4$  and  $CP_2$  are also unique spaces allowing twistor space with Kähler structure.

### 1.2.2 TGD as a Generalization of the Hadronic String Model

The second approach was based on the generalization of the mesonic string model describing mesons as strings with quarks attached to the ends of the string. In the 3-dimensional generalization 3-surfaces correspond to free particles and the boundaries of the 3-surface correspond to partons in the sense that the quantum numbers of the elementary particles reside on the boundaries. Various boundary topologies (number of handles) correspond to various fermion families so that one obtains an explanation for the known elementary particle quantum numbers. This approach leads also to a natural topological description of the particle reactions as topology changes: for instance, two-particle decay corresponds to a decay of a 3-surface to two disjoint 3-surfaces.

This decay vertex does not however correspond to a direct generalization of trouser vertex of string models. Indeed, the important difference between TGD and string models is that the analogs of string world sheet diagrams do not describe particle decays but the propagation of particles via different routes. Particle reactions are described by generalized Feynman diagrams for which 3-D light-like surface describing particle propagating join along their ends at vertices. As 4-manifolds the space-time surfaces are therefore singular like Feynman diagrams as 1-manifolds.

Quite recently, it has turned out that fermionic strings inside space-time surfaces define an exact part of quantum TGD and that this is essential for understanding gravitation in long length scales. Also the analog of AdS/CFT duality emerges in that the Kähler metric can be defined either in terms of Kähler function identifiable as Kähler action assignable to Euclidian space-time regions or Kähler action + string action assignable to Minkowskian regions.

The recent view about construction of scattering amplitudes is very “stringy”. By strong form of holography string world sheets and partonic 2-surfaces provide the data needed to construct scattering amplitudes. Space-time surfaces are however needed to realize quantum-classical correspondence necessary to understand the classical correlates of quantum measurement. There is a huge generalization of the duality symmetry of hadronic string models.

The proposal is that scattering amplitudes can be regarded as sequences of computational operations for the Yangian of super-symplectic algebra. Product and co-product define the basic vertices and realized geometrically as partonic 2-surfaces and algebraically as multiplication for the elements of Yangian identified as super-symplectic Noether charges assignable to strings. Any computational sequences connecting given collections of algebraic objects at the opposite boundaries of causal diamond (CD) produce identical scattering amplitudes.

### 1.2.3 Fusion of the Two Approaches via a Generalization of the Space-Time Concept

The problem is that the two approaches to TGD seem to be mutually exclusive since the orbit of a particle like 3-surface defines 4-dimensional surface, which differs drastically from the topologically trivial macroscopic space-time of General Relativity. The unification of these approaches forces a considerable generalization of the conventional space-time concept. First, the topologically trivial 3-space of General Relativity is replaced with a “topological condensate” containing matter as particle like 3-surfaces “glued” to the topologically trivial background 3-space by connected sum operation. Secondly, the assumption about connectedness of the 3-space is given up. Besides the “topological condensate” there could be “vapor phase” that is a “gas” of particle like 3-surfaces and string like objects (counterpart of the “baby universes” of GRT) and the non-conservation of energy in GRT corresponds to the transfer of energy between different sheets of the space-time and possible existence vapour phase.

. What one obtains is what I have christened as many-sheeted space-time (see **Fig.** <http://tgdtheory.fi/appfigures/manysheeted.jpg> or **Fig. ??** in the appendix of this book). One particular aspect is topological field quantization meaning that various classical fields assignable to a physical system correspond to space-time sheets representing the classical fields to that particular system. One can speak of the field body of a particular physical system. Field body consists of topological light rays, and electric and magnetic flux quanta. In Maxwell's theory the physical system does not possess this kind of field identity. The notion of the magnetic body is one of the key players in TGD inspired theory of consciousness and quantum biology. The existence of monopole flux tubes requiring no current as a source of the magnetic field makes it possible to understand the existence of magnetic fields in cosmological and astrophysical scales.

This picture became more detailed with the advent of zero energy ontology (ZEO). The basic notion of ZEO is causal diamond (CD) identified as the Cartesian product of  $CP_2$  and of the intersection of future and past directed light-cones and having scale coming as an integer multiple of  $CP_2$  size is fundamental. CDs form a fractal hierarchy and zero energy states decompose to products of positive and negative energy parts assignable to the opposite boundaries of CD defining the ends of the space-time surface. The counterpart of zero energy state in positive energy ontology is the pair of initial and final states of a physical event, say particle reaction.

At space-time level ZEO means that 3-surfaces are pairs of space-like 3-surfaces at the opposite light-like boundaries of CD. Since the extremals of Kähler action connect these, one can say that by holography the basic dynamical objects are the space-time surface connecting these 3-surfaces and identifiable as analogs of Bohr orbits. This changes totally the vision about notions like self-organization: self-organization by quantum jumps does not take for a 3-D system but for the entire 4-D field pattern associated with it.

General Coordinate Invariance (GCI) allows to identify the basic dynamical objects as space-like 3-surfaces at the ends of space-time surface at boundaries of CD: this means that space-time surface is analogous to Bohr orbit. An alternative identification of the lines of generalized Feynman diagrams is as light-like 3-surfaces at which the signature of the induced metric changes from Minkowskian to Euclidian. Also the Euclidian 4-D regions can have a similar interpretation. The requirement that the two interpretations are equivalent, leads to a strong form of General Coordinate Invariance. The outcome is effective 2-dimensionality stating that the partonic 2-surfaces identified as intersections of the space-like ends of space-time surface and light-like wormhole throats are the fundamental objects. That only effective 2-dimensionality is in question is due to the effects caused by the failure of strict determinism of Kähler action. In finite length scale resolution these effects can be neglected below UV cutoff and above IR cutoff. One can also speak about a strong form of holography.

The understanding of the super symplectic invariance leads to the proposal that super symplectic algebra and other Kac-Moody type algebras labelled by non-negative multiples of basic conformal weights allow a hierarchy of symmetry breakings in which the analog of gauge symmetry breaks down to a genuine dynamical symmetry. This gives rise to fractal hierarchies of algebras and symmetry breakings. This breaking can occur also for ordinary conformal algebras if one restricts the conformal weights to be non-negative integers.

### 1.3 Basic Objections

Objections are the most powerful tool in theory building. The strongest objection against TGD is the observation that all classical gauge fields are expressible in terms of four embedding space coordinates only- essentially  $CP_2$  coordinates. The linear superposition of classical gauge fields taking place independently for all gauge fields is lost. This would be a catastrophe without many-sheeted space-time. Instead of gauge fields, only the effects such as gauge forces are superposed. Particles topologically condense to several space-time sheets simultaneously and experience the sum of gauge forces. This transforms the weakness to extreme economy: in a typical unified theory the number of primary field variables is countered in hundreds if not thousands, now it is just four.

Second objection is that TGD space-time is quite too simple as compared to GRT space-time due to the embeddability to 8-D embedding space. One can also argue that Poincare invariant theory of gravitation cannot be consistent with General Relativity. The above interpretation makes it possible to understand the relationship to GRT space-time and how the Equivalence Principle



(EP) follows from Poincare invariance of TGD. The interpretation of GRT space-time is as effective space-time obtained by replacing many-sheeted space-time with Minkowski space with effective metric determined as a sum of Minkowski metric and sum over the deviations of the induced metrics of the space-time sheets from Minkowski metric. Poincare invariance strongly suggests classical EP for the GRT limit in long length scales at least. One can also consider other kinds of limits such as the analog of GRT limit for Euclidian space-time regions assignable to elementary particles. In this case deformations of  $CP_2$  metric define a natural starting point and  $CP_2$  indeed defines a gravitational instanton with a very large cosmological constant in Einstein-Maxwell theory. Also gauge potentials of the standard model correspond classically to superpositions of induced gauge potentials over space-time sheets.

### 1.3.1 Topological Field Quantization

Topological field quantization distinguishes between TGD based and more standard - say Maxwellian - notion of field. In Maxwell's fields created by separate systems superpose and one cannot tell which part of field comes from which system except theoretically. In TGD these fields correspond to different space-time sheets and only their effects on test particle superpose. Hence physical systems have well-defined field identifies - field bodies - in particular magnetic bodies.

The notion of magnetic body carrying dark matter with non-standard large value of Planck constant has become central concept in TGD inspired theory of consciousness and living matter, and by starting from various anomalies of biology one ends up to a rather detailed view about the role of magnetic body as intentional agent receiving sensory input from the biological body and controlling it using EEG and its various scaled up variants as a communication tool. Among other things this leads to models for cell membrane, nerve pulse, and EEG.

## 1.4 Quantum TGD as Spinor Geometry of World of Classical Worlds

A turning point in the attempts to formulate a mathematical theory was reached after seven years from the birth of TGD. The great insight was "Do not quantize". The basic ingredients to the new approach have served as the basic philosophy for the attempt to construct Quantum TGD since then and have been the following ones.

### 1.4.1 World of Classical Worlds

The notion of WCW reduces the interacting quantum theory to a theory of free WCW spinor fields.

1. Quantum theory for extended particles is free(!), classical(!) field theory for a generalized Schrödinger amplitude identified as WCW spinor in the configuration space  $CH$  ("world of classical worlds", WCW) consisting of all possible 3-surfaces in  $H$ . "All possible" means that surfaces with arbitrary many disjoint components and with arbitrary internal topology and also singular surfaces topologically intermediate between two different manifold topologies are included.
2. 4-D general coordinate invariance forces holography and replaces the ill-defined path integral over all space-time surfaces with a discrete sum over 4-D analogs of Bohr orbits for particles identified as 3-surfaces. Holography means that basic objects are these analogs of Bohr orbits. Since there is no quantization at the level of WCW, one has an analog of wave mechanics with point-like particles replaced with 4-D Bohr orbits.
3. One must geometrize WCW as the space of Bohr orbits. In an infinite-dimensional situation the existence of geometry requires maximal symmetries already in the case of loop spaces. Physics is unique from its mathematical existence.

WCW is endowed with metric and spinor structure so that one can define various metric related differential operators, say Dirac operators, appearing in the field equations of the theory <sup>1</sup>

---

<sup>1</sup>There are four kinds of Dirac operators in TGD. The geometrization of quantum theory requires Kähler metric definable either in terms of Kähler function identified as a the bosonic action for Euclidian space-time regions

### 1.4.2 Identification of Kähler function

The evolution of these basic ideas has been rather slow but has gradually led to a rather beautiful vision. One of the key problems has been the definition of Kähler function. Kähler function is Kähler action for a preferred extremal assignable to a given 3-surface but what this preferred extremal is? The obvious first guess was as absolute minimum of Kähler action but could not be proven to be right or wrong. One big step in the progress was boosted by the idea that TGD should reduce to almost topological QFT in which braids would replace 3-surfaces in finite measurement resolution, which could be inherent property of the theory itself and imply discretization at partonic 2-surfaces with discrete points carrying fermion number.

It took long time to realize that there is no discretization in 4-D sense - this would lead to difficulties with basic symmetries. Rather, the discretization occurs for the parameters characterizing co-dimension 2 objects representing the information about space-time surface so that they belong to some algebraic extension of rationals. These 2-surfaces - string world sheets and partonic 2-surfaces - are genuine physical objects rather than a computational approximation. Physics itself approximates itself, one might say! This is of course nothing but strong form of holography.

1. TGD as almost topological QFT vision suggests that Kähler action for preferred extremals reduces to Chern-Simons term assigned with space-like 3-surfaces at the ends of space-time (recall the notion of causal diamond (CD)) and with the light-like 3-surfaces at which the signature of the induced metric changes from Minkowskian to Euclidian. Minkowskian and Euclidian regions would give at wormhole throats the same contribution apart from coefficients and in Minkowskian regions the  $\sqrt{g_4}$  factor coming from metric would be imaginary so that one would obtain sum of real term identifiable as Kähler function and imaginary term identifiable as the ordinary Minkowskian action giving rise to interference effects and stationary phase approximation central in both classical and quantum field theory.

Imaginary contribution - the presence of which I realized only after 33 years of TGD - could also have topological interpretation as a Morse function. On physical side the emergence of Euclidian space-time regions is something completely new and leads to a dramatic modification of the ideas about black hole interior.

2. The way to achieve the reduction to Chern-Simons terms is simple. The vanishing of Coulomb contribution to Kähler action is required and is true for all known extremals if one makes a general ansatz about the form of classical conserved currents. The so called weak form of electric-magnetic duality defines a boundary condition reducing the resulting 3-D terms to Chern-Simons terms. In this way almost topological QFT results. But only "almost" since the Lagrange multiplier term forcing electric-magnetic duality implies that Chern-Simons action for preferred extremals depends on metric.

### 1.4.3 WCW spinor fields

Classical WCW spinor fields are analogous to Schrödinger amplitudes and the construction of WCW Kähler geometry reduces to the second quantization of free spinor fields of  $H$ .

1. The WCW metric is given by anticommutators of WCW gamma matrices which also have interpretation as supercharges assignable to the generators of WCW isometries and allowing expression as non-conserved Noether charges. Holography implies zero energy ontology (ZEO) meaning that zero energy states are superpositions of Bohr orbits connecting boundaries of causal diamond (CD). CDs form a fractal hierarchy and their space forming the spine of WCW is finite-dimensional and can be geometrized. The alternative interpretation is as a superposition of pairs of ordinary 3-D fermionic states assignable to the ends of the space-time surfaces.
2. There are several Dirac operators. WCW Dirac operator  $D_{WCW}$  appears in Super-symplectic gauge conditions analogous to Super Virasoro conditions. The algebraic variant of the  $H$

---

or as anti-commutators for WCW gamma matrices identified as conformal Noether super-charges associated with the second quantized modified Dirac action consisting of string world sheet term and possibly also modified Dirac action in Minkowskian space-time regions. These two possible definitions reflect a duality analogous to AdS/CFT duality.

Dirac operator  $D_H$  appears in fermionic correlation functions: this is due to the fact that free fermions appearing as building bricks of WCW gamma matrices are modes of  $D_H$ . The modes of  $D_H$  define the ground states of super-symplectic representations. There is also the modified Dirac operator  $D_{X^4}$  acting on the induced spinors at space-time surfaces and it is dictated by symmetry one the action fixing the space-time surfaces as Bohr orbits is fixed.  $D_H$  is needed since it determines the expressions of WCW gamma matrices as Noether charges assignable to 3-surfaces at the ends of WCW.

#### 1.4.4 The role of modified Dirac action

1. By quantum classical correspondence, the construction of WCW spinor structure in sectors assignable to CDs reduces to the second quantization of the induced spinor fields of  $H$ . The basic action is so called modified Dirac action in which gamma matrices are replaced with the modified gamma matrices defined as contractions of the canonical momentum currents of the bosonic action defining the space-time surfaces with the embedding space gamma matrices. In this way one achieves super-conformal symmetry and conservation of fermionic currents among other things and a consistent Dirac equation.

Modified Dirac action is needed to define WCW gamma matrices as super charges assignable to WCW isometry generators identified as generators of symplectic transformations and by holography are needed only at the 3-surface at the boundaries of WCW. It is important to notice that the modified Dirac equation does not determine propagators since induced spinor fields are obtained from free second quantized spinor fields of  $H$ . This means enormous simplification and makes the theory calculable.

2. An important interpretational problem relates to the notion of the induced spinor connection. The presence of classical W boson fields is in conflict with the classical conservation of em charge since the coupling to classical W fields changes em charge.

One way out of the problem is the fact that the quantum averages of weak and gluon fields vanish unlike the quantum average of the em field. This leads to a rather precise understanding of electroweak symmetry breaking as being due the fact that color symmetries rotate space-time surfaces and also affect the induced weak fields.

One can also consider a stronger condition. If one requires that the spinor modes have well-defined em charge, one must assume that the modes in the generic situation are localized at 2-D surfaces - string world sheets or perhaps also partonic 2-surfaces - at which classical W boson fields vanish. Covariantly constant right handed neutrinos generating super-symmetries forms an exception. The vanishing of the  $Z^0$  field is possible for Kähler-Dirac action and should hold true at least above weak length scales. This implies that the string model in 4-D space-time becomes part of TGD. Without these conditions classical weak fields can vanish above weak scale only for the GRT limit of TGD for which gauge potentials are sums over those for space-time sheets.

The localization would simplify the mathematics enormously and one can solve exactly the Kähler-Dirac equation for the modes of the induced spinor field just like in super string models.

At the light-like 3-surfaces the signature of the induced metric changes from Euclidian to Minkowskian so that  $\sqrt{g_4}$  vanishes. One can pose the condition that the algebraic analog of the massless Dirac equation is satisfied by the modes of the modified-Dirac action assignable to the Chern-Simons-Kähler action.

## 1.5 Construction of scattering amplitudes

### 1.5.1 Reduction of particle reactions to space-time topology

Particle reactions are identified as topology changes [A8, A10, A11]. For instance, the decay of a 3-surface to two 3-surfaces corresponds to the decay  $A \rightarrow B+C$ . Classically this corresponds to a path of WCW leading from 1-particle sector to 2-particle sector. At quantum level this corresponds to the dispersion of the generalized Schrödinger amplitude localized to 1-particle sector to two-particle

sector. All coupling constants should result as predictions of the theory since no nonlinearities are introduced.

During years this naïve and very rough vision has of course developed a lot and is not anymore quite equivalent with the original insight. In particular, the space-time correlates of Feynman graphs have emerged from theory as Euclidian space-time regions and the strong form of General Coordinate Invariance has led to a rather detailed and in many respects un-expected visions. This picture forces to give up the idea about smooth space-time surfaces and replace space-time surface with a generalization of Feynman diagram in which vertices represent the failure of manifold property. I have also introduced the word “world of classical worlds” (WCW) instead of rather formal “configuration space”. I hope that “WCW” does not induce despair in the reader having tendency to think about the technicalities involved!

### 1.5.2 Construction of the counterparts of S-matrices

What does one mean with the counterpart of S-matrix in the TGD framework has been a long standing problem. The development of ZEO based quantum measurement theory has led to a rough overall view of the situation.

1. There are two kinds of state function reductions (SFRs). ”Small” SFRs (SSFRs) following the TGD counterpart of a unitary time evolution defines a sequence of SFRs, which is analogous to a sequence of repeated quantum measurements associated with the Zeno effect. In wave mechanics nothing happens in these measurements. In quantum optics these measurements correspond to weak measurements. In TGD SSFR affects the zero energy state but leaves the 3-D state at the passive boundary of CD unaffected.
2. In TGD framework each SSFR is preceded by a counterpart of a unitary time evolution, which means dispersion in the space of CDs and unitary time evolution in fermionic degrees of freedom such that the passive boundary of CDs and 3-D states at it are unaffected but a superposition of CDs with varying active boundaries in the space of CDs is formed. In SSFR a localization in the space of CDs occurs such that the active is fixed. In a statistical sense the size of the CD increases and the increasing distance between the tips of the CD gives rise to the arrow of geometric time.
3. Also ”big” SFRs (BSFRs) can occur and they correspond to ordinary SFRs. In BSFR the roles of the active and passive boundary are changed and this means that the arrow of time is changed. Big SFR occurs when the SSFR corresponds to a quantum measurement, which does not commute with the operators, which define the states at the passive boundary of CD as their eigenstates. This means a radical deviation from standard quantum measurement theory and has predictions in all scales.
4. One can assign the counterpart of S-matrix to the unitary time evolution between two subsequent SSFRs and also to the counterpart of S-matrix associated with BSFR. At least in the latter case the dimension of the state space can increase since at least BSFRs lead to the increase of the dimension of algebraic extension of rationals assignable to the space-time surface by  $M^8 - H$  duality. Unitarity is therefore replaced with isometry.
5. I have also considered the possibility that unitary S-matrix could be replaced in the fermionic degrees of freedom with Kähler metric of the state space satisfying analogs of unitarity conditions but it seems that this is un-necessary and also too outlandish an idea.

### 1.5.3 The notion of M-matrix

1. The most ambitious dream is that zero energy states correspond to a complete solution basis for the Dirac operators associated with WCWs associated with the spaces of CDs with fixed passive boundary: this would define an S-matrix assignable to SFR. Also the analog of S-matrix for the localizations of the states to the active boundary assignable to the BSFR changing the state at the passive boundary of CD is needed.

2. If one allows entanglement between positive and energy parts of the zero energy state but assumes that the states at the passive boundary are fixed, one must introduce the counterpart of the density matrix, or rather its square root. This classical free field theory would dictate what I have called M-matrices defined between positive and negative energy parts of zero energy states which form orthonormal rows of what I call U-matrix as a matrix defined between zero energy states. A given M-matrix in turn would decompose to a product of a hermitian square root of density matrix and unitary S-matrix.
3. M-matrix would define time-like entanglement coefficients between positive and negative energy parts of zero energy states (all net quantum numbers vanish for them) and can be regarded as a hermitian square root of density matrix multiplied by a unitary S-matrix. Quantum theory would be in a well-defined sense a square root of thermodynamics. The orthogonality and hermiticity of the M-matrices commuting with S-matrix means that they span infinite-dimensional Lie algebras acting as symmetries of the S-matrix. Therefore quantum TGD would reduce to group theory in a well-defined sense.
4. In fact the Lie algebra of Hermitian M-matrices extends to Kac-Moody type algebra obtained by multiplying hermitian square roots of density matrices with powers of the S-matrix. Also the analog of Yangian algebra involving only non-negative powers of S-matrix is possible and would correspond to a hierarchy of CDs with the temporal distances between tips coming as integer multiples of the  $CP_2$  time.

The M-matrices associated with CDs are obtained by a discrete scaling from the minimal CD and characterized by integer  $n$  are naturally proportional to a representation matrix of scaling:  $S(n) = S^n$ , where  $S$  is unitary S-matrix associated with the minimal CD [K10]. This conforms with the idea about unitary time evolution as exponent of Hamiltonian discretized to integer power of  $S$  and represented as scaling with respect to the logarithm of the proper time distance between the tips of CD.

5. I have also considered the notion of U-matrix. U-matrix elements between M-matrices for various CDs are proportional to the inner products  $Tr[S^{-n_1} \circ H^i H^j \circ S^{n_2} \lambda]$ , where  $\lambda$  represents unitarily the discrete Lorentz boost relating the moduli of the active boundary of CD and  $H^i$  form an orthonormal basis of Hermitian square roots of density matrices.  $\circ$  tells that  $S$  acts at the active boundary of CD only. I have proposed a general representation for the U-matrix, reducing its construction to that of the S-matrix.

## 1.6 TGD as a generalized number theory

Quantum T(opological)D(ynamics) as a classical spinor geometry for infinite-dimensional configuration space (“world of classical worlds”, WCW), p-adic numbers and quantum TGD, and TGD inspired theory of consciousness, have been for last ten years the basic three strongly interacting threads in the tapestry of quantum TGD. The fourth thread deserves the name “TGD as a generalized number theory”. It involves three separate threads: the fusion of real and various p-adic physics to a single coherent whole by requiring number theoretic universality discussed already, the formulation of quantum TGD in terms of complexified counterparts of classical number fields, and the notion of infinite prime. Note that one can identify subrings such as hyper-quaternions and hyper-octonions as sub-spaces of complexified classical number fields with Minkowskian signature of the metric defined by the complexified inner product.

### 1.6.1 The Threads in the Development of Quantum TGD

The development of TGD has involved several strongly interacting threads: physics as infinite-dimensional geometry; TGD as a generalized number theory, the hierarchy of Planck constants interpreted in terms of dark matter hierarchy, and TGD inspired theory of consciousness. In the following these threads are briefly described.

1. Quantum T(opological) G(eometro)D(ynamics) as a classical spinor geometry for infinite-dimensional WCW, p-adic numbers and quantum TGD, and TGD inspired theory of consciousness and of quantum biology have been for last decade of the second millenium the basic three strongly interacting threads in the tapestry of quantum TGD.

2. The discussions with Tony Smith initiated a fourth thread which deserves the name “TGD as a generalized number theory”. The basic observation was that classical number fields might allow a deeper formulation of quantum TGD. The work with Riemann hypothesis made time ripe for realization that the notion of infinite primes could provide, not only a reformulation, but a deep generalization of quantum TGD. This led to a thorough and rather fruitful revision of the basic views about what the final form and physical content of quantum TGD might be. Together with the vision about the fusion of p-adic and real physics to a larger coherent structure these sub-threads fused to the “physics as generalized number theory” thread.
3. A further thread emerged from the realization that by quantum classical correspondence TGD predicts an infinite hierarchy of macroscopic quantum systems with increasing sizes, that it is not at all clear whether standard quantum mechanics can accommodate this hierarchy, and that a dynamical quantized Planck constant might be necessary and strongly suggested by the failure of strict determinism for the fundamental variational principle. The identification of hierarchy of Planck constants labelling phases of dark matter would be natural. This also led to a solution of a long standing puzzle: what is the proper interpretation of the predicted fractal hierarchy of long ranged classical electro-weak and color gauge fields. Quantum classical correspondences allows only single answer: there is infinite hierarchy of p-adically scaled up variants of standard model physics and for each of them also dark hierarchy. Thus TGD Universe would be fractal in very abstract and deep sense.

The chronology based identification of the threads is quite natural but not logical and it is much more logical to see p-adic physics, the ideas related to classical number fields, and infinite primes as sub-threads of a thread which might be called “physics as a generalized number theory”. In the following I adopt this view. This reduces the number of threads to three corresponding to geometric, number theoretic and topological views of physics.

TGD forces the generalization of physics to a quantum theory of consciousness, and TGD as a generalized number theory vision leads naturally to the emergence of p-adic physics as physics of cognitive representations.

### 1.6.2 Number theoretic vision very briefly

Number theoretic vision about quantum TGD involves notions like adelic physics,  $M^8 - H$  duality and number theoretic universality. A short review of the basic ideas that have developed during years is in order.

1. The physical interpretation of  $M^8$  is as an analog of momentum space and  $M^8 - H$  duality is analogous to momentum-position duality of ordinary wave mechanics.
2. Adelic physics means that all classical number fields, all p-adic number fields and their extensions induced by extensions of rationals and defining adeles, and also finite number fields are basic mathematical building bricks of physics.

The complexification of  $M^8$ , identified as complexified octonions, would provide a realization of this picture and  $M^8 - H$  duality would map the algebraic physics in  $M^8$  to the ordinary physics in  $M^4 \times CP_2$  described in terms of partial differential equations.

3. Negentropy Maximization Principle (NMP) states that the conscious information assignable with cognition representable measured in terms of p-adic negentropy increases in statistical sense.

NMP is mathematically completely analogous to the second law of thermodynamics and number theoretic evolution as an unavoidable statistical increase of the dimension of the algebraic extension of rationals characterizing a given space-time region implies it. There is no paradox involved: the p-adic negentropy measures the conscious information assignable to the entanglement of two systems regarded as a conscious entity whereas ordinary entropy measures the lack of information about the quantum state of either entangled system.

4. Number theoretical universality requires that space-time surfaces or at least their  $M^8 - H$  duals in  $M_c^8$  are defined for both reals and various p-adic number fields. This is true if they are

defined by polynomials with integer coefficients as surfaces in  $M^8$  obeying number theoretic holography realized as associativity of the normal space of 4-D surface using as holographic data 3-surfaces at mass shells identified in terms of roots of a polynomial. A physically motivated additional condition is that the coefficients of the polynomials are smaller than their degrees.

5. Galois confinement is a key piece of the number theoretic vision. It states that the momenta of physical states are algebraic integers in the extensions of rationals assignable to the space-time region considered. These numbers are in general complex and are not consistent with particle in box quantization. The proposal is that physical states satisfy Galois confinement being thus Galois singlets and having therefore total momenta, whose components are ordinary integers, when momentum unit defined by the scale of causal diamond (CD) is used.
6. The notion of p-adic prime was introduced in p-adic mass calculations that started the developments around 1995. p-Adic length scale hypothesis states that p-adic primes near powers of 2 have a special physical role (as possibly also the powers of other small primes such as  $p = 3$ ).

The proposal is that p-adic primes correspond to ramified primes assignable to the extension and identified as divisors of the polynomial defined by the products of the root differences for the roots of the polynomial defining space-time space and having interpretation as values of, in general complex, virtual mass squared.

### 1.6.3 p-Adic TGD and fusion of real and p-adic physics to single coherent whole

The p-adic thread emerged for roughly ten years ago as a dim hunch that p-adic numbers might be important for TGD. Experimentation with p-adic numbers led to the notion of canonical identification mapping reals to p-adics and vice versa. The breakthrough came with the successful p-adic mass calculations using p-adic thermodynamics for Super-Virasoro representations with the super-Kac-Moody algebra associated with a Lie-group containing standard model gauge group. Although the details of the calculations have varied from year to year, it was clear that p-adic physics reduces not only the ratio of proton and Planck mass, the great mystery number of physics, but all elementary particle mass scales, to number theory if one assumes that primes near prime powers of two are in a physically favored position. Why this is the case, became one of the key puzzles and led to a number of arguments with a common gist: evolution is present already at the elementary particle level and the primes allowed by the p-adic length scale hypothesis are the fittest ones.

It became very soon clear that p-adic topology is not something emerging in Planck length scale as often believed, but that there is an infinite hierarchy of p-adic physics characterized by p-adic length scales varying to even cosmological length scales. The idea about the connection of p-adics with cognition motivated already the first attempts to understand the role of the p-adics and inspired “Universe as Computer” vision but time was not ripe to develop this idea to anything concrete (p-adic numbers are however in a central role in TGD inspired theory of consciousness). It became however obvious that the p-adic length scale hierarchy somehow corresponds to a hierarchy of intelligences and that p-adic prime serves as a kind of intelligence quotient. Ironically, the almost obvious idea about p-adic regions as cognitive regions of space-time providing cognitive representations for real regions had to wait for almost a decade for the access into my consciousness.

In string model context one tries to reduce the physics to Planck scale. The price is the inability to say anything about physics in long length scales. In TGD p-adic physics takes care of this shortcoming by predicting the physics also in long length scales.

There were many interpretational and technical questions crying for a definite answer.

1. What is the relationship of p-adic non-determinism to the classical non-determinism of the basic field equations of TGD? Are the p-adic space-time region genuinely p-adic or does p-adic topology only serve as an effective topology? If p-adic physics is direct image of real physics, how the mapping relating them is constructed so that it respects various symmetries? Is the basic physics p-adic or real (also real TGD seems to be free of divergences) or both? If it is both, how should one glue the physics in different number field together to get *the* Physics? Should one perform p-adicization also at the level of the WCW? Certainly the p-adicization at the level of super-conformal representation is necessary for the p-adic mass calculations.

2. Perhaps the most basic and most irritating technical problem was how to precisely define p-adic definite integral which is a crucial element of any variational principle based formulation of the field equations. Here the frustration was not due to the lack of solution but due to the too large number of solutions to the problem, a clear symptom for the sad fact that clever inventions rather than real discoveries might be in question. Quite recently I however learned that the problem of making sense about p-adic integration has been for decades central problem in the frontier of mathematics and a lot of profound work has been done along same intuitive lines as I have proceeded in TGD framework. The basic idea is certainly the notion of algebraic continuation from the world of rationals belonging to the intersection of real world and various p-adic worlds.

Despite various uncertainties, the number of the applications of the poorly defined p-adic physics has grown steadily and the applications turned out to be relatively stable so that it was clear that the solution to these problems must exist. It became only gradually clear that the solution of the problems might require going down to a deeper level than that represented by reals and p-adics.

The key challenge is to fuse various p-adic physics and real physics to single larger structure. This has inspired a proposal for a generalization of the notion of number field by fusing real numbers and various p-adic number fields and their extensions along rationals and possible common algebraic numbers. This leads to a generalization of the notions of embedding space and space-time concept and one can speak about real and p-adic space-time sheets. One can talk about adelic space-time, embedding space, and WCW.

The corresponds of real 4-surfaces with the p-adic ones is induced by number theoretical discretization using points of 4-surfaces  $Y^4 \subset M_c^8$  identifiable as 8-momenta, whose components are assumed to be algebraic integers in an extension of rationals defined by the extension of rationals associated with a polynomial  $P$  with integer coefficients smaller than the degree of  $P$ . These points define a cognitive representation, which is universal in the sense that it exists also in the algebraic extensions of p-adic numbers. The points of the cognitive representations associated with the mass shells with mass squared values identified as roots of  $P$  are enough since  $M^8 - H$  duality can be used at both  $M^8$  and  $H$  sides and also in the p-adic context. The mass shells are special in that they allow for Minkowski coordinates very large cognitive representations unlike the interiors of the 4-surfaces determined by holography by using the data defined by the 3-surfaces at the mass shells. The higher the dimension of the algebraic extension associated with  $P$ , the better the accuracy of the cognitive representation.

Adelization providing number theoretical universality reduces to algebraic continuation for the amplitudes from this intersection of reality and various p-adicities - analogous to a back of a book - to various number fields. There are no problems with symmetries but canonical identification is needed: various group invariant of the amplitude are mapped by canonical identification to various p-adic number fields. This is nothing but a generalization of the mapping of the p-adic mass squared to its real counterpart in p-adic mass calculations.

This leads to surprisingly detailed predictions and far reaching conjectures. For instance, the number theoretic generalization of entropy concept allows negentropic entanglement central for the applications to living matter (see **Fig.** <http://tgdtheory.fi/appfigures/cat.jpg> or **Fig. ??** in the appendix of this book). One can also understand how preferred p-adic primes could emerge as so called ramified primes of algebraic extension of rationals in question and characterizing string world sheets and partonic 2-surfaces. Preferred p-adic primes would be ramified primes for extensions for which the number of p-adic continuations of two-surfaces to space-time surfaces (imaginings) allowing also real continuation (realization of imagination) would be especially large. These ramifications would be winners in the fight for number theoretical survival. Also a generalization of p-adic length scale hypothesis emerges from NMP [K9].

The characteristic non-determinism of the p-adic differential equations suggests strongly that p-adic regions correspond to “mind stuff”, the regions of space-time where cognitive representations reside. This interpretation implies that p-adic physics is physics of cognition. Since Nature is probably a brilliant simulator of Nature, the natural idea is to study the p-adic physics of the cognitive representations to derive information about the real physics. This view encouraged by TGD inspired theory of consciousness clarifies difficult interpretational issues and provides a clear interpretation for the predictions of p-adic physics.



### 1.6.4 Infinite primes

The discovery of the hierarchy of infinite primes and their correspondence with a hierarchy defined by a repeatedly second quantized arithmetic quantum field theory gave a further boost for the speculations about TGD as a generalized number theory.

After the realization that infinite primes can be mapped to polynomials possibly representable as surfaces geometrically, it was clear how TGD might be formulated as a generalized number theory with infinite primes forming the bridge between classical and quantum such that real numbers, p-adic numbers, and various generalizations of p-adics emerge dynamically from algebraic physics as various completions of the algebraic extensions of complexified quaternions and octonions. Complete algebraic, topological and dimensional democracy would characterize the theory.

The infinite primes at the first level of hierarchy, which represent analogs of bound states, can be mapped to irreducible polynomials, which in turn characterize the algebraic extensions of rationals defining a hierarchy of algebraic physics continuable to real and p-adic number fields. The products of infinite primes in turn define more general algebraic extensions of rationals. The interesting question concerns the physical interpretation of the higher levels in the hierarchy of infinite primes and integers mappable to polynomials of  $n > 1$  variables.

## 1.7 An explicit formula for $M^8 - H$ duality

$M^8 - H$  duality is a generalization of momentum-position duality relating the number theoretic and geometric views of physics in TGD and, despite that it still involves poorly understood aspects, it has become a fundamental building block of TGD. One has 4-D surfaces  $Y^4 \subset M_c^8$ , where  $M_c^8$  is complexified  $M^8$  having interpretation as an analog of complex momentum space and 4-D spacetime surfaces  $X^4 \subset H = M^4 \times CP_2$ .  $M_c^8$ , equivalently  $E_c^8$ , can be regarded as complexified octonions.  $M_c^8$  has a subspace  $M_c^4$  containing  $M^4$ .

**Comment:** One should be very cautious with the meaning of "complex". Complexified octonions involve a complex imaginary unit  $i$  commuting with the octonionic imaginary units  $I_k$ .  $i$  is assumed to also appear as an imaginary unit also in complex algebraic numbers defined by the roots of polynomials  $P$  defining holographic data in  $M_c^8$ .

In the following  $M^8 - H$  duality and its twistor lift are discussed and an explicit formula for the dualities are deduced. Also possible variants of the duality are discussed.

### 1.7.1 Holography in $H$

$X^4 \subset H$  satisfies holography and is analogous to the Bohr orbit of a particle identified as a 3-surface. The proposal is that holography reduces to a 4-D generalization of holomorphy so that  $X^4$  is a simultaneous zero of two functions of complex  $CP_2$  coordinates and of what I have called Hamilton-Jacobi coordinates of  $M^4$  with a generalized Kähler structure.

The simplest choice of the Hamilton-Jacobi coordinates is defined by the decomposition  $M^4 = M^2 \times E^2$ , where  $M^2$  is endowed with hypercomplex structure defined by light-like coordinates  $(u, v)$ , which are analogous to  $z$  and  $\bar{z}$ . Any analytic map  $u \rightarrow f(u)$  defines a new set of light-like coordinates and corresponds to a solution of the massless d'Alembert equation in  $M^2$ .  $E^2$  has some complex coordinates with imaginary unit defined by  $i$ .

The conjecture is that also more general Hamilton-Jacobi structures for which the tangent space decomposition is local are possible. Therefore one would have  $M^4 = M^2(x) \times E^2(x)$ . These would correspond to non-equivalent complex and Kähler structures of  $M^4$  analogous to those possessed by 2-D Riemann surfaces and parametrized by moduli space.

### 1.7.2 Number theoretic holography in $M_c^8$

$Y^4 \subset M_c^8$  satisfies number theoretic holography defining dynamics, which should reduce to associativity in some sense. The Euclidian complexified normal space  $N^4(y)$  at a given point  $y$  of  $Y^4$  is required to be associative, i.e. quaternionic. Besides this,  $N^4(i)$  contains a preferred complex Euclidian 2-D subspace  $Y^2(y)$ . Also the spaces  $Y^2(x)$  define an integrable distribution. I have assumed that  $Y^2(x)$  can depend on the point  $y$  of  $Y^4$ .

These assumptions imply that the normal space  $N(y)$  of  $Y^4$  can be parameterized by a point of  $CP_2 = SU(3)/U(2)$ . This distribution is always integrable unlike quaternionic tangent space

distributions.  $M^8 - H$  duality assigns to the normal space  $N(y)$  a point of  $CP_2$ .  $M_c^4$  point  $y$  is mapped to a point  $x \in M^4 \subset M^4 \times CP_2$  defined by the real part of its inversion (conformal transformation): this formula involves effective Planck constant for dimensional reasons.

The 3-D holographic data, which partially fixes 4-surfaces  $Y^4$  is partially determined by a polynomial  $P$  with real integer coefficients smaller than the degree of  $P$ . The roots define mass squared values which are in general complex algebraic numbers and define complex analogs of mass shells in  $M_c^4 \subset M_c^8$ , which are analogs of hyperbolic spaces  $H^3$ . The 3-surfaces at these mass shells define 3-D holographic data continued to a surface  $Y^4$  by requiring that the normal space of  $Y^4$  is associative, i.e. quaternionic. These 3-surfaces are not completely fixed but an interesting conjecture is that they correspond to fundamental domains of tessellations of  $H^3$ .

What does the complexity of the mass shells mean? The simplest interpretation is that the space-like  $M^4$  coordinates (3-momentum components) are real whereas the time-like coordinate (energy) is complex and determined by the mass shell condition. One would have  $Re^2(E) - Im(E)^2 - p^2 = Re(m^2)$  and  $2Re(E)Im(E) = Im(m^2)$ . The condition for the real parts gives  $H^3$  when  $\sqrt{Re^2(E) - Im(E)^2}$  is taken as a time coordinate. The second condition allows to solve  $Im(E)$  in terms of  $Re(E)$  so that the first condition reduces to an equation of mass shell when  $\sqrt{(Re(E)^2 - Im(E)^2)}$ , expressed in terms of  $Re(E)$ , is taken as new energy coordinate  $E_{eff} = \sqrt{(Re(E)^2 - Im(E)^2)}$ . Is this deformation of  $H^3$  in imaginary time direction equivalent with a region of the hyperbolic 3-space  $H^3$ ?

One can look at the formula in more detail. Mass shell condition gives  $Re^2(E) - Im(E)^2 - p^2 = Re(m^2)$  and  $2Re(E)Im(E) = Im(m^2)$ . The condition for the real parts gives  $H^3$ , when  $\sqrt{Re^2(E) - Im(E)^2}$  is taken as an effective energy. The second condition allows to solve  $Im(E)$  in terms of  $Re(E)$  so that the first condition reduces to a dispersion relation for  $Re(E)^2$ .

$$Re(E)^2 = \frac{1}{2}(Re(m^2) - Im(m^2) + p^2)(1 \pm \sqrt{1 + \frac{2Im(m^2)^2}{(Re(m^2) - Im(m^2) + p^2)^2}}). \quad (1.1)$$

Only the positive root gives a non-tachyonic result for  $Re(m^2) - Im(m^2) > 0$ . For real roots with  $Im(m^2) = 0$  and at the high momentum limit the formula coincides with the standard formula. For  $Re(m^2) = Im(m^2)$  one obtains  $Re(E)^2 \rightarrow Im(m^2)/\sqrt{2}$  at the low momentum limit  $p^2 \rightarrow 0$ . Energy does not depend on momentum at all: the situation resembles that for plasma waves.

### 1.7.3 Can one find an explicit formula for $M^8 - H$ duality?

The dream is an explicit formula for the  $M^8 - H$  duality mapping  $Y^4 \subset M_c^8$  to  $X^4 \subset H$ . This formula should be consistent with the assumption that the generalized holomorphy holds true for  $X^4$ .

The following proposal is a more detailed variant of the earlier proposal for which  $Y^4$  is determined by a map  $g$  of  $M_c^4 \rightarrow SU(3)_c \subset G_{2,c}$ , where  $G_{2,c}$  is the complexified automorphism group of octonions and  $SU(3)_c$  is interpreted as a complexified color group.

This map defines a trivial  $SU(3)_c$  gauge field. The real part of  $g$  however defines a non-trivial real color gauge field by the non-linearity of the non-abelian gauge field with respect to the gauge potential. The quadratic terms involving the imaginary part of the gauge potential give an additional condition to the real part in the complex situation and cancel it. If only the real part of  $g$  contributes, this contribution would be absent and the gauge field is non-vanishing.

How could the automorphism  $g(x) \subset SU(3) \subset G_2$  give rise to  $M^8 - H$  duality?

1. The interpretation is that  $g(y)$  at given point  $y$  of  $Y^4$  relates the normal space at  $y$  to a fixed quaternionic/associative normal space at point  $y_0$ , which corresponds is fixed by some subgroup  $U(2)_0 \subset SU(3)$ . The automorphism property of  $g$  guarantees that the normal space is quaternionic/associative at  $y$ . This simplifies the construction dramatically.
2. The quaternionic normal sub-space (which has Euclidian signature) contains a complex sub-space which corresponds to a point of sphere  $S^2 = SO(3)/O(2)$ , where  $SO(3)$  is the quaternionic automorphism group. The interpretation could be in terms of a selection of spin quantization axes. The local choice of the preferred complex plane would not be unique

and is analogous to the possibility of having non-trivial Hamilton Jacobi structures in  $M^4$  characterized by the choice of  $M^2(x)$  and equivalently its normal subspace  $E^2(x)$ .

These two structures are independent apart from dependencies forced by the number theoretic dynamics. Hamilton-Jacobi structure means a selection of the quantization axis of spin and energy by fixing a distribution of light-like tangent vectors of  $M^4$  and the choice of the quaternionic normal sub-space fixes a choice of preferred quaternionic imaginary unit defining a quantization axis of the weak isospin.

3. The real part  $Re(g(y))$  defines a point of  $SU(3)$  and the bundle projection  $SU(3) \rightarrow CP_2$  in turn defines a point of  $CP_2 = SU(3)/U(2)$ . Hence one can assign to  $g$  a point of  $CP_2$  as  $M^8 - H$  duality requires and deduce an explicit formula for the point. This means a realization of the dream.
4. The construction requires a fixing of a quaternionic normal space  $N_0$  at  $y_0$  containing a preferred complex subspace at a single point of  $Y^4$  plus a selection of the function  $g$ . If  $M^4$  coordinates are possible for  $Y^4$ , the first guess is that  $g$  as a function of complexified  $M^4$  coordinates obeys generalized holomorphy with respect to complexified  $M^4$  coordinates in the same sense and in the case of  $X^4$ . This might guarantee that the  $M^8 - H$  image of  $Y^4$  satisfies the generalized holomorphy.
5. Also space-time surfaces  $X^4$  with  $M^4$  projection having a dimension smaller than 4 are allowed. I have proposed that they might correspond to singular cases for the above formula: a kind of blow-up would be involved. One can also consider a more general definition of  $Y^4$  allowing it to have a  $M^4$  projection with dimension smaller than 4 (say cosmic strings). Could one have implicit equations for the surface  $Y^4$  in terms of the complex coordinates of  $SU(3)_c$  and  $M^4$ ? Could this give for instance cosmic strings with a 2-D  $M^4$  projection and  $CP_2$  type extremals with 4-D  $CP_2$  projection and 1-D light-like  $M^4$  projection?

#### 1.7.4 What could the number theoretic holography mean physically?

What could be physical meaning of the number theoretic holography? The condition that has been assumed is that the  $CP_2$  coordinates at the mass shells of  $M_c^4 \subset M_c^8$  mapped to mass shells  $H^3$  of  $M^4 \subset M^4 \times CP_2$  are constant at the  $H^3$ . This is true if the  $g(y)$  defines the same  $CP_2$  point for a given component  $X_i^3$  of the 3-surface at a given mass shell.  $g$  is therefore fixed apart from a local  $U(2)$  transformation leaving the  $CP_2$  point invariant. A stronger condition would be that the  $CP_2$  point is the same for each component of  $X_i^3$  and even at each mass shell but this condition seems to be unnecessarily strong.

**Comment:** One can criticize this condition as too strong and one can consider giving up this condition. The motivation for this condition is that the number of algebraic points at the 3-surfaces associated with  $H^3$  explodes since the coordinates associated with normal directions vanish. Kind of cognitive explosion would be in question.

$SU(3)$  corresponds to a subgroup of  $G_2$  and one can wonder what the fixing of this subgroup could mean physically.  $G_2$  is 14-D and the coset space  $G_2/SU(3)$  is 6-D and a good guess is that it is just the 6-D twistor space  $SU(3)/U(1) \times U(1)$  of  $CP_2$ : at least the isometries are the same. The fixing of the  $SU(3)$  subgroup means fixing of a  $CP_2$  twistor. Physically this means the fixing of the quantization axis of color isospin and hypercharge.

#### 1.7.5 Twistor lift of the holography

What is interesting is that by replacing  $SU(3)$  with  $G_2$ , one obtains an explicit formula from the generalization of  $M^8 - H$  duality to that for the twistorial lift of TGD!

One can also consider a twistorial generalization of the above proposal for the number theoretic holography by allowing local  $G_2$  automorphisms interpreted as local choices of the color quantization axis.  $G_2$  elements would be fixed apart from a local  $SU(3)$  transformation at the components of 3-surfaces at mass shells. The choice of the color quantization axes for a connected 3-surface at a given mass shell would be the same everywhere. This choice is indeed very natural physically since 3-surface corresponds to a particle.

Is this proposal consistent with the boundary condition of the number theoretical holography mean in the case of 4-surfaces in  $M_c^8$  and  $M^4 \times CP_2$ ?

1. The selection of  $SU(3) \subset G_2$  for ordinary  $M^8 - H$  duality means that the  $G_{2,c}$  gauge field vanishes everywhere and the choice of color quantization axis is the same at all points of the 4-surface. The fixing of the  $CP_2$  point to be constant at  $H^3$  implies that the color gauge field at  $H^3 \subset M_c^8$  and its image  $H^3 \subset H$  vanish. One would have color confinement at the mass shells  $H_i^3$ , where the observations are made. Is this condition too strong?
2. The constancy of the  $G_2$  element at mass shells makes sense physically and means a fixed color quantization axis. The selection of a fixed  $SU(3) \subset G_2$  for entire space-time surface is in conflict with the non-constancy of  $G_2$  element unless  $G_2$  element differs at different points of 4-surface only by a multiplication of a local  $SU(3)_0$  element, that is local  $SU(3)$  transformation. This kind of variation of the  $G_2$  element would mean a fixed color group but varying choice of color quantization axis.
3. Could one consider the possibility that the local  $G_{2,c}$  element is free and defines the twistor lift of  $M^8 - H$  duality as something more fundamental than the ordinary  $M^8 - H$  duality based on  $SU(3)_c$ . This duality would make sense only at the mass shells so that only the spaces  $H^3 \times CP_2$  assignable to mass shells would make sense physically? In the interior  $CP_2$  would be replaced with the twistor space  $SU(3)/U(1) \times U(1)$ . Color gauge fields would be non-vanishing at the mass shells but outside the mass shells one would have  $G_2$  gauge fields. There is also a physical objection against the  $G_2$  option. The 14-D Lie algebra representation of  $G_2$  acts on the imaginary octonions which decompose with respect to the color group to  $1 \oplus 3 \oplus \bar{3}$ . The automorphism property requires that 1 can be transformed to 3 or  $\bar{3}$  to themselves: this requires that the decomposition contains  $3 \oplus \bar{3}$ . Furthermore, it must be possible to transform 3 and  $\bar{3}$  to themselves, which requires the presence of 8. This leaves only the decomposition  $8 \oplus 3 \oplus \bar{3}$ .  $G_2$  gluons would both color octet and triplets. In the TDG framework the only conceivable interpretation would be in terms of ordinary gluons and leptoquark-like gluons. This does not fit with the basic vision of TGD.

The choice of twistor as a selection of quantization axes should make sense also in the  $M^4$  degrees of freedom.  $M^4$  twistor corresponds to a choice of light-like direction at a given point of  $M^4$ . The spatial component of the light-like vector fixes the spin quantization axis. Its choice together with the light-likeness fixes the time direction and therefore the rest system and energy quantization axis. Light-like vector fixes also the choice of  $M^2$  and of  $E^2$  as its orthogonal complement. Therefore the fixing of  $M^4$  twistor as a point of  $SU(4)/SU(3) \times U(1)$  corresponds to a choice of the spin quantization axis and the time-like axis defining the rest system in which the energy is measured. This choice would naturally correspond to the Hamilton-Jacobi structure fixing the decompositions  $M^2(x) \times E^2(x)$ . At a given mass shell the choice of the quantization axis would be constant for a given  $X_i^3$ .

## 1.8 Hierarchy of Planck Constants and Dark Matter Hierarchy

By quantum classical correspondence space-time sheets can be identified as quantum coherence regions. Hence the fact that they have all possible size scales more or less unavoidably implies that Planck constant must be quantized and have arbitrarily large values. If one accepts this then also the idea about dark matter as a macroscopic quantum phase characterized by an arbitrarily large value of Planck constant emerges naturally as does also the interpretation for the long ranged classical electro-weak and color fields predicted by TGD. Rather seldom the evolution of ideas follows simple linear logic, and this was the case also now. In any case, this vision represents the fifth, relatively new thread in the evolution of TGD and the ideas involved are still evolving.

### 1.8.1 Dark Matter as Large $\hbar$ Phases

D. Da Rocha and Laurent Nottale [E1] have proposed that Schrödinger equation with Planck constant  $\hbar$  replaced with what might be called gravitational Planck constant  $\hbar_{gr} = \frac{GmM}{v_0}$  ( $\hbar = c = 1$ ).  $v_0$  is a velocity parameter having the value  $v_0 = 144.7 \pm .7$  km/s giving  $v_0/c = 4.6 \times 10^{-4}$ . This is rather near to the peak orbital velocity of stars in galactic halos. Also subharmonics and harmonics of  $v_0$  seem to appear. The support for the hypothesis coming from empirical data is impressive.

Nottale and Da Rocha believe that their Schrödinger equation results from a fractal hydrodynamics. Many-sheeted space-time however suggests that astrophysical systems are at some levels of the hierarchy of space-time sheets macroscopic quantum systems. The space-time sheets in question would carry dark matter.

Nottale's hypothesis would predict a gigantic value of  $h_{gr}$ . Equivalence Principle and the independence of gravitational Compton length on mass  $m$  implies however that one can restrict the values of mass  $m$  to masses of microscopic objects so that  $h_{gr}$  would be much smaller. Large  $h_{gr}$  could provide a solution of the black hole collapse (IR catastrophe) problem encountered at the classical level. The resolution of the problem inspired by TGD inspired theory of living matter is that it is the dark matter at larger space-time sheets which is quantum coherent in the required time scale [K18].

It is natural to assign the values of Planck constants postulated by Nottale to the space-time sheets mediating gravitational interaction and identifiable as magnetic flux tubes (quanta) possibly carrying monopole flux and identifiable as remnants of cosmic string phase of primordial cosmology. The magnetic energy of these flux quanta would correspond to dark energy and magnetic tension would give rise to negative "pressure" forcing accelerate cosmological expansion. This leads to a rather detailed vision about the evolution of stars and galaxies identified as bubbles of ordinary and dark matter inside magnetic flux tubes identifiable as dark energy.

Certain experimental findings suggest the identification  $h_{eff} = n \times h_{gr}$ . The large value of  $h_{gr}$  can be seen as a way to reduce the string tension of fermionic strings so that gravitational (in fact all!) bound states can be described in terms of strings connecting the partonic 2-surfaces defining particles (analogous to AdS/CFT description). The values  $h_{eff}/h = n$  can be interpreted in terms of a hierarchy of breakings of super-conformal symmetry in which the super-conformal generators act as gauge symmetries only for a sub-algebras with conformal weights coming as multiples of  $n$ . Macroscopic quantum coherence in astrophysical scales is implied. If also Kähler-Dirac action is present, part of the interior degrees of freedom associated with the Kähler-Dirac part of conformal algebra become physical. A possible is that fermionic oscillator operators generate super-symmetries and sparticles correspond almost by definition to dark matter with  $h_{eff}/h = n > 1$ . One implication would be that at least part if not all gravitons would be dark and be observed only through their decays to ordinary high frequency graviton ( $E = hf_{high} = h_{eff}f_{low}$ ) of bunch of  $n$  low energy gravitons.

### 1.8.2 Hierarchy of Planck Constants from the Anomalies of Neuroscience and Biology

The quantal ELF effects of ELF em fields on vertebrate brain have been known since seventies. ELF em fields at frequencies identifiable as cyclotron frequencies in magnetic field whose intensity is about 2/5 times that of Earth for biologically important ions have physiological effects and affect also behavior. What is intriguing that the effects are found only in vertebrates (to my best knowledge). The energies for the photons of ELF em fields are extremely low - about  $10^{-10}$  times lower than thermal energy at physiological temperatures- so that quantal effects are impossible in the framework of standard quantum theory. The values of Planck constant would be in these situations large but not gigantic.

This inspired the hypothesis that these photons correspond to so large a value of Planck constant that the energy of photons is above the thermal energy. The proposed interpretation was as dark photons and the general hypothesis was that dark matter corresponds to ordinary matter with non-standard value of Planck constant. If only particles with the same value of Planck constant can appear in the same vertex of Feynman diagram, the phases with different value of Planck constant are dark relative to each other. The phase transitions changing Planck constant can however make possible interactions between phases with different Planck constant but these interactions do not manifest themselves in particle physics. Also the interactions mediated by classical fields should be possible. Dark matter would not be so dark as we have used to believe.

The hypothesis  $h_{eff} = h_{gr}$  - at least for microscopic particles - implies that cyclotron energies of charged particles do not depend on the mass of the particle and their spectrum is thus universal although corresponding frequencies depend on mass. In bio-applications this spectrum would correspond to the energy spectrum of bio-photons assumed to result from dark photons by  $h_{eff}$  reducing phase transition and the energies of bio-photons would be in visible and UV range

associated with the excitations of bio-molecules.

Also the anomalies of biology (see for instance [K15, K16, K14] ) support the view that dark matter might be a key player in living matter.

### 1.8.3 Dark Matter as a Source of Long Ranged Weak and Color Fields

Long ranged classical electro-weak and color gauge fields are unavoidable in TGD framework. The smallness of the parity breaking effects in hadronic, nuclear, and atomic length scales does not however seem to allow long ranged electro-weak gauge fields. The problem disappears if long range classical electro-weak gauge fields are identified as space-time correlates for massless gauge fields created by dark matter. Also scaled up variants of ordinary electro-weak particle spectra are possible. The identification explains chiral selection in living matter and unbroken  $U(2)_{ew}$  invariance and free color in bio length scales become characteristics of living matter and of bio-chemistry and bio-nuclear physics.

The recent view about the solutions of Kähler- Dirac action assumes that the modes have a well-defined em charge and this implies that localization of the modes to 2-D surfaces (right-handed neutrino is an exception). Classical  $W$  boson fields vanish at these surfaces and also classical  $Z^0$  field can vanish. The latter would guarantee the absence of large parity breaking effects above intermediate boson scale scaling like  $h_{eff}$ .

## 1.9 Twistors in TGD and connection with Veneziano duality

The twistorialization of TGD has two aspects. The attempt to generalize twistor Grassmannian approach emerged first. It was however followed by the realization that also the twistor lift of TGD at classical space-time level is needed. It turned out that the progress in the understanding of the classical twistor lift has been much faster - probably this is due to my rather limited technical QFT skills.

### 1.9.1 Twistor lift at space-time level

8-dimensional generalization of ordinary twistors is highly attractive approach to TGD [K23]. The reason is that  $M^4$  and  $CP_2$  are completely exceptional in the sense that they are the only 4-D manifolds allowing twistor space with Kähler structure [A7]. The twistor space of  $M^4 \times CP_2$  is Cartesian product of those of  $M^4$  and  $CP_2$ . The obvious idea is that space-time surfaces allowing twistor structure if they are orientable are representable as surfaces in  $H$  such that the properly induced twistor structure coincides with the twistor structure defined by the induced metric.

In fact, it is enough to generalize the induction of spinor structure to that of twistor structure so that the induced twistor structure need not be identical with the ordinary twistor structure possibly assignable to the space-time surface. The induction procedure reduces to a dimensional reduction of 6-D Kähler action giving rise to 6-D surfaces having bundle structure with twistor sphere as fiber and space-time as base. The twistor sphere of this bundle is imbedded as sphere in the product of twistor spheres of twistor spaces of  $M^4$  and  $CP_2$ .

This condition would define the dynamics, and the original conjecture was that this dynamics is equivalent with the identification of space-time surfaces as preferred extremals of Kähler action. The dynamics of space-time surfaces would be lifted to the dynamics of twistor spaces, which are sphere bundles over space-time surfaces. What is remarkable that the powerful machinery of complex analysis becomes available.

It however turned out that twistor lift of TGD is much more than a mere technical tool. First of all, the dimensionally reduction of 6-D Kähler action contained besides 4-D Kähler action also a volume term having interpretation in terms of cosmological constant. This need not bring anything new, since all known extremals of Kähler action with non-vanishing induced Kähler form are minimal surfaces. There is however a large number of embeddings of twistor sphere of space-time surface to the product of twistor spheres. Cosmological constant has spectrum and depends on length scale, and the proposal is that coupling constant evolution reduces to that for cosmological constant playing the role of cutoff length. That cosmological constant could transform from a mere nuisance to a key element of fundamental physics was something totally new and unexpected.

1. The twistor lift of TGD at space-time level forces to replace 4-D Kähler action with 6-D dimensionally reduced Kähler action for 6-D surface in the 12-D Cartesian product of 6-D twistor spaces of  $M^4$  and  $CP_2$ . The 6-D surface has bundle structure with twistor sphere as fiber and space-time surface as base.

Twistor structure is obtained by inducing the twistor structure of 12-D twistor space using dimensional reduction. The dimensionally reduced 6-D Kähler action is sum of 4-D Kähler action and volume term having interpretation in terms of a dynamical cosmological constant depending on the size scale of space-time surface (or of causal diamond CD in zero energy ontology (ZEO)) and determined by the representation of twistor sphere of space-time surface in the Cartesian product of the twistor spheres of  $M^4$  and  $CP_2$ .

2. The preferred extremal property as a representation of quantum criticality would naturally correspond to minimal surface property meaning that the space-time surface is separately an extremal of both Kähler action and volume term almost everywhere so that there is no coupling between them. This is the case for all known extremals of Kähler action with non-vanishing induced Kähler form.

Minimal surface property could however fail at 2-D string world sheets, their boundaries and perhaps also at partonic 2-surfaces. The failure is realized in minimal sense if the 3-surface has 1-D edges/folds (strings) and 4-surface 2-D edges/folds (string world sheets) at which some partial derivatives of the embedding space coordinates are discontinuous but canonical momentum densities for the entire action are continuous.

There would be no flow of canonical momentum between interior and string world sheet and minimal surface equations would be satisfied for the string world sheet, whose 4-D counterpart in twistor bundle is determined by the analog of 4-D Kähler action. These conditions allow the transfer of canonical momenta between Kähler- and volume degrees of freedom at string world sheets. These no-flow conditions could hold true at least asymptotically (near the boundaries of CD).

$M^8 - H$  duality suggests that string world sheets (partonic 2-surfaces) correspond to images of complex 2-sub-manifolds of  $M^8$  (having tangent (normal) space which is complex 2-plane of octonionic  $M^8$ ).

3. Cosmological constant would depend on p-adic length scales and one ends up to a concrete model for the evolution of cosmological constant as a function of p-adic length scale and other number theoretic parameters (such as Planck constant as the order of Galois group): this conforms with the earlier picture.

Inflation is replaced with its TGD counterpart in which the thickening of cosmic strings to flux tubes leads to a transformation of Kähler magnetic energy to ordinary and dark matter. Since the increase of volume increases volume energy, this leads rapidly to energy minimum at some flux tube thickness. The reduction of cosmological constant by a phase transition however leads to a new expansion phase. These jerks would replace smooth cosmic expansion of GRT. The discrete coupling constant evolution predicted by the number theoretical vision could be understood as being induced by that of cosmological constant taking the role of cutoff parameter in QFT picture [?].

### 1.9.2 Twistor lift at the level of scattering amplitudes and connection with Veneziano duality

The classical part of twistor lift of TGD is rather well-understood. Concerning the twistorialization at the level of scattering amplitudes the situation is much more difficult conceptually - I already mentioned my limited QFT skills.

1. From the classical picture described above it is clear that one should construct the 8-D twistorial counterpart of theory involving space-time surfaces, string world sheets and their boundaries, plus partonic 2-surfaces and that this should lead to concrete expressions for the scattering amplitudes.

The light-like boundaries of string world sheets as carriers of fermion numbers would correspond to twistors as they appear in twistor Grassmann approach and define the analog for the massless sector of string theories. The attempts to understand twistorialization have been restricted to this sector.

2. The beautiful basic prediction would be that particles massless in 8-D sense can be massive in 4-D sense. Also the infrared cutoff problematic in twistor approach emerges naturally and reduces basically to the dynamical cosmological constant provided by classical twistor lift.

One can assign 4-momentum both to the spinor harmonics of the embedding space representing ground states of super-conformal representations and to light-like boundaries of string world sheets at the orbits of partonic 2-surfaces. The two four-momenta should be identical by quantum classical correspondence: this could be seen as a concretization of Equivalence Principle. Also a connection with string model emerges.

3. As far as symmetries are considered, the picture looks rather clear. Ordinary twistor Grassmannian approach boils down to the construction of scattering amplitudes in terms of Yangian invariants for conformal group of  $M^4$ . Therefore a generalization of super-symplectic symmetries to their Yangian counterpart seems necessary. These symmetries would be gigantic but how to deduce their implications?
4. The notion of positive Grassmannian is central in the twistor approach to the scattering amplitudes in  $calN = 4$  SUSYs. TGD provides a possible generalization and number theoretic interpretation of this notion. TGD generalizes the observation that scattering amplitudes in twistor Grassmann approach correspond to representations for permutations. Since 2-vertex is the only fermionic vertex in TGD, OZI rules for fermions generalizes, and scattering amplitudes are representations for braidings.

Braid interpretation encourages the conjecture that non-planar diagrams can be reduced to ordinary ones by a procedure analogous to the construction of braid (knot) invariants by gradual un-braiding (un-knotting).

This is however not the only vision about a solution of non-planarity. Quantum criticality provides different view leading to a totally unexpected connection with string models, actually with the Veneziano duality, which was the starting point of dual resonance model in turn leading via dual resonance models to super string models.

1. Quantum criticality in TGD framework means that coupling constant evolution is discrete in the sense that coupling constants are piecewise constant functions of length scale replaced by dynamical cosmological constant. Loop corrections would vanish identically and the recursion formulas for the scattering amplitudes (allowing only planar diagrams) deduced in twistor Grassmann would involve no loop corrections. In particular, cuts would be replaced by sequences of poles mimicking them like sequences of point charge mimic line charges. In momentum discretization this picture follows automatically.
2. This would make sense in finite measurement resolution realized in number theoretical vision by number-theoretic discretization of the space-time surface (cognitive representation) as points with coordinates in the extension of rationals defining the adèle [?]. Similar discretization would take place for momenta. Loops would vanish at the level of discretization but what would happen at the possibly existing continuum limit: does the sequence of poles integrate to cuts? Or is representation as sum of resonances something much deeper?
3. Maybe it is! The basic idea of behind the original Veneziano amplitudes (see <http://tinyurl.com/yyhwvqb>) was Veneziano duality. This 4-particle amplitude was generalized by Yoshiro Nambu, Holber-Beck Nielsen, and Leonard Susskind to N-particle amplitude (see <http://tinyurl.com/yyvnx7as>) based on string picture, and the resulting model was called dual resonance model. The model was forgotten as QCD emerged. Later came superstring models and led to M-theory. Now it has become clear that something went wrong, and it seems that one must return to the roots. Could the return to the roots mean a careful reconsideration of the dual resonance model?
4. Recall that Veneziano duality (1968) was deduced by assuming that scattering amplitude can be described as sum over s-channel resonances or t-channel Regge exchanges and Veneziano duality stated that hadronic scattering amplitudes have representation as sums over s- or t-channel resonance poles identified as excitations of strings. The sum over exchanges defined by t-channel resonances indeed reduces at larger values of  $s$  to Regge form.

The resonances had zero width, which was not consistent with unitarity. Further, there were no counterparts for the *sum* of s-, t-, and u-channel diagrams with continuous cuts



in the kinematical regions encountered in QFT approach. What puts bells ringing is the u-channel diagrams would be non-planar and non-planarity is the problem of twistor Grassmann approach.

5. Veneziano duality is true only for s- and t- channels but not been s- and u-channel. Stringy description makes t-channel and s-channel pictures equivalent. Could it be that in fundamental description u-channels diagrams cannot be distinguished from s-channel diagrams or t-channel diagrams? Could the stringy representation of the scattering diagrams make u-channel twist somehow trivial if handles of string world sheet representing stringy loops in turn representing the analog of non-planarity of Feynman diagrams are absent? The permutation of external momenta for tree diagram in absence of loops in planar representation would be a twist of  $\pi$  in the representation of planar diagram as string world sheet and would not change the topology of the string world sheet and would not involve non-trivial world sheet topology.

For string world sheets loops would correspond to handles. The presence of handle would give an edge with a loop at the level of 3-surface (self energy correction in QFT). Handles are not allowed if the induced metric for the string world sheet has Minkowskian signature. If the stringy counterparts of loops are absent, also the loops in scattering amplitudes should be absent.

This argument applies only inside the Minkowskian space-time regions. If string world sheets are present also in Euclidian regions, they might have handles and loop corrections could emerge in this manner. In TGD framework strings (string world sheets) are identified to 1-D edges/folds of 3-surface at which minimal surface property and topological QFT property fails (minimal surfaces as calibrations). Could the interpretation of edge/fold as discontinuity of some partial derivatives exclude loopy edges: perhaps the branching points would be too singular?

A reduction to a sum over s-channel resonances is what the vanishing of loops would suggest. Could the presence of string world sheets make possible the vanishing of continuous cuts even at the continuum limit so that continuum cuts would emerge only in the approximation as the density of resonances is high enough?

The replacement of continuous cut with a sum of *infinitely* narrow resonances is certainly an approximation. Could it be that the stringy representation as a sum of resonances with *finite* width is an essential aspect of quantum physics allowing to get rid of infinities necessarily accompanying loops? Consider now the arguments against this idea.

1. How to get rid of the problems with unitarity caused by the zero width of resonances? Could *finite* resonance widths make unitarity possible? Ordinary twistor Grassmannian approach predicts that the virtual momenta are light-like but complex: obviously, the imaginary part of the energy in rest frame would have interpretation as resonance width.

In TGD framework this generalizes for 8-D momenta. By quantum-classical correspondence (QCC) the classical Noether charges are equal to the eigenvalues of the fermionic charges in Cartan algebra (maximal set of mutually commuting observables) and classical TGD indeed predicts complex momenta (Kähler coupling strength is naturally complex). QCC thus supports this proposal.

2. Sum over resonances/exchanges picture is in conflict with QFT picture about scattering of particles. Could *finite* resonance widths due to the complex momenta give rise to the QFT type scattering amplitudes as one develops the amplitudes in Taylor series with respect to the resonance width? Unitarity condition indeed gives the first estimate for the resonance width. QFT amplitudes should emerge in an approximation obtained by replacing the discrete set of finite width resonances with a cut as the distance between poles is shorter than the resolution for mass squared.

In superstring models string tension has single very large value and one cannot obtain QFT type behavior at low energies (for instance, scattering amplitudes in hadronic string model are concentrated in forward direction). TGD however predicts an entire hierarchy of p-adic length scales with varying string tension. The hierarchy of mass scales corresponding roughly to the lengths and thickness of magnetic flux tubes as thickened cosmic strings and characterized by the value of cosmological constant predicted by twistor lift of TGD. Could this give rise

to continuous QCT type cuts at the limit when measurement resolution cannot distinguish between resonances?

The dominating term in the sum over sums of resonances in  $t$ -channel gives near forward direction approximately the lowest mass resonance for strings with the smallest string tension. This gives the behavior  $1/(t - m_{min}^2)$ , where  $m_{min}$  corresponds to the longest mass scale involved (the largest space-time sheet involved), approximating the  $1/t$ -behavior of massless theories. This also brings in IR cutoff, the lack of which is a problem of gauge theories. This should give rise to continuous QFT type cuts at the limit when measurement resolution cannot distinguish between resonances.

## 2 Bird's Eye of View about the Topics of "Physics in Many-sheeted Space-time: Part II"

The key ideas are explained in the first part of the books so that it is enough to explain the organization of the book.

In this book many-sheeted-cosmology and astrophysics are summarized. As in all books in this collection, the material is ordered along the time-line of TGD, and the views of the earliest chapters are not necessarily completely consistent with the views of latest chapters. The book consists of 2 parts.

p-Adic and dark matter hierarchies imply that TGD inspired cosmology has a kind of Russian doll structure containing cosmologies within cosmologies. Cosmic strings and their deformations are basic objects of TGD inspired cosmology and of galactic physics - even astrophysics. They form a fractal hierarchy continuing down to the level of elementary particles representable as closed flux tubes carrying monopole flux. Very early cosmology would be dominated by cosmic strings with 2-D  $M^4$  projection and one can say that the GRT based cosmology with macroscopic space-time identifiable as a small metric deformation of Minkowski space emerged during the analog of inflationary period leading to radiation dominated cosmology. In the chapters devoted to the TGD inspired cosmology the imbeddings of Robertson-Walker cosmology are studied. Both critical and over-critical cosmology are found to be unique apart from the parameter characterizing its duration. It must be emphasized that RW cosmologies make sense as preferred extremals only at the limit of vanishing cosmological constant. Furthermore, the metrics obtained at the QFT limit of TGD need not be embeddable as 4-surfaces in  $M^4 \times CP_2$ . The idea about dark matter hierarchy with levels labeled by the values of Planck constant was originally motivated by the observation that planetary orbits could be interpreted as Bohr orbits with enormous value of Planck constant (introduced originally by Nottale), whose value is fixed to a high degree by Equivalence Principle and which is assignable to the flux tubes mediating gravitational interaction. One ends up to a rather detailed view about macroscopically quantum coherent dark matter in astrophysics and cosmology. In particular, dark matter could be in anyonic phase at light-like 2-surfaces with complex topology and astrophysical size and visible matter would condense around it. Dark matter hierarchy allows interpreting critical cosmologies as correlates for the phase transitions, increasing Planck constant and involving a relatively rapid expansion of space-time sheets. The quantum counterpart of the smooth cosmological expansion would be a series of phase transitions increasing the value of Planck constant and these phase transitions are predicted to take place also at planetary level, which provides a new theoretical basis for Expanding Earth hypothesis and suggests totally unexpected connections between biology and geology. Twistor lift of TGD provides justification for the idea about cosmological constant depending on length scale. It replaces Kähler action with its sum with a volume term having interpretation in terms of dynamical cosmological constant, which depends on length scale and decreases like one over p-adic length scale squared. One can see cosmological evolution as a sequence of phase transitions reducing the value of cosmological constant followed by an expansion of the thickness of flux tubes reducing their Kähler magnetic energy but increasing volume energy until energy minimum is achieved. Twistor lift of TGD predicts that also the  $M^4$  analog of Kähler action is present in the dimensionally reduced 6-D Kähler action. One can understand why the physical effects such as the

violation of CP and T symmetries are small and an explanation for the small CP breaking effects of hadron physics and for the matter antimatter asymmetry emerges. Long cosmic strings would develop galaxies as tangles with topology of dipole magnetic field (roughly) and these flux tubes associated with these tangles would in turn generate stars and planets as similar tangles. Also flux tubes within flux tubes are possible. This hierarchy could continue to even smaller length scales. The thickening of cosmic strings and the sequence of phase transitions reducing cosmological constant would induce transformation of the magnetic and volume energy of cosmic strings to ordinary matter generating the visible matter of galaxies and also stars and planets. The decay of magnetic energy to ordinary matter during the thickening period of the flux tube would be analogous to the decay of the inflaton field. The flat velocity spectrum of stars rotating around galaxies is predicted correctly without need for dark matter halo. A new view about blackholes and white holes as tangles associated with cosmic strings suggests itself. For instance quasars could be white hole-like objects created as Kähler magnetic tangle evolves from straight cosmic string and thickens by decays to ordinary matter. Black hole-like entity would be a time reversal of this process and indeed possible in ZEO predicting that “big” state function reductions inducing reversal of the arrow of time are possible even in astrophysical and cosmological scales.

### 3 Sources

The eight online books about TGD [K25, K24, K17, K12, K4, K11, K6, K20] and nine online books about TGD inspired theory of consciousness and quantum biology [K22, K3, K13, K2, K5, K7, K8, K19, K21] are warmly recommended for the reader willing to get overall view about what is involved.

My homepage (<http://tinyurl.com/ybv8dt4n>) contains a lot of material about TGD. In particular, a TGD glossary at <http://tinyurl.com/yd6j3o7>.

I have published articles about TGD and its applications to consciousness and living matter in *Journal of Non-Locality* (<http://tinyurl.com/ycyrxj4o> founded by Lian Sidorov and in *Prespacetime Journal* (<http://tinyurl.com/ycvktjhn>), *Journal of Consciousness Research and Exploration* (<http://tinyurl.com/yba4f672>), and *DNA Decipher Journal* (<http://tinyurl.com/y9z52khg>), all of them founded by Huping Hu. One can find the list about the articles published at <http://tinyurl.com/ybv8dt4n>. I am grateful for these far-sighted people for providing a communication channel, whose importance one cannot overestimate.

#### 3.1 PART I: MANY-SHEETED COSMOLOGY

##### 3.1.1 Cosmic Strings

Cosmic strings belong to the basic extremals of the Kähler action. The upper bound for string tension of the cosmic strings is  $T \simeq .5 \times 10^{-6}/G$  and in the same range as the string tension of GUT strings and this makes them very interesting cosmologically although TGD cosmic strings have otherwise practically nothing to do with their GUT counterparts.

###### 1. Basic ideas

The understanding of cosmic strings has developed only slowly and has required dramatic modifications of existing views.

1. Zero energy ontology implies that the energy and all quantum numbers of the Universe vanishes and physical states are zero energy states decomposing into pairs of positive and negative energy states localizable to the light-like boundaries of causal diamonds defined as intersections of future and past directed light-cones. Positive energy ontology is a good approximation under certain assumptions.
2. Dark matter hierarchy whose levels are labeled by gigantic values of gravitational Planck constant associated with dark matter is second essential piece of the picture.
3. The second variation of Kähler action vanishes for preferred extremals - at least the second variations associated with dynamical symmetries. This guarantees that Noether currents assignable to the Kähler-Dirac action are conserved. The properties of the preferred extremals

suggest a dimensional reduction providing formulations of quantum TGD in terms of possibly dual slicings of space-time surface by string world sheets and partonic 2-surfaces. The localization of the modes of the Kähler-Dirac equation to 2-D surfaces - string world sheets and possibly partonic 2-surfaces) suggests something similar although it might be that both kind of objects are necessary for a full description.

4. GRT limit of can be understood as an outcome of the replacement of sheets of the many-sheeted space-time with single sheet endowed with effective metric given by the sum of Minkowski metric and deviations of the induced metrics of space-time sheets from Minkowski metric. Gauge theory limit can be understood in an analogous manner. Equivalence Principle in Einsteinian sense follows from Poincare invariance of TGD. The additional assumption made before a real understanding of GRT limit was that the the most important GRT space-times can be represented as vacuum extremals of Kähler action. This assumption can be of course questioned.
5. The basic question whether one can model the exterior region of the topologically condensed cosmic string using General Relativity. The exterior metric of the cosmic string corresponds to a small deformation of a vacuum extremal assuming the identification of the most important GRT space-times as vacuum extremals of Kähler action. The angular defect and surplus associated with the exterior metrics extremizing curvature scalar can be much smaller than assuming vacuum Einstein's equations. The conjecture is that the exterior metric of galactic string conforms with the Newtonian intuitions and thus explains the constant velocity spectrum of distant stars if one assumes that galaxies are organized to linear structures along long strings like pearls in a necklace.

### *2. Critical and over-critical cosmologies involve accelerated cosmic expansion*

In TGD framework critical and over-critical cosmologies are unique apart from single parameter telling their duration and predict the recently discovered accelerated cosmic expansion. Critical cosmologies are naturally associated with quantum critical phase transitions involving the change of gravitational Planck constant. A natural candidate for such a transition is the increase of the size of a large void as galactic strings have been driven to its boundary. During the phase transitions connecting two stationary cosmologies (extremals of curvature scalar) also determined apart from single parameter, accelerated expansion is predicted to occur. These transitions are completely analogous to quantum transitions at atomic level.

The proposed microscopic model predicts that the TGD counterpart of the quantity  $\rho + 3p$  for cosmic strings is negative during the phase transition which implies accelerated expansion. Dark energy is replaced in TGD framework with dark matter indeed predicted by TGD and its fraction is .74 as in standard scenario. Cosmological constant thus characterizes phenomenologically the density of dark matter rather than energy in TGD Universe.

The sizes of large voids stay constant during stationary periods which means that also cosmological constant is piecewise constant. p-Adic length fractality predicts that  $\Lambda$  scales as  $1/L^2(k)$  as a function of the p-adic scale characterizing the space-time sheet of void. The order of magnitude for the recent value of the cosmological constant comes out correctly. The gravitational energy density described by the cosmological constant is identifiable as that associated with topologically condensed cosmic strings and of magnetic flux tubes to which they are gradually transformed during cosmological evolution.

### *3. Cosmic strings and generation of structures*

1. In zero energy ontology cosmic strings must be created from vacuum as zero energy states consisting of pairs of strings with opposite time orientation and inertial energy.
2. The counterpart of Hawking radiation provides a mechanism by which cosmic strings can generate ordinary matter. The splitting of cosmic strings followed by a "burning" of the string ends provides a second manner to generate visible matter. Matter-antimatter symmetry would result if antimatter is inside cosmic strings and matter in the exterior region. A justification for CP asymmetry comes from basic quantum TGD. One can add to Kähler function of the WCW an imaginary part defined by instanton term  $J \wedge J$ . This term does not affect Kähler metric but implies CP breaking.

3. Zero energy ontology has deep implications for the cosmic and ultimately also for biological evolution (magnetic flux tubes play a fundamental role in TGD inspired biology and cosmic strings are limiting cases of them). The arrows of geometric time are opposite for the strings and also for positive energy matter and negative energy antimatter. This implies a competition between two dissipative time developments proceeding in different directions of geometric time and looking self-organization and even self-assembly from the point of view of each other. This resolves paradoxes created by gravitational self-organization contra second law of thermodynamics. So called super-symplectic matter at cosmic strings implies large p-adic entropy resolves the well-known entropy paradox.
4. p-Adic fractality and simple quantitative observations lead to the hypothesis that cosmic strings are responsible for the evolution of astrophysical structures in a very wide length scale range. Large voids with size of order  $10^8$  light years can be seen as structures cosmic strings wound around the boundaries of the void. Galaxies correspond to same structure with smaller size and linked around the supra-galactic strings. This conforms with the finding that galaxies tend to be grouped along linear structures. Simple quantitative estimates show that even stars and planets could be seen as structures formed around cosmic strings of appropriate size. Thus Universe could be seen as fractal cosmic necklace consisting of cosmic strings linked like pearls around longer cosmic strings linked like...

#### 4. Cosmic strings, gamma ray bursts, and supernovae

During year 2003 two important findings related to cosmic strings were made.

1. A correlation between supernovae and gamma ray bursts was observed.
2. Evidence that some unknown particles of mass  $m \simeq 2m_e$  and decaying to gamma rays and/or electron positron pairs annihilating immediately serve as signatures of dark matter. These findings challenge the identification of cosmic strings and/or their decay products as dark matter, and also the idea that gamma ray bursts correspond to cosmic fire crackers formed by the decaying ends of cosmic strings.

This forces the updating of the more than decade old rough vision about topologically condensed cosmic strings and about gamma ray bursts described in this chapter. According to the updated model, cosmic strings transform in topological condensation to magnetic flux tubes about which they represent a limiting case. Primordial magnetic flux tubes forming ferro-magnet like structures become seeds for gravitational condensation leading to the formation of stars and galaxies. The TGD based model for the asymptotic state of a rotating star as dynamo leads to the identification of the predicted magnetic flux tube at the rotation axis of the star as  $Z^0$  magnetic flux tube of primordial origin. Besides  $Z^0$  magnetic flux tube structure also magnetic flux tube structure exists at different space-time sheet but is in general not parallel to the  $Z^0$  magnetic structure. This structure cannot have primordial origin (the magnetic field of star can even flip its polarity).

The flow of matter along  $Z^0$  magnetic (rotation) axis generates synchrotron radiation, which escapes as a precisely targeted beam along magnetic axis and leaves the star. The identification is as the rotating light beam associated with ordinary neutron stars. During the core collapse leading to the supernova this beam becomes gamma ray burst. The mechanism is very much analogous to the squeezing of the tooth paste from the tube. The fact that all nuclei are fully ionized  $Z^0$  ions, the  $Z^0$  charge unbalance caused by the ejection of neutrinos, and the radial compression make the effect extremely strong so that there are hopes to understand the observed incredibly high polarization of  $80 \pm 20$  per cent.

The  $W$  fields experienced by fundamental fermions at 2-D surfaces at which they are localized vanish by the well-definedness of em charge, and one can also require that  $Z^0$  fields vanish at least above weak scale. The vanishing of effective weak fields is an obvious objection against the model unless one allows the possibility of large values of  $h_{eff} = n \times h$  strongly suggested by the identification  $h_{eff} = h_{gr}$ , where  $h_{gr}GMm/v_0$  is the gravitational Planck constant inspired by Nottale's considerations: here  $M$  and  $m$  would correspond to masses of supernova and of microscopic system.

TGD suggests the identification of particles of mass  $m \simeq 2m_e$  accompanying dark matter as lepto-pions formed by color excited leptons, and topologically condensed at magnetic flux tubes having thickness of about lepto-pion Compton length. Lepto-pions would serve as signatures of

dark matter whereas dark matter itself would correspond to the magnetic energy of topologically condensed cosmic strings transformed to magnetic flux tubes.

### 3.1.2 TGD and Cosmology

A proposal for what might be called TGD inspired cosmology is made. The basic ingredient of this cosmology is the TGD counter part of the cosmic string. It is found that many-sheeted space-time concept, the new view about the relationship between inertial and gravitational four-momenta, the basic properties of the cosmic strings, zero energy ontology, the hierarchy of dark matter with levels labeled by arbitrarily large values of Planck constant: the existence of the limiting temperature (as in string model, too), the assumption about the existence of the vapor phase dominated by cosmic strings, and quantum criticality imply a rather detailed picture of the cosmic evolution, which differs from that provided by the standard cosmology in several respects but has also strong resemblances with inflationary scenario.

TGD inspired cosmology in its recent form relies on an ontology differing dramatically from that of GRT based cosmologies. Zero energy ontology (ZEO) states that all physical states have vanishing net quantum numbers so that all matter is creatable from vacuum. The hierarchy of dark matter identified as macroscopic quantum phases labeled by arbitrarily large values of Planck constant is second aspect of the new ontology. The values of the gravitational Planck constant assignable to space-time sheets mediating gravitational interaction are gigantic. This implies that TGD inspired late cosmology might decompose into stationary phases corresponding to stationary quantum states in cosmological scales and critical cosmologies corresponding to quantum transitions changing the value of the gravitational Planck constant and inducing an accelerated cosmic expansion.

#### 1. Zero energy ontology

The construction of quantum theory leads naturally to ZEO stating that everything is creatable from vacuum. Zero energy states decompose into positive and negative energy parts having identification as initial and final states of particle reaction in time scales of perception longer than the geometro-temporal separation  $T$  of positive and negative energy parts of the state. If the time scale of perception is smaller than  $T$ , the usual positive energy ontology applies.

In ZEO inertial four-momentum is a quantity depending on the temporal time scale  $T$  used and in time scales longer than  $T$  the contribution of zero energy states with parameter  $T_1 < T$  to four-momentum vanishes. This scale dependence alone implies that it does not make sense to speak about conservation of inertial four-momentum in cosmological scales. Hence it would be in principle possible to identify inertial and gravitational four-momenta and achieve strong form of Equivalence Principle. It however seems that this is not the correct approach to follow.

#### 2. Dark matter hierarchy and hierarchy of Planck constants

Dark matter revolution with levels of the hierarchy labeled by values of Planck constant forces a further generalization of the notion of imbedding space and thus of space-time. One can say, that embedding space is a book like structure obtained by gluing together infinite number of copies of the embedding space like pages of a book: two copies characterized by singular discrete bundle structure are glued together along 4-dimensional set of common points. These points have physical interpretation in terms of quantum criticality. Particle states belonging to different sectors (pages of the book) can interact via field bodies representing space-time sheets which have parts belonging to two pages of this book. Dark matter hierarchy follows naturally from the non-determinism of Kähler action.

#### 3. Quantum criticality

TGD Universe is quantum counterpart of a statistical system at critical temperature. As a consequence, topological condensate is expected to possess hierarchical, fractal like structure containing topologically condensed 3-surfaces with all possible sizes. Both Kähler magnetized and Kähler electric 3-surfaces ought to be important and string like objects indeed provide a good example of Kähler magnetic structures important in TGD inspired cosmology. In particular space-time is expected to be many-sheeted even at cosmological scales and ordinary cosmology must be replaced with many-sheeted cosmology. The presence of vapor phase consisting of free

cosmic strings containing topologically condensed fermions is second crucial aspect of TGD inspired cosmology.

Quantum criticality of TGD Universe, which corresponds to the vanishing of second variation of Kähler action for preferred extremals - at least of the variations related to dynamical symmetries - supports the view that many-sheeted cosmology is in some sense critical. Criticality in turn suggests fractality. Phase transitions, in particular the topological phase transitions giving rise to new space-time sheets, are (quantum) critical phenomena involving no scales. If the curvature of the 3-space does not vanish, it defines scale: hence the flatness of the cosmic time=constant section of the cosmology implied by the criticality is consistent with the scale invariance of the critical phenomena. This motivates the assumption that the new space-time sheets created in topological phase transitions are in good approximation modelable as critical Robertson-Walker cosmologies for some period of time at least.

These phase transitions are between stationary quantum states having stationary cosmologies as space-time correlates: also these cosmologies are determined uniquely apart from single parameter.

#### *4. Only sub-critical cosmologies are globally imbeddable*

It should be made clear that TGD inspired cosmology involves a vulnerable assumption. It is assumed that single-sheeted space-time surface is enough to model the cosmology. This need not to be the case. GRT limit of TGD is obtained by lumping together the sheets of many-sheeted space-time to a piece of Minkowski space and endowing it with an effective metric, which is sum of Minkowski metric and deviations of the induced metrics of space-time sheets from Minkowski metric. Hence the proposed models make sense only if GRT limits allowing imbedding as a vacuum extremal of Kähler action have special physical role.

TGD allows global imbedding of subcritical cosmologies. A partial imbedding of one-parameter families of critical and overcritical cosmologies is possible. The infinite size of the horizon for the imbeddable critical cosmologies is in accordance with the presence of arbitrarily long range fluctuations at criticality and guarantees the average isotropy of the cosmology. Imbedding is possible for some critical duration of time. The parameter labeling these cosmologies is scale factor characterizing the duration of the critical period. These cosmologies have the same optical properties as inflationary cosmologies. Critical cosmology can be regarded as a “Silent Whisper amplified to Bang” rather than “Big Bang” and transformed to hyperbolic cosmology before its imbedding fails. Split strings decay to elementary particles in this transition and give rise to seeds of galaxies. In some later stage the hyperbolic cosmology can decompose to disjoint 3-surfaces. Thus each sub-cosmology is analogous to biological growth process leading eventually to death.

#### *5. Fractal many-sheeted cosmology*

The critical cosmologies can be used as a building blocks of a fractal cosmology containing cosmologies containing ... cosmologies. p-Adic length scale hypothesis allows a quantitative formulation of the fractality. Fractal cosmology predicts cosmos to have essentially same optic properties as inflationary scenario but avoids the prediction of unknown vacuum energy density. Fractal cosmology explains the paradoxical result that the observed density of the matter is much lower than the critical density associated with the largest space-time sheet of the fractal cosmology. Also the observation that some astrophysical objects seem to be older than the Universe, finds a nice explanation.

#### *6. Cosmic strings as basic building blocks of TGD inspired cosmology*

Cosmic strings are the basic building blocks of TGD inspired cosmology and all structures including large voids, galaxies, stars, and even planets can be seen as pearls in a cosmic fractal necklaces consisting of cosmic strings containing smaller cosmic strings linked around them containing... During cosmological evolution the cosmic strings are transformed to magnetic flux tubes with smaller Kähler string tension and these structures are also key players in TGD inspired quantum biology.

The observed large voids would contain galactic cosmic strings at their boundaries. These voids would participate cosmic expansion only in average sense. During stationary periods the quantum states would be modelable using stationary cosmologies and during phase transitions increasing gravitational Planck constant and thus size of the large void they critical cosmologies would be the appropriate description. The acceleration of cosmic expansion predicted by critical cosmologies

can be naturally assigned with these periods. Classically the quantum phase transition would be induced when galactic strings are driven to the boundary of the large void. The mechanism forcing the phase transition could be repulsive Coulomb energy associated with dark matter at strings if cosmic strings generate net em charge as a consequence of CP breaking (antimatter could reside inside cosmic strings) or a repulsive gravitational acceleration. The large values of Planck constant are crucial for understanding of living matter so that gravitation would play fundamental role also in the evolution of life and intelligence.

Some sections are devoted to the TGD counterpart of inflationary cosmology. From the beginning it has been clear that quantum criticality implying flatness of 3-space and thus criticality is the TGD counterpart for inflationary cosmology. Only after the recent findings about evidence for the polarization of CMB I realized that critical cosmology contains a period of very fast accelerating expansion and that both inflation and accelerating expansion much later are special cases of criticality. This leads to a rather detailed view about how the temperature fluctuations could emerged in TGD framework. The predecessor of inflationary cosmology would be cosmic string gas in the light-cone of Minkowski space and critical period would mean the emergence of space-time as we know it.

### 3.1.3 More about TGD and Cosmology

This chapter can be regarded as second part of the previous chapter and is developed to various applications and problems of cosmology. Much of the text is written decade or two ago.

1. The anomalies of CMB are discussed as a natural continuation of discussion of the counterpart of inflationary cosmology in TGD framework.
2. Simulating Big Bang in laboratory is the title of the next section. The motivation comes from the observation that critical cosmology could serve as a universal model for phase transitions.
3. Some problems of existing cosmology are considered in TGD framework. Discussion includes certain problems of the cosmology such as the questions why some stars seem to be older than the Universe, the claimed time dependence of the fine structure constant, the generation of matter antimatter asymmetry, the problem of the fermion families, and the redshift anomaly of quasars. A mechanism for accelerated expansion of Universe is also considered. In the recent framework this reduces to the critical cosmology and cosmological constant can be assigned to the effective space-time defining GRT limit of TGD.
4. There is a section about matter-antimatter asymmetry, baryogenesis, leptogenesis and TGD discussing whether right-handed neutrino suggested to generate SUSY in TGD framework could be the key entity in fermiogenesis.
5. The remaining sections are devoted to Hogan's theory about quantum fluctuations as new kind of noise and the question whether hyperbolic 3-manifolds emerging naturally in Zero Energy Ontology might be useful in TGD inspired cosmology and explain some redshift anomalies.

### 3.1.4 Breaking of $CP$ , $P$ , and $T$ in cosmological scales in TGD Universe

The twistor lift of TGD forces the analog of Kähler form for  $M^4$ . Covariantly constant self-dual Kähler form  $J(CD)$  depends on causal diamond of  $M^4$  and defines rest frame and spin quantization axis. This implies a violation of  $CP$ ,  $P$ , and  $T$ . By introducing a moduli space for the Kähler forms one avoids the loss of Poincare invariance. The natural question is whether  $J(CD)$  could relate to CP breaking for K and B type mesons, to matter antimatter asymmetry and the large scale parity breaking suggested by CMB data.

The simplest guess for the coupling strength of  $U(1)$  interaction associated with  $J(CD)$  predicts a correct order of magnitude for CP violation for K meson and for the antimatter asymmetry and inspires a more detailed discussion. A general mechanism for the generation of matter asymmetry is proposed, and a model for the formation of disk- and elliptic galaxies is considered. The matter antimatter asymmetry would be apparent in the sense that the CP asymmetry would force matter-antimatter separation: antimatter would reside as dark matter (in TGD sense) inside magnetic flux tubes and matter outside them. Also the angular momenta of dark matter and matter would compensate each other.



### 3.1.5 Comparing Electric Universe hypothesis and TGD

Electric Universe scenario in its extreme form postulates that electromagnetic fields are enough to explain gravitation and even nuclear fusion. From TGD viewpoint this vision is unrealistic. Wes Johnson however gave links to two Youtube videos related to Electric Universe telling about extremely interesting physical findings providing applications for TGD if take seriously. The first video was about the anomalies related to the craters of the Moon and second describe the claimed findings of SAFIRE team having a nice interpretation in TGD framework using the notions of monopole flux tubes and dark matter as hierarchy of phases of ordinary matter with non-standard value of Planck constant implying that electromagnetism has deep implications in arbitrarily long scales. The question in TGD is therefore not about whether electromagnetism (of gauge interactions in general) or gravitation is enough to understand cosmology and astrophysics: both are needed and in the sense of TGD.

## 3.2 PART II: MANY-SHEETED ASTROPHYSICS

### 3.2.1 TGD and Astrophysics

Astrophysics in TGD Universe is the basic topics of this chapter. The topics discussed are following.

1. p-Adic length scale hypothesis can be applied also in astrophysical length scales, and some examples of possible applications are discussed. One of the most interesting implications of p-adicity is the possibility of series of phase transitions changing the value of cosmological constant behaving as  $\Lambda \propto 1/L^2(k)$  as a function of p-adic length scale characterizing the size of the space-time sheet.
2. A model for the solar magnetic field as a bundle of topological magnetic flux tubes is constructed and a model of Sunspot cycle is proposed. This model is also shown to explain the mysteriously high temperature of solar corona and also some other mysterious phenomena related to the solar atmosphere. A direct connection with the TGD based explanation of the dark energy as magnetic and  $Z^0$  magnetic energy of the magnetic flux tubes containing dark matter as ordinary matter, emerges. The matter in the solar corona is simply dark matter leaked from the highly curved portions of the magnetic flux tubes to the space-time sheets where it becomes visible. The generation of anomalous  $Z^0$  charge caused by the runoff of dark neutrinos in Super Nova could provide a first principle explanation for the avoidance of collapse to black-hole in Super Nova explosion.

The recent view about fermions is based on the condition that spinor modes have well-define em charge predicts that induced spinor fields are in the generic case localized to 2-D surfaces at which the classical  $W$  field vanishes as does also  $Z^0$  field above weak scale (proportional to effective Planck constant  $h_{eff}$ ). Hence fermions would feel weak  $Z^0$  field only if they are at space-time sheets with large  $h_{eff}$ .

3. One section is devoted to some astrophysical and cosmological anomalies such as the apparent shrinking of solar system observed by Masreliez, Pioneer anomaly and Flyby anomaly.
4. The astrophysics of solar system involves also an anomaly related to the precession of equinoxes suggesting that Sun might have a companion. TGD suggests a model for anomalies as being due to interaction magnetic flux tube connecting Sun to its companion.
5. The TGD variant of the model of Nottale involved gravitational Planck constant  $h_{gr}$  is discussed in detail. Also further indications for large values of Planck constant are discussed and also the argument that  $h_{gr} = GMm/v_0 = h_{eff} = n \times h$  holds true at least microscopially. If so, the dependence of the effective Planck constant on particle mass can be predicted.

### 3.2.2 Quantum Astrophysics

In this chapter the topics relates to what might be called quantum astrophysics. Motivation comes from the model for Nottale's findings suggesting Bohr quantization of planetary orbits. The model leads to the introduction of gravitational Planck constant  $h_{gr} = GMm/v_0$ , where  $v_0$  corresponds to a typical rotational velocity in two particle system.  $h_{gr}$  characterizes the interaction of masses  $M$  and  $m$  and assigned to the magnetic flux tube connecting them and carrying the massless

extremals mediating gravitational interaction. If  $m$  is planetary mass, the value of  $h_{gr}$  is gigantic. Since gravitational acceleration and gravitational Compton length do not depend on particle mass, one can however assume only that microscopic objects have gravitational flux tube connections to the central mass  $M$ . In this case the values of  $h_{gr}$  could be even identical to the corresponding values of  $h_{eff} = n \times h$  in living matter and  $h_{eff} = h_{gr}$  identification makes sense.

The topics discussed in this chapter are following.

1. An updated view about hierarchy of Planck constants is discussed and the connection  $h_{eff} = h_{gr}$  is shown to be consistent with TGD inspired quantum biology. Quantum gravity would be in key role in biology as intuited also by Penrose.
2. Vision about formation of structures and quantum chaos is astrophysical scales is discussed. Also a speculative view about gravitational radiation based on  $h_{gr}$  is considered.
3. A simple model for gravitational radiation assuming that the emission occurs as dark gravitons is considered.  $h \rightarrow h_{gr}$  implies that the energy of graviton is scaled from that in standard model by  $h_{gr}/h$  factor. Realistic model might correspond to  $h_{gr}$  for microscopic particles. The basic prediction is that gravitons would be detected as bunches of ordinary gravitons.
4. TGD suggests that cosmological evolution involves a series of phase transitions changing the value of  $h_{gr}$  occurring via periods of quantum criticality. The critical cosmology is fixed apart from its duration. This suggests a piecewise accelerated expansion. Also inflationary period would be example of this phenomenon as also accelerating expansion much later.
5. Expanding Earth model has been proposed for long time ago to explain why the continents seem to fit nicely to form a complete covering of the Earth's surface. The model however makes sense if the radius of Earth is one half of its recent value. TGD based interpretation for the expansion is in terms of a phase transition increasing  $h_{gr}$  by factor 2.
6. Blackholes in TGD is the topic of the last two sections.

#### Magnetic Bubbles in TGD Universe: Part I

I received a link to a video summarizing the properties of the Local Bubble surrounding the solar system. The Local Bubble represents only one example of magnetic bubbles. The magnetic bubble carries a magnetic field with field lines along its surface. Star formation and interstellar gas seems to concentrate on the bubble.

It is believed that the Local Bubble has been formed in a burst of star formation in the center of the bubble. These stars would have died as supernovae and the matter from supernova explosions would have pushed gas and compressed it to form the Local Bubble.

These bubbles bring in mind the large voids, whose boundaries carry galaxies. I have discussed this from the TGD point of view already earlier. One ends up with a question, whether galaxies are formed at the surfaces of large voids and stars at the surfaces of the magnetic bubbles. Could also the formation of planets be understood in this way? TGD predicts that cosmic expansion takes place as rapid "jerks" and this view has application to the mystery of Cambrian Explosion. Could these local Big-Bangs give rise to a universal mechanism for the formation of structures? If so, then Earth and Moon must have the same composition. The finding that this is indeed the case, came as a total surprise.

The fusion of dark protons at monopole flux tubes to dark proton sequences identified as dark nuclei, which then transform to ordinary nuclei and liberate nuclei binding energy and in this way induce explosion, is the basic step in the formation of astrophysical objects. Dark fusion was originally proposed as a model of "cold fusion" but later generalized to a model for the first step in the formation of stars not yet involving ordinary fusion. The recently found candidates for population III stars could correspond to these prestellar objects.

Galactic blackholes have been recently found to receive a new contribution to their mass from dark energy identifiable as the energy of cosmic strings in the TGD framework. The second discovery is that galaxies, which should be the oldest ones on the basis of their distance, are oldest ones on the basis of their age: zero energy ontology explains this.

A detailed model emerges for the formation of a planetary system as a series of solar explosions as analogs of supernova explosions throwing out a layer of dark matter transforming to ordinary matter, possibly forming a planet. Both the generalization of Nottale's model for planetary orbits involving gravitational Planck constant and a generalization of the Expanding Earth model are

involved. The model explains the composition differences between giant planets and Earth-like planets and also the Kuiper belt as a failed planet and is also applied to giant exoplanets.

Magnetic Bubbles in TGD Universe: Part II Solar flares involving mass eruptions accompany sunspots and the reversal of the magnetic polarity of the solar magnetic field. The models however have several problems. For instance, it is believed that reconnections of magnetic field lines are essential for the process but the prediction for the rate of the process is by 13-14 orders of magnitude too low. The TGD view of space-time provides a new view of electromagnetic fields based on the notion of a field body. Dark matter as phases of ordinary matter with a large value of effective Planck constant is the second new idea and zero energy ontology (ZEO) provides a third new ingredient.

The recent advances in the understanding of the formation of astrophysical structures in various scales in the TGD framework inspire the attempt to understand the structure of the solar magnetic field and its dynamics involving solar cycle, solar flares, reconnections and reversal of the solar magnetic field. By fractality, the general vision leads to a concrete model for the solar cycle and strongly suggests a concrete analogy of the solar cycle with the basic rhythms appearing in biological systems and the identification of the counterparts of anabolism and catabolism at the fundamental level.

The general picture also leads to a model for the reversals of the Earth's magnetic field and to interesting speculations concerning their connection with the evolutionary leaps. In zero energy ontology, the reversal involves the decay and re-organization of the magnetic body in zero energy ontology. The decay is analogous to the decay of the biological body after death and induces it. This interpretation provides an understanding of the so-called Tuktam phenomenon.

Some Solar Mysteries

This article was inspired by the article "Is the Sun a Black Hole?" by Nassim Hamein. The article describes a collection of various anomalies related to the physics of the Sun, which I have also considered from the TGD point of view. The most important anomalies are the gamma ray anomalies and the missing nuclear matter of about 1500 Earth masses. There is also evidence that the solar surface contains a solid layer: something totally implausible in the standard atomic physics. The idea that the Sun could contain a blackhole led in the TGD framework to a refinement of the earlier model for blackhole-like objects (BHs) as maximally dense flux tube spaghettis predicting also their mass spectrum in terms of Mersenne primes and their Gaussian counterparts. The fact that the masses of the Earth and Sun belong to the spectrum, puts bells ringing.

It however turned out that the TGD based model for the missing nuclear matter assigns the gamma ray anomalies to a magnetic bubble as a layer covering the surface of the Sun and consisting of closed monopole flux tube loops running in North-South direction and carrying  $M_{89}$  nucleons with a mass which is 512 times the mass of the ordinary nucleon. This structure could be seen as a 2-D surface variant of the TGD counterpart of blackhole and under very natural assumptions its mass is the missing 1500 Earth masses of ordinary nuclear matter. This model conforms with the earlier model of the sunspot activity related to the reversal of the solar magnetic field. It also explains the gamma ray anomaly below 35 GeV.

A possible explanation for the TeV anomaly is in terms of  $M_{79}$  nuclei generated in the TGD counterpart for the formation of quark gluon plasma, which in the TGD Universe would generate  $M_{89}$  hadrons from  $M_{107}$  hadrons. Now  $M_{79}$  nuclei would be generated from  $M_{89}$  hadrons in a process analogous to high energy nuclear collision, which would correspond to the collision of the  $M_{89}$  flux tubes, whose distance would be larger than 2 Compton lengths of  $M_{89}$  nucleons.

### 3.2.3 TGD View about Quasars

The work of Rudolph Schild and his colleagues Darryl Letier and Stanley Robertson (among others) suggests that quasars are not supermassive blackholes but something else - MECOs, magnetic eternally collapsing objects having no horizon and possessing magnetic moment. Schild et al argue that the same applies to galactic blackhole candidates and active galactic nuclei, perhaps even to ordinary blackholes as Abhas Mitra, the developer of the notion of MECO proposes.

In the sequel TGD inspired view about quasars relying on the general model for how galaxies are generated as the energy of thickened cosmic strings decays to ordinary matter is proposed. Quasars would not be blackhole like objects but would serve as an analog of the decay of

inflaton field producing the galactic matter. The energy of the string like object would replace galactic dark matter and automatically predict a flat velocity spectrum.

TGD is assumed to have standard model and GRT as QFT limit in long length scales. Could MECOs provide this limit? It seems that the answer is negative: MECOs represent still collapsing objects. The energy of inflaton field is replaced with the sum of the magnetic energy of cosmic string and positive volume energy, which both decrease as the thickness of flux tube increases. The liberated energy transforms to ordinary particles and their dark variants in TGD sense. Time reversal of blackhole would be more appropriate interpretation. One can of course ask, whether the blackhole candidates in galactic nuclei are time reversals of quasars in TGD sense.

The writing of the article led also to a considerable understanding of two key aspects of TGD. The understanding of twistor lift and p-adic evolution of cosmological constant improved considerably. Also the understanding of gravitational Planck constant and the notion of space-time as a covering space became much more detailed in turn allowing much more refined view about the anatomy of magnetic body.

### 3.2.4 Cosmic string model for the formation of galaxies and stars

The view about the role of new nuclear physics predicted by TGD in the model of solar interior gives excellent guidelines for attempts to develop a more detailed understanding about TGD counterparts of blackholes as volume filling flux tube tangles. One ends up to rather detailed picture making correct predictions about minimum radii of blackholes and neutron stars. The idea about ordinary stars as blackhole like objects emerges.

The standard blackhole thermodynamics is replaced by two thermodynamics. The first thermodynamics is assignable to the flux tubes as string like entities having Hagedorn temperature  $T_H$  as maximal temperature. The second thermodynamics is assignable to gravitational flux tubes characterized by the gravitational Planck constant  $h_{gr}$ : Hawking temperature  $T_B$  is scaled up by the ratio  $h_{gr}/\hbar$  to  $T_{B,D}$  and is gigantic as compared to the ordinary Hawking temperature but the intensity of dark Hawking radiation is extremely low.

The condition  $T_H = T_{B,D}$  for thermodynamical equilibrium fixes the velocity parameter  $\beta_0 = v_0/c$  appearing in the Nottale formula for  $h_{gr}$  and suggests  $\beta_0 = 1/h_{eff}$  for the dark nuclei at flux tubes defining star as blackhole like entity in TGD sense. This also predicts the Hagedorn temperature of the counterpart of blackhole in GRT sense to be hadronic Hagedorn temperature assignable to the flux tube containing dark nuclei as dark nucleon sequences so that there is a remarkable internal consistency. In zero energy ontology (ZEO) quasars and galactic blackholes can be seen as time reversals of each other.

The cosmological time anomalies such as stars older than the Universe can be understood. In ZEO the time evolution for the zero energy states associated with causal diamonds (CDs) by sequences of small state function reductions (weak measurements) gives rise to conscious entity, self. Self dies and re-incarnates with an opposite arrow of time in big (ordinary) state function reduction reversing the arrow of time. These reincarnations define kind of universal Karma's cycle. If the Karma's cycle leaves the sizes of CDs bounded and their position in  $M^4$  unaffected, quantum dynamics reduces to a local dynamics inside CDs defining sub-cosmologies. In particular, the age distributions and properties of stars depend only weakly on the value of cosmic time - stars older than the Universe become possible in standard view about time.

The flux tube picture about galaxies and larger structures is discussed with application to some anomalies strongly suggesting the presence of coherence in scales of even billion light years. Also "too" fast spinning galaxies are discussed. The local galaxy supercluster Laniakea is discussed in the flux tube picture as a flux tube tangle in scale of .5 Gly.

### 3.2.5 Could solar system be modelled as a miniature version of spiral galaxy?

The fractality of the TGD Universe motivates a model for planetary systems as miniature version of the model of spiral galaxy. The first two key elements are many-sheeted space-time, the notion of magnetic flux tubes - both monopole flux tubes and gravitational flux tubes without monopole flux - and the identification of dark matter as phases of ordinary matter labelled by effective Planck constant  $h_{eff} = n \times h_0$  ( $h = 6h_0$  is a good guess). Also the TGD generalization of Nottale's model for planetary system as analog of Bohr atom is in key role. A further key aspect is the prediction of

twistor lift of TGD that cosmological constant is length scale dependent and characterizes various systems.

I did not originally end up with this model from general considerations. The first input were the problems related to the collision and accretion models for the formation of planets - TGD could replace these with quantal model. The discovery of "too" heavy blackholes and neutron stars by LIGO suggesting that TGD view about the formation of also planets could provide understanding about the role of angular momentum.

There are also problems related to the understanding of the entire planetary system: the dramatic difference between terrestrial and giant planets is not really understood. The problematic aspects of the Bohr orbit model together with the poorly understood differences between terrestrial and giant planets lead to a proposal that phase transition increasing the  $\hbar_{gr}$  by factor 5 and accompanying a transition reducing the length scale dependent cosmological constant  $\Lambda$  could have scaled up the orbital radii of former inner planets. The transition could have also scaled up the radii of the former inner planets so that they became giant planets.

### 3.2.6 TGD View of the Engine Powering Jets from Active Galactic Nuclei

The identification of the energy source (central engine) explaining the energy loss associated with the jets from active galactic nuclei (AGNs) is a long-standing problem of astrophysics. In the model of Blandford and Znajek (BZ model) for the central engine as a blackhole, the Penrose process would provide the energy. The energy would come basically from the blackhole mass.

Empirical support for the BZ model emerges from the study of the supermassive blackhole associated with a galaxy known as Messier 87 (M87). The finding is that the magnetic field associated with the jet structure is tightly wound helical structure and so strong that it would control the dynamics of the matter from falling to blackhole except by occasional leakages. Electron-positron pairs created in the annihilation of photons would accelerate in the force-free helical electromagnetic field having also an electric component.

The TGD based model involves several aspects of the new physics predicted by TDG. TGD leads to a model of galaxies and other astrophysical structures. Inflation decay is replaced with the thickening of cosmic strings to flux tubes liberating as ordinary matter. Hierarchy of Planck constants  $h_{eff} = nh_0$ , in particular Nottale's hypothesis predicts quantum coherence in the exterior of in scales at least of order Schwarzschild radius of the blackhole-like entity. Zero energy ontology (ZEO) predicts that the arrow of time changes in ordinary state function reductions. TGD replaces black-holes with blackhole-like entities (BHs) and white-holes with their time reversals (WHs) allowed in ZEO.

BH (WH) would be a volume filling flux tube but with a relatively small value of  $h_{eff}$ . In the case of WH, it would provide "metabolic energy" for jets and take care that the value of  $h_{eff}$  is preserved (the analogy with living systems is very strong). The jets would be analogous to laser beams/supracurrents with a huge value of  $h_{eff} = h_{gr}$ . The model would also explain the ultrahigh energy cosmic rays. The force-free fields would be generalized Beltrami fields associated with flux tubes and identifiable as minimal surfaces in the Minkowskian regions of space-time surface. The absence of classical dissipation would be a correlate for the absence of dissipation for supra-currents and dark photon laser beams.

### 3.2.7 Solar Metallicity Problem from TGD Perspective

For ten years ago it was thought that Sun is a well-understood system but more precise computations demonstrated a problem. The metallicities deduced from the spectroscopic data deviate strongly from those deduced from helio-seismology and solar neutrino data.

The abundances used are determined from meteorites and these estimates are more accurate and consistent with the values determined by Asplund et al using 3-D modelling of solar surface used also to extrapolate the metallicities in core.

1. The metallicity of Sun deduced from spectroscopy by Asplund et al would be 1.3 per cent whereas the older model and also helio-seismology give 1.8 per cent metallicity. Is the metallicity indeed 1.3 per cent using standard model to extrapolate the spectroscopic data at surface? Or is it 1.8 per cent deeper in the interior in which case the extrapolation used to deduce metallicity in the interior would not be realistic.

2. There are also other discrepancies. The height of convective zone at which radiative energy transfer is replaced with convection is given by  $R_{CZ} = .724R$ . The predicted He abundance at surface is  $Y_{surf} = .231$ . These values are in conflict with  $R_{CZ} = .713R$  and  $Y_{surf} = .248$  deduced from helio-seismological data. Also density and sound velocity profiles deviate from those deduced from the helio-seismology. Ironically, the earlier model approximating solar surface as 2-D structure is in excellent accordance with the helio-seismological data.

When one has a paradox one must challenge the basic assumptions. Do the metallicities outside Sun and inside solar core really mean same thing? Dark matter identified as  $h_{eff} = nh_0$  phases has become key player in TGD inspired new physics being now a crucial element of TGD based view about living matter. Dark nuclear fusion is proposed to provide the new physics allowing to understand “cold fusion”. In the following it will be found that dark matter in TGD associated with solar core could provide an elegant solution also to the solar metallicity problem.

In TGD classical physics is an exact part of quantum physics. The tunnelling phenomenon essential for nuclear physics based model of solar nuclear fusion would correspond in TGD to a state function reduction creating a phase consisting of dark nuclei which can fuse without tunnelling due to the reduction of the binding energy scale. State function reduction to ordinary phase leads to the final state of the reaction. In ZEO “big” (ordinary) state function reduction (BFSR) would reverse the arrow of time so that if tunnelling phenomenon is assignable to BFSR rather than “small” state function reduction (SFSR) as TGD counterpart of “weak” measurement, ZEO would make possible nuclear fusion.

The missing nuclear matter inside core would be dark variants of nuclei associated with dark flux tubes. This would explain the conflict between the metallicities deduced from spectroscopic and meteoritic data on one hand and those derived from helio-seismic data. The reason is that sound waves and photons in the core couple to both ordinary and dark matter so that helio-seismology gives metallicities as sums of ordinary and dark metallicities. Using the estimate for the thickness of the dark flux tube coming from the TGD based model of “cold fusion”, one can estimate the length of dark flux tube inside solar core and it turns out to fill about 30 per cent of its volume.

One can relate the model also to the model for the formation of galaxies, stars, and planets as tangles assignable to cosmic strings thickened to flux tubes implying the decay of their Kähler magnetic energy to ordinary matter in analogy with the decay of inflaton field and nice quantitative estimates follow. Also a connection with twistor lift of TGD predicting hierarchy of cosmological constants emerges and the radius of solar core turns out to corresponds to the value of cosmological constant implied by the amount of missing matter identified as dark matter at flux tubes.

The view about the role of new nuclear physics predicted by TGD in the model of solar interior gives excellent guidelines for attempts to develop a more detailed understanding about TGD counterparts of blackholes as volume filling flux tube tangles. One ends up to rather detailed picture making correct predictions about minimum radii of blackholes and neutron stars. The idea about ordinary stars as blackhole like objects emerges.

The standard blackhole thermodynamics is replaced by two thermodynamics. The first thermodynamics is assignable to the flux tubes as string like entities having Hagedorn temperature  $T_H$  as maximal temperature. The second thermodynamics is assignable to gravitational flux tubes characterized by the gravitational Planck constant  $h_{gr}$ : Hawking temperature  $T_B$  is scaled up by the ratio  $h_{gr}/\hbar$  to  $T_{B,D}$  and is gigantic as compared to the ordinary Hawking temperature but the intensity of dark Hawking radiation is extremely low. The condition  $T_H = T_{B,D}$  for thermodynamical equilibrium fixes the velocity parameter  $\beta_0 = v_0/c$  appearing in the Nottale formula for  $h_{gr}$  and suggests  $\beta_0 = 1/h_{eff}$  for the dark nuclei at flux tubes defining star as blackhole like entity in TGD sense. This also predicts the Hagedorn temperature of the counterpart of blackhole in GRT sense to be hadronic Hagedorn temperature assignable to the flux tube containing dark nuclei as dark nucleon sequences so that there is a remarkable internal consistency. In zero energy ontology (ZEO) quasars and galactic blackholes can be seen as time reversals of each other.

The flux tube picture about galaxies and larger structures is discussed with application to some anomalies strongly suggesting the presence of coherence in scales of even billion light years. Also “too” fast spinning galaxies are discussed. The local galaxy supercluster Laniakea is discussed in the flux tube picture as a flux tube tangle in scale of .5 Gly.

The flux tube picture about galaxies and larger structures is discussed with application to some

anomalies strongly suggesting the presence of coherence in scales of even billion light years. Also “too” fast spinning galaxies are discussed. The local galaxy supercluster Laniakea is discussed in the flux tube picture as a flux tube tangle in scale of .5 Gly.

### 3.2.8 Solar Surprise

The detection of gamma rays from Sun has yielded a surprises. There are 5 times more gamma rays than expected and the spectrum has a deep and narrow dip around 30-50 GeV. Spectrum continues to much higher energies than expected, at least up to 100 GeV. One proposal is that there could be dark matter in the interior of Sun yielding the gamma rays but is unclear how they could get to the surface without experiencing the same fate as ordinary gammas from nuclear reactions.

The findings provide a test bench for TGD based view about magnetic fields and the first challenge is to understand the solar cycle. The model follows from the model for the formation of galaxies, stars, and planets as tangles of long cosmic strings thickened to flux tube. Wormhole magnetic fields correspond to closed flux tubes with monopole flux returning along different sheet. If  $M^4$  projections of the sheets co-incide and test particle touching them experiences no net magnetic force but the energy of flux tubes is dark making itself visible through gravitational fields. For disjoint projections sheets carry measurable magnetic fields.

Polarization reversal could be understood as a quantum analog of spontaneous magnetization generating first dipole loops of type II (I) taking measured  $B$  to zero. After this dipole loops of type I (II) would split by reconnection and decay to smaller loops and leave Sun. This defines first half-cycle and for second half-cycle the roles of loops are changed.

The model discussed explains qualitatively the findings in terms of cosmic rays entering to the flux tubes of dipole fields and accelerated in the electric field of the closed flux tube and making possibly several cycles before being detected. This predicts band structure of the spectrum.

The model suggests also inversion as a  $Z_2$  symmetry changing the roles of the flux tube portions in the interior and exterior of the solar surface. Inversion symmetry is also a symmetry of Maxwell's equations. The notions of of monopole flux tube and associated approximate  $Z_2$  symmetry acting either as reflection or inversion could be universal.  $Z_2$  can be also represented as a subgroup of the group of Galois symmetries predicted by adelic physics.

This picture leads to highly non-trivial predictions. For instance, the “Axis of Evil” anomaly of CMB can be understood. For instance, quantum correlations in cosmological scales explain why the plane of planetary system makes itself visible in CMB. One can also add highly non-trivial detail to the TGD inspired view about quantum biology and consciousness.

### 3.2.9 LIGO and TGD

The recent detection of gravitational radiation from a merger of blackholes by LIGO detector initiated a new era in astronomy. The detection allows to sharpen the TGD based view about gravitational radiation, in particular to test the proposal that gravitons propagate as dark gravitons with very large value of Planck constant along magnetic flux tubes. Since classical (no dependence of  $h_{eff}$ ) detection of gravitational waves rather than direct detection of gravitons is in question, it is not too surprising that the TGD picture survives. Also a gamma ray burst was observed .4 seconds after the merger and is very probably associated with it. In TGD framework the natural proposal is that this burst arrived as dark cyclotron radiation along the dark flux tubes carrying also the dark gravitons. The energy conserving transformation of the ordinary cyclotron radiation created in the ultra-strong magnetic field of the blackhole to dark photons could have generated the gamma ray pulse. The hypothesis allows to estimate the strength of magnetic field at magnetic flux tubes. The value is consistent with the order of magnitude for intergalactic magnetic fields.

### 3.2.10 About the Recent TGD Based View Concerning Cosmology and Astrophysics

This article is about various topics related to cosmology and to the physics of galaxies, stars and planets and was inspired by several inputs. The first section is about primordial cosmology and describes the TGD counterpart of inflation. The proposal is that the fluctuations of CMB background can be understood number-theoretically as being induced by the fluctuations of the effective Planck constant  $h_{eff} = nh_0$  around  $h_{eff} = h = n_0h_0$ . This also suggests a solution of

the problem posed by two different values of Hubble constants in terms of especially large local upwards fluctuation of the value of  $h_{eff}$  from  $h_{eff} = h$ .

The second section is about several important aspects of TGD inspired cosmology. The findings of JWST force us to ask which came first: supermassive blackholes of galaxies. The recently discovered weak lensing effects lend support for the very long cosmic strings, which represent a key notion in TGD inspired cosmology. Besides dark matter there is also the problem of missing baryonic matter: for some reason 30 per cent of baryons are missing. Furthermore, the quite recent finding of JWST related to supermassive blackhole challenges the GRT based notion of blackhole.

The third section is about the recent TGD view of the physics of stars and planets. The stimulus came from the discovery of a planet that should not exist: the planet has the mass scale of Neptune but the mass of the star is 1/9:th of the solar mass. TGD based model for the formation of galaxies, stars and planets is based on the notion of cosmic strings which produce monopole flux tubes and explanation for the finding and leads to considerably more detailed model for the evolution of stars making a rather dramatic prediction: the element abundances should depend only weakly on cosmic time: the first support for this prediction came already 20 years ago and JWST has provides additional support for it.

In the last section a model for planets and stars as gravitational oscillators inspired by the TGD variant of Nottale's proposal is discussed. It turns out that the radius of the core of Earth corresponds to the Bohr radius for the first orbital, which suggests that the core of Earth, and more generally of the inner planets and Mars corresponds to an S-wave ground state. For the Sun the  $n = 1$  S-wave orbital is 1.5 times the solar radius. The model applies also to the outer planets. Also the rings of giant planets can be understood at a rough quantitative level.

In the last section a cosmic string model for the spiral galaxies predicting the branched spiral structure is considered. Also a quantum model for the galactic blackhole-like objects allowing to understand the weird properties of Sagittarius A\* is discussed.

## REFERENCES

### Mathematics

- [A1] Yangian symmetry. Available at: <https://en.wikipedia.org/wiki/Yangian>.
- [A2] Pope CN. Eigenfunctions and  $Spin^c$  Structures on  $CP_2$ , 1980.
- [A3] Hanson J Eguchi T, Gilkey B. *Phys Rep*, 66, 1980.
- [A4] Eisenhart. *Riemannian Geometry*. Princeton University Press, 1964.
- [A5] Pope CN Gibbons GW.  $CP_2$  as gravitational instanton. *Comm Math Phys*, 55, 1977.
- [A6] Pope CN Hawking SW. Generalized Spin Structures in Quantum Gravity. *Phys Lett*, (1), 1978.
- [A7] N. Hitchin. Kählerian twistor spaces. *Proc London Math Soc*, 8(43):133–151, 1981.. Available at: <https://tinyurl.com/pb8zpqo>.
- [A8] Milnor J. *Topology form Differential Point of View*. The University Press of Virginia, Virginia, 1965.
- [A9] Spivak M. *Differential Geometry I,II,III,IV*. Publish or Perish, Boston, 1970.
- [A10] Thom R. *Comm Math Helvet*, 28, 1954.
- [A11] Wallace. *Differential Topology*. W. A. Benjamin, New York, 1968.

### Theoretical Physics

- [B1] Rapoport D. Stochastic processes in conformal Riemann-Cartan-Weyl gravitation, 1991. Available at: <https://link.springer.com/article/10.1007/BF00675614>.
- [B2] Witten E Dolan L, Nappi CR. Yangian Symmetry in  $D = 4$  superconformal Yang-Mills theory, 2004. Available at: <https://arxiv.org/abs/hep-th/0401243>.



- [B3] Plefka J Drummond J, Henn J. Yangian symmetry of scattering amplitudes in  $\mathcal{N} = 4$  super Yang-Mills theory, 2009. Available at: <https://cdsweb.cern.ch/record/1162372/files/jhep052009046.pdf>.
- [B4] Arkani-Hamed N et al. The All-Loop Integrand For Scattering Amplitudes in Planar N=4 SYM, 2010. Available at: <https://arxiv.org/abs/1008.2958>.

## Cosmology and Astro-Physics

- [E1] Nottale L Da Rocha D. Gravitational Structure Formation in Scale Relativity, 2003. Available at: <https://arxiv.org/abs/astro-ph/0310036>.

## Books related to TGD

- [K1] Pitkänen M. *Topological Geometrostatics*. 1983. Thesis in Helsinki University 1983.
- [K2] Pitkänen M. *Bio-Systems as Conscious Holograms*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/holography.html>, 2023.
- [K3] Pitkänen M. *Bio-Systems as Self-Organizing Quantum Systems*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/bioselforg.html>, 2023.
- [K4] Pitkänen M. Classical TGD. In *Topological Geometrostatics: Overview: Part I:https://tgdtheory.fi/tgdhtml/Btgdview1.html*. Available at: <https://tgdtheory.fi/pdfpool/tgdclass.pdf>, 2023.
- [K5] Pitkänen M. *Genes and Memes*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/genememe.html>, 2023.
- [K6] Pitkänen M. *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/neuplanck.html>, 2023.
- [K7] Pitkänen M. *Magnetospheric Consciousness*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/magnconsc.html>, 2023.
- [K8] Pitkänen M. *Mathematical Aspects of Consciousness Theory*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/mathconsc.html>, 2023.
- [K9] Pitkänen M. Negentropy Maximization Principle. In *TGD Inspired Theory of Consciousness: Part I. https://tgdtheory.fi/tgdhtml/Btgdconsc1.html*. Available at: <https://tgdtheory.fi/pdfpool/nmpc.pdf>, 2023.
- [K10] Pitkänen M. Number theoretic vision, Hyper-finite Factors and S-matrix. In *Quantum TGD: Part I. https://tgdtheory.fi/tgdhtml/Btgdquantum1.html*. Available at: <https://tgdtheory.fi/pdfpool/UandM.pdf>, 2023.
- [K11] Pitkänen M. *p-Adic length Scale Hypothesis*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/padphys.html>, 2023.
- [K12] Pitkänen M. Physics as a Generalized Number Theory. In *Topological Geometrostatics: Overview: Part I. https://tgdtheory.fi/tgdhtml/Btgdview1.html*. Available at: <https://tgdtheory.fi/pdfpool/tgdnumber.pdf>, 2023.
- [K13] Pitkänen M. *Quantum Hardware of Living Matter*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/bioware.html>, 2023.
- [K14] Pitkänen M. Quantum Mind and Neuroscience. In *TGD and EEG: Part I. https://tgdtheory.fi/tgdhtml/Btgd EEG1.html*. Available at: <https://tgdtheory.fi/pdfpool/lianPN.pdf>, 2023.
- [K15] Pitkänen M. Quantum Model for Bio-Superconductivity: I. In *TGD and Quantum Biology: Part I. https://tgdtheory.fi/tgdhtml/Bqbio1.html*. Available at: <https://tgdtheory.fi/pdfpool/biosupercondI.pdf>, 2023.
- [K16] Pitkänen M. Quantum Model for Bio-Superconductivity: II. In *TGD and Quantum Biology: Part I. https://tgdtheory.fi/tgdhtml/Bqbio1.html*. Available at: <https://tgdtheory.fi/pdfpool/biosupercondII.pdf>, 2023.

- [K17] Pitkänen M. *Quantum TGD*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/tgdquantum.html>, 2023.
- [K18] Pitkänen M. TGD and Astrophysics. In *Physics in Many-Sheeted Space-Time: Part II*. <https://tgdtheory.fi/tgdhtml/Btgdclass2.html>. Available at: <https://tgdtheory.fi/pdfpool/astro.pdf>, 2023.
- [K19] Pitkänen M. *TGD and EEG*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/tgdeeg.html>, 2023.
- [K20] Pitkänen M. *TGD and Fringe Physics*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/freenergy.html>, 2023.
- [K21] Pitkänen M. *TGD Based View About Living Matter and Remote Mental Interactions*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/tgdlian.html>, 2023.
- [K22] Pitkänen M. *TGD Inspired Theory of Consciousness*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/tgdconsc.html>, 2023.
- [K23] Pitkänen M. The classical part of the twistor story. In *Quantum TGD: Part III*. <https://tgdtheory.fi/tgdhtml/Btgdquantum3.html>. Available at: <https://tgdtheory.fi/pdfpool/twistorstory.pdf>, 2023.
- [K24] Pitkänen M. The Geometry of the World of the Classical Worlds. In *Topological Geometro-dynamics: Overview: Part I*:<https://tgdtheory.fi/tgdhtml/Btgdview1.html>. Available at: <https://tgdtheory.fi/pdfpool/tgdgeom.pdf>, 2023.
- [K25] Pitkänen M. *Topological Geometro-dynamics: an Overview*. Online book. Available at: <https://www.tgdtheory.fi/tgdhtml/tgdview.html>, 2023.