

Expanding Earth Hypothesis and Pre-Cambrian Earth

June 29, 2025

Matti Pitkänen

orcid:0000-0002-8051-4364.

email: matpitka6@gmail.com,

url: http://tgdtheory.com/public_html/,

address: Valtatie 8, as. 2, 03600, Karkkila, Finland.

Contents

1	Introduction	2
1.1	Vision of the evolution of life on pre-Cambrian Earth	3
1.2	A model for the phase transition increasing the radius of Earth by factor 2	4
2	A possible view of the expansion of Earth?	5
2.1	Did the inner core participate the expansion?	5
2.2	How the outer core was formed?	5
2.3	The energetics of the transition	6
3	TGD view of the pre-Cambrian era	7
3.1	A rough overall view of pre-Cambrian era	7
3.1.1	Energetics during the pre-Cambrian era	7
3.1.2	The atmosphere of Earth before CE	8
3.1.3	Maybe deep oceans are not necessary?	8
3.2	Comparison with the recent Mars	9
3.3	Some questions related to the pre-Cambrian period	10
3.3.1	Why so few multicellular fossils from pre-Cambrian period?	10
3.3.2	Questions related to energetics	11
3.3.3	Did large oceans really exist?	11
3.3.4	Snowball Earth hypothesis from TGD point of view	11
3.3.5	The Great Oxygenation Event	12
3.3.6	Was CE a quantum phase transition in Earth scale?	12
3.3.7	Oxygenation of oceans did not precede Cambrian Explosion as TGD strongly suggests and empirical study concludes	13
3.3.8	Strange correlation between the strength of the Earth's magnetic field and the atmospheric oxygen content	14
3.3.9	Anomalously high rates of tectonic plate motion and the TGD view of Cambrian explosion	15
3.3.10	Does the existence of an underground ocean floor at the mantle-core boundary relate to underground life?	16
3.3.11	What are the mysterious structures observed in the lower mantle?	18

3.4	A possible mechanism of radiative energy transfer from the Earth's core to underground oceans near the surface of Earth	20
3.4.1	Ultralow velocity zones	20
3.4.2	Ordinary water cannot survive near mantle	21
3.4.3	Pollack effect for superionic water and metabolic energy feed from the core of Earth	21
3.4.4	A mechanism of photosynthesis which does not involve biomolecules	22
3.5	Did Earth have a ring before the Cambrian Explosion and did the rapidly expanding Earth catch the ring?	23
3.6	Could the proposed Expansion of Earth during Cambrian Explosion have lead to the formation of Moon?	23
4	Some mysteries of the biological evolution from the TGD point of view	26
4.1	Three mysteries	26
4.1.1	The mystery of life's origin deepens	26
4.1.2	The mystery of Cambrian Explosion	27
4.1.3	Perplexing findings about asteroid Ryugu	27
4.2	LUCA from the TGD perspective	27
4.2.1	Some basic facts about evolution	28
4.2.2	The number of genes for LUCA is upper bound for the genes of archaea: what does this mean?	28
4.2.3	Two mysteries with a common solution: Cambrian Explosion and the complexity of LUCA	28
4.3	The perplexing findings about the asteroid Ryugu from the TGD perspective	29
4.3.1	Some facts about Ryugu	29
4.3.2	Objections against the Panspermia hypothesis as explanation of the findings about Ryugu	30
4.3.3	The TGD view of Ryugu	30

Abstract

In this article I continue to develop a more detailed TGD version of the Expanding Earth hypothesis explaining Cambrian Explosion (CE). A more detailed view of the pre-Cambrian biology, geology, and thermal evolution emerges and one can relate it to the standard view. This involves topics like faint Sun paradox, the mechanism of Great Oxygenation Event, understanding the TGD counterparts of supercontinents Rodinia and Pannotia preceding CE, snowball Earth, and CE that led to a sudden emergence of highly advanced multicellulars.

Also a more detailed view of what happened in the Cambrian explosion induced by the increase of the radius of Earth by factor 2 emerges (in the TGD Universe, a smooth continuous cosmological expansion is replaced with a sequence of short lasting and fast expansions). One ends up with a detailed model for the phase transition leading to the increase of the Earth radius. This phase transition requires a considerable energy feed provided by the phase transition thickening monopole flux tubes of the magnetic body of Earth and liberating energy. In analogy with the recent Mars, pre-Cambrian Earth had a solid core analogous to the inner core. In the phase transition to a liquid outer core with much larger volume. Part of the newly formed outer core could in turn have transformed to form a part of the mantle increasing its thickness.

Contents

1 Introduction

In this article I continue to develop the TGD version of the Expanding Earth hypothesis [L3]. More detailed views of the pre-Cambrian biology, geology, and thermal evolution of Earth and of what happened in the Cambrian explosion induced by the increase of the radius of Earth by factor 2 will be discussed.

The Expanding Earth Hypothesis (EEH) is discussed in various articles [L11, L7, L15].

1. Cosmic expansion according to general relativity (GRT) predicts that astrophysical objects should expand smoothly. This does not happen. In the TGD Universe, the expansion would be basically a quantum phenomenon and take place in rapid jerks and such a jerk would have induced Cambrian Explosion (CE).
2. Expanding Earth Hypothesis (EEH) stated that the radius of Earth increased rather rapidly by a factor of about two in Cambrian Explosion (<https://cutt.ly/x2zaWAe>) (CE) that started about 541 million years ago and lasted about 13-25 million years.

1.1 Vision of the evolution of life on pre-Cambrian Earth

The recent view of pre-Cambrian era has problems. How to solve faint Sun paradox (<https://rb.gy/mfhavz>): was some additional source of energy present and heat the surface of Earth to make liquid water possible? What happened in Great Oxygenation event (GOE) (<https://cutt.ly/K2jAxV9>)? Did deep oceans really exist? Did Snowball Earth (<https://rb.gy/qkoiah>) precede Cambrian Explosion (CE) (<https://cutt.ly/x2zaWAe>)? What happened in CE?

TGD view of EEH leads to a vision of how underground oceans could have served as seats for highly evolved photosynthesizing life, which bursted to the surface and formed the recent oceans.

1. Life would have evolved in underground oceans shielded from meteoritic bombardment and cosmic rays. The radius of Earth increased rapidly by a factor of about 2 during the Cambrian explosion (CE). The multicellular life utilizing photosynthesis bursted to the surface of Earth and formed recent oceans.

There would have been no oceans before the CE. Hydrothermal vents could have existed. The possible lifeforms were very simple bacteria, which photosynthesized using H_2S since there was now water and oxygen.

Earth was like Mars now: Mars has no oceans and no oxygen. There are indications of underground reservoirs of water and signs of simple life forms.

2. Highly developed multicellular animals and photosynthesizing algae bursted to the surface. Note that algae are responsible for the production of most oxygen also in the recent oceans. If hydrothermal vents contained sulphur based life it disappeared because the generation of the basic building blocks of biomolecules was too slow.

Interestingly, the radius of Mars is roughly 1/2 of that for Earth. Could Mars have underground oceans teeming with life? When does the radius increase by factor two?

3. There is however a problem. How is photosynthesis possible underground? It is dark there! The basic proposal is that solar photons with energies in the visible and possibly infrared range arrive as dark photons along monopole flux tubes, which extend above the Earth and carry dark matter. The strength of the magnetic field would be about .2 Gauss and fraction 2/5 of the nominal value of the Earth's total magnetic field involving also a non-monopole part.
4. Also dark photons from the interior of Earth propagating along the flux tubes or associated with them could have served as an energy source. The temperature in the Earth's inner core (with radius about 20 percent of the Earth's radius) corresponds to about 5,500 K, which corresponds to a thermal energy scale of about .55 eV, which corresponds to the nominal value of the metabolic energy quantum.

The energy at the maximum of the energy distribution is roughly 3 times larger than this energy and would be around 1.65 eV. The energy at the maximum wavelength of thermal energy distribution is 5 times higher and about 2.75 eV, which is the upper bound for the energy range 2-2.75 eV of visible photons.

If the temperature of the inner core before CE has not differed appreciably from that now, which could hold true if the inner core was already before CE in the expanded state as also water containing regions, the idea about dark photons from the inner core as a metabolic energy source, which would make possible the evolution of photosynthesis in underground oceans, makes sense.

1.2 A model for the phase transition increasing the radius of Earth by factor 2

The idea about relatively fast growth of the Earth radius by factor 2 (during 13-25 million years) raises the eyebrows of standard physicists. How can such a large change of density make sense? It seems safe to exclude the possibility that the mass of Earth has increased roughly by a factor of 8 (mass should have arrived from dark magnetic flux tube structure to which the core of Earth is associated as a tangle). It must be admitted that the question of how this scaling could have occurred has remained poorly understood.

In this article, a model for how this scaling could have occurred is proposed. The key question that I was not previously aware of, relates to the energetics. Where did the required energy compensating for the decrease of the gravitational binding energy and providing the energy required by the expansion come from? Some new physics seems to be necessary.

In the TGD framework, the quantum phase transitions of the magnetic body (MB) increases the thickness of the flux tubes and reduce their string tension determined by Kähler magnetic energy and volume energy (to which a hierarchy of values of cosmological constant can be assigned). This leads to a liberation of energy and this energy feed could have made possible to induce ordinary phase transitions requiring energy.

The analogy between recent Mars and pre-Cambrian Earth suggests that the Earth had only a single core analogous to the recent inner core which is solid and consists mostly of Fe and Ni. The most natural phase transition would have transformed part of this core to the outer core which is liquid and has a smaller density and has also lower temperature. This could have forced at least the horizontal expansion of the mantle. The thickness of the mantle could have increased in a further transformation of the newly formed outer core to mantle or transfer of material from the outer core to the mantle.

In this and earlier articles [L3, L11, L21] a rather detailed view of the pre-Cambrian biology, geology, and thermal evolution is developed and one can relate it to the standard view. This involves topics like faint Sun paradox, the mechanism of the Great Oxygenation Event, understanding the TGD counterparts of supercontinents Rodinia and Pannotia preceding CE, snowball Earth, and CE that led to a sudden emergence of highly advanced multicellulars.

I have become recently aware of new empirical support for the TGD view of CE.

1. There is empirical support for the view that the oxygenation of oceans did not occur before CE. This conforms with the prediction that oxygenation was due to photosynthesis in underground oceans. TGD provides the new physics needed: dark photons from either Earth's core or Sun could have provided the metabolic energy making photosynthesis and therefore oxygenation possible.
2. Anomalously high recession velocities for the tectonic plates during CE have been observed and could be due to the radial expansion of the Earth lasting about 30 million years which corresponds to the duration of CE. A quantitative estimate for the expansion velocity gives an estimate consistent with the findings. CE would correspond to quantum tunnelling in astrophysical scale and involve "big" state function reductions and a temporary change of the arrow of time. One could even understand the plant fossils with age about 600 million years conflicting with the fact that the CE (CE) occurred about 540 million years ago.
3. The finding that the mantle-core boundary looks like a seafloor having even mountains has a rather convincing explanation in terms of the subduction of tectonic plates, which sink to the mantle. This however inspired the question whether life in underground oceans as porous structures containing water in some exotic form, most naturally the fifth phase of water studied by Pollack playing a key role in the TGD inspired view of biology, could make possible the needed thermal and chemical isolation. Pollack effect could provide this isolation and is certainly needed even if the temperature of the underground ocean is not far from the physiological temperature.

Assuming that the Sun was faint so that the temperature at the surface of Earth was below the freezing point, one ends up with conflict with the isotopic determination of the temperature giving a temperature of oceans slightly higher than the temperature 38 C above which marine invertebrates cannot survive. The temperature about 30 degrees allows life but this requires

a slightly lower amount of O^{18} isotope than prevailing in the recent oceans. The paradox can be solved if the warm water originated from underground oceans and mixed with the non-oxygenated water (or actually ice) at the surface of Earth so that the isotopic fraction was reduced. The optimal situation for life would have been at depths of order kilometer and one can say that life had no other option than developing underground.

2 A possible view of the expansion of Earth?

The most natural TGD inspired guess is that a phase transition at the level of MB, increasing flux tube thickness, induces a phase transition at the level of ordinary matter by providing energy in the case that it requires energy. There are several questions to be answered.

2.1 Did the inner core participate the expansion?

Did the inner core (<https://cutt.ly/P2jS1jB>) participate in the expansion?

1. Could it be that the outer core emerged and this led to a generation of convective currents giving rise to the Maxwellian part of the magnetic field. The temperature at the boundary of the inner core is the same as the solar surface temperature. Note that in the "standard model" the weak interactions within the mantle are assumed to produce energy.

The recent radius of the inner core is $R/5$ and so small that there is no need for it to participate in the expansion: one would $4R/5 \rightarrow 8R/5$ for the outer and the radius would increase by factor $9/5$: not far from 2.

2. Could it be that the outer core with the recent thickness 2400 km emerged in a phase transition transforming the Fe-Li solid of the inner core to Fe-liquid of the outer core so that the inner core could have reduced in size? Was the mantle+crust with recent thickness 2390 km (the recent Earth radius is $2890+2400+1220=6370$ km) scaled in the transition by factor 2 also in the radial direction or only horizontally?

The scaling of the Earth radius by factor 2 gives the condition

$$d(\text{mantle}, i) + d(\text{inner}, i) = (1/2)[d(\text{mantle}, f) + d(\text{outer}, f) + d(\text{inner}, f)] .$$

One has $d(\text{mantle}, f) = 2890$ km, $d(\text{outer}, f) = 2200$ km, $d(\text{inner}, f) = 1220$ km.

- (a) If the mantle thickness was scaled by factor 2 ($d(\text{mantle}, i) = d(\text{mantle}, f)/2$), one has $d(\text{inner}, i) = 1810$ km, which is larger than 1220 km as required. For this option the initial value of the mantle thickness could have been small and the thickness of the inner core correspondingly larger.
- (b) The second option is that the newly formed outer core partially transformed to mantle and increased its thickness so that no radial scaling of the mantle was needed. This option is perhaps the most plausible one.
- (c) For the no-scaling option ($d(\text{mantle}, i) = d(\text{mantle}, f)$) one would have $d(\text{inner}, i) = 365$ km, which is suspiciously small and smaller than 1220 km. Therefore the scaling of the mantle thickness is the more realistic option.

2.2 How the outer core was formed?

What happened in the formation of the outer core?

1. The proposal is that regions of dark water with an increased value of $h_{eff} = h_{gr}$ were generated in the mantle. Inside these regions photosynthesis occurred using the dark photons carrying metabolic energy from the inner core and outer core. The range of energies is the same as in the radiation from Sun [L15].

2. Did the expansion of the volume force a formation of underground oceans in mantle containing dark water, which then bursted to the surface? Was the formation of oceans necessary? Were dark matter blobs enough? Did they condense to form larger dark water volumes, which eventually bursted to the surface?
3. The material in the core of Earth derives basically from chondrites, which contain water and also organic molecules. This suggests that the water of the underground oceans in the mantle derives from the chondrites and that the presence of the basic biomolecules in chondrites was essential for the evolution of life inside Earth.
4. In the case of water, superionic ice [D2] (<https://cutt.ly/uXUIkUQ> and <https://cutt.ly/3XUIWhX>) existing at extreme pressures is a possible candidate for the exotic phase of water. Superionic ice is proposed to appear in the mantles of giant planets such as Uranus and Neptune and in [L11, L7] the possibility that it could occurring the Earth's mantle was considered. The density of superionic ice is slightly less than 4 times the density of ordinary ice.

Could superionic ice in the mantle have transformed to dark water with a volume larger a factor $4^{2/3}$. This would have contributed to the increase of the volume of the mantle. Note that the transition could have led to expansion only in horizontal directions increasing the 2-dimensional volume by a factor 4.

5. One can compare the situation with that in recent Mars. For Mars, the inner and outer core are one and the same thing so that the situation corresponds to that in pre-Cambrian Earth. Also the radii were nearly the same.

On Mars, the temperature at the surface of the core is about 2000 K ($E = .21$ eV). If the temperature for the pre-CE Earth was the same, the temperature of the core came 1.5 times higher and inner core 2.5 times higher in the phase transition. This could be understood if energy was liberated in the thickening of the flux tubes.

2.3 The energetics of the transition

What can one say of the energetics of the phase transition?

1. Expansion requires energy. Where did the energy come from? The fraction of gravitational energy Earth from its mass is of the order of $GM/R \sim 10^{-9}$ and its reduction in the expansion was of the same order of magnitude 10^{-9} . Energy was needed to induce the expansion. Also the reduction of the average density and the increase of temperature required energy. Where did the energy come from? Did it come from the increase of the flux tube thickness reducing string tension?

2. Magnetic and volume energy should have been liberated in the model based on the thickening of cosmic strings or flux tubes. Generation of ordinary matter or a phase transition for an ordinary matter can be imagined.

The liberated energy could have driven the expansion as an explosion. Also the heating of the matter would be an outcome.

3. The increase of h_{gr} by a factor 2 is one option that one can imagine. The reduction of the quantized velocity parameter β_0 by factor 1/2 could have happened $\beta_0 \rightarrow \beta_0/2$? This process should have had as a counterpart ordinary phase transition liberating energy.

One can consider two options for the phase transition at the level of the MB. The first option would correspond to a thickening of cosmic string to monopole flux tube and second option to that of an already thickened cosmic string. A more realistic mechanism involves the expansion of an already thickened flux tube with much smaller liberated energy. Factor of order 10^{-9} is enough.

1. The phase transition for the ordinary matter would have been induced by a phase transition at the level of the MB of the system involving thickening of the flux tubes. The ordinary matter would have been in the form Fe and Ni inside in the inner core.

2. The ordinary phase transition could have been melting of the inner core at its outer surface so that it would have gradually generated the outer core as a liquid layer between outer and inner cores. The process would still continue.

Temperature would have increased in the melting from 2000 K to about 5000 K in the inner core and from 2000 K to 3000 in the outer core. The energy liberated in the thickening of the flux tubes would have provided this energy.

The expansion of the core and the liberated energy would have driven the expansion of the volume above the contracting inner core. This process could have also forced the expansion of the outermost layer with recent thickness about $.444 R_E$. It is also possible that the newly formed outer core transformed partially to mantle so that no radial expansion of the mantle was not needed.

3 TGD view of the pre-Cambrian era

In the sequel a TGD view general view of pre-Cambrian era is proposed followed by a discussion as a series of questions. The basic idea is to take the recent Mars as a guideline in attempts to understand pre-Cambrian Earth.

3.1 A rough overall view of pre-Cambrian era

The recent view of pre-Cambrian era has problems. How to solve faint Sun paradox (<https://rb.gy/mfhavz>): was some additional source of energy present and heat the surface of Earth to make liquid water possible? What happened in Great Oxygenation even (GOE) (<https://cutt.ly/K2jAxV9>)? Did deep oceans really exist? Did Snowball Earth (<https://rb.gy/qkoiah>) precede CE?

3.1.1 Energetics during the pre-Cambrian era

The basic assumption is thermal flow equilibrium in which the energy from the Sun is temporarily stored and leaves the system in time determined by the planet. In the TGD framework, the flow of energy from the interior of Earth forces us to challenge this picture.

1. The total energy flux from the Sun would have been by a factor $1/4$ smaller than now although local flux would have been the same. One can wonder whether this had some implications. If the biosphere is controlled by MB having quantum coherence in the scale of Earth, this might be the case. As a matter of fact, during CE the phase transition generating inner core would have fed energy liberated in the thickening of flux tubes of the MB of Earth to also to the surface of Earth. Inner core and core could have also served as source of metabolic energy for the photosynthesizing life forms in underground oceans [L15].
2. Solar energy input and reradiation of energy by greenhouse gases storing the energy temporarily. Oceans bind 90 percent of the solar energy in the recent Earth. In the TGD framework, oceans would have been absent during the pre-Cambrian era and could not have stored energy in large scales so that the situation would have been similar to that in recent Mars. This would explain why multicellular life was virtually absent during pre-Cambrian period.

Lakes and small oceans could have been present and the energy from the Earth interior could have warmed them. Also volcanic activity could have transferred energy to the Earth's surface as it does also nowadays. Reradiation would have been missing unless greenhouse gases were present locally.

3. Sunlight absorption depends on various factors. The spin of Earth and the tilt of the orbit with respect to the rotation plane affect the absorption in the time scale of $10^4 - 10^5$ years. Milankovitch cycle for the ice ages relates to this dependence.

The duration of the Carbon cycle corresponds to millions of years. Also weathering and volcanic activity affect the absorption. Greenhouse gases are an important factor on recent Earth. Also the distribution of continents affects the absorption (<https://cutt.ly/62jSQU>).

3.1.2 The atmosphere of Earth before CE

What about the surface pressure before CE?

1. Pressure is determined by the concentrations of molecules and atoms in the atmosphere. The abundances of O and N were small before CE: the situation could have been as in Mars today.
2. The composition of the Earth's earliest atmosphere is not known with certainty. Present nitrogen, N_2 , and carbon dioxide, CO_2 , which are also the predominant nitrogen- and carbon-bearing gases produced by volcanism today. These are relatively inert gases.

The leakage of molecules in CE Earth was not as intense as in Mars today since the gravitational field was 10 times stronger than in Mars. Leakage partially explains low surface pressure if the production of gases has roughly the same rate.

3. It is interesting to compare the percentages of various gases in the recent Earth and recent Mars .

In the recent Earth: the fractions of N and O are 78 % and 21 % respectively. The variation of Oxygen fraction is between 10 and 35 per cent during the last 541 million years (<https://cutt.ly/22jSUzR>).

In recent Mars the fractions of CO_2 , N and O are 96 %, 1.9 %, and .15 % respectively. The partial pressure of Oxygen present in the Martian atmosphere is $pO = .95 \times 10^{-2}$ atm which corresponds to the value of pO in the beginning of the second stage of GOE at Earth.

Recent Mars could give some hints of the situation in CE Earth. CO_2 or other greenhouse gases should have been present. Also methane CH_4 was present. These gases could have been produced in the interior of Earth as in Mars and responsible for the warm Earth.

4. Ideal gas approximation gives the estimate $P = nT$. Hydrodynamic equilibrium requires that the pressure gradient is equal to gravitational force density, which determined by the density of the atmosphere and by the strength of the gravitational field, which was 4 times stronger for pre-Cambrian Earth according to TGD. Note that the gravitational energy of protons in the Earth's gravitational field is of order 1 eV and twice the nominal value .5 eV of the metabolic energy quantum [L14].

The transfer of oxygenated fluid from underground oceans would have made oxygenation possible in pre-CE Earth. This would have caused local oxidation of iron without deep oceans.

5. The climax of the GOE caused by accelerating expansion of Earth corresponds CE. Before GOEs there is a slow oxygenation, which in the standard model is assumed to be caused by the photosynthesis of cyanobacteria. What really happened during CE is a mystery as is also CE. In the TGD framework, the underground oceans would have played a key role. Also cyanobacteria could have emerged from the underground oceans.

3.1.3 Maybe deep oceans are not necessary?

According to the Wikipedia article (<https://cutt.ly/K2jAxV9>), the end of the deposition of banded iron formation at 1.85 billion years ago is interpreted as marking the oxygenation of the deep ocean. Banded iron formation largely disappears from the geological record at 1.85 billion years ago, after peaking at about 2.5 billion years ago. Banded iron formations can form only when abundant dissolved ferrous iron is transported into depositional basins, and an oxygenated ocean blocks such transport by oxidizing the iron to form insoluble ferric iron compounds.

But are deep oceans really needed? One can compare the situation with the situation in recent Mars. Mars is red and contains oxidized iron.

1. Rivers and water flow at its surface and there are no deep oceans in the recent Mars. The simplest assumption is that the situation could have been the same always. This applies also to the existence of the Maxwellian part of the magnetic field requiring an inner core.

Although the Martian atmosphere has a low Oxygen content, the iron rich material is oxidized in presence of water and develops rust giving the color red. Also in Mars, band iron formations consisting of non-oxidized iron and dating to times before the beginning of Martian GOE, should exist. The flow of a non-oxygenated water could have transported ferrous iron to the depositional basis.

2. The amount of oxygen in the Martian atmosphere is very low but the oxygenated water percolating from the underground oceans could induce the oxidation of iron. The same mechanism could be at work on the early Earth. The beginning of the oxygenation would correspond to the emergence of oxygen based life forms to the underground oceans.

3.2 Comparison with the recent Mars

The resemblances between Mars <https://cutt.ly/s2jA0FJ> and pre-Cambrian Earth according to TGD inspire the question whether the recent Mars could be like the pre-Cambrian Earth after the initiation of GOE. Could one use these resemblance to understand pre-Cambrian Earth in the TGD Universe.

1. For Mars the radius and distance from the Sun are $R = .53 \times R_E$ and $d = 1.523AU$. Similar tilt of a rotation axis.

The mass of Mars is $M = .107M_E$ so that the surface gravitation is .38 g. For pre-CE Earth surface gravitation is roughly 10 times stronger.

2. Surface temperatures in Mars vary from 20 C to -153 C at poles. The average temperature is -62 C. The temperature drops rapidly with height and with the time of day since the atmosphere is very thin and cannot store heat energy. The ratio of energy fluxes to Mars and Earth is equal $d/d(Mars) = .43$ so that there is no big difference between Mars and Earth.
3. Surface pressure at Mars is $p(Mars) = 0.0628$ atm which is 6.3 % of the pressure $p(Eart) = 1$ atm.

The Martian atmosphere contains 0.174 % Oxygen and 2.8 % Nitrogen. At the Earth, the corresponding abundances are 29 % O and 78 % N. Oxygen partial pressures for Oxygen is $pO(Mars) = 0.01$ atm. There are active sources of gases, such as methane, in the interior of Mars.

4. On Mars there are no oceans and continents are absent. If Earth is like Mars, the oceans could have been absent also on Earth. Instead of them small lakes and rivers could have been present as in the recent Mars. They would have contained oxygenated water from oceans inside the Earth interior, where photosynthesis was producing oxygen. Continents would have been absent so that the existing vision about the history of the continental drift before CE, should be obsolete albeit natural if one assumes that the radius of Earth has always been the recent one.
5. Could Mars be experiencing the analog of GOE? During the second stage of the Great Oxidation Event (GOE) (see the picture <https://cutt.ly/d2jAEd3> in Wikipedia article <https://cutt.ly/K2jAxV9>), the oxygen content of the atmosphere gradually increased during period 2.45-.84 Ga to values of .02 and .04 atm but it is assumed that Oxygen was absorbed by the oceans and seabed rock. During the third stage of GOE, the oxygen content did not change. It is assumed that the oxygen started to gas out from oceans but was absorbed by land surfaces. At the fourth state of GOE, CE occurred and the oxygen content increased rapidly to the recent 21 per cent.

The recent value $pO(Mars) = 0.01$ atm is below .02 atm at the end of the second phase of GOE but would correspond to the value of $pO \simeq .01$ on Earth in the beginning of the second state of GOE. Could it be that GOE on Mars has begun? The naive guess is that there is still 2.45 Gy to the recent situation on Earth.

6. The interior of Mars consists of a crust, a mantle consisting of silicates and a solid metal core consisting of Fe and Ni as the solid inner core of Earth. Crust has a typical thickness of 50 km, which is not far from that on Earth.

There is no liquid outer core, which in the case of Earth is responsible for the convective ionic currents creating the Maxwellian part of the magnetic field (note that the monopole part needs no source). In Mars the magnetic field is indeed absent in large scales. Martian Auroras have been observed however. In the TGD framework, this suggests that only the dark part of the Martian magnetic field has a considerable strength whereas the Maxwellian part is very small.

This suggests that in the case of Earth the emergence of the Maxwellian part of the magnetic field made possible the shielding of life against cosmic rays and that life did not have any other option than to evolve in the womb of Gaia.

7. For the recent Earth, the volume fraction of the outer core of Earth 16.5 %. The outer core carries the convective currents giving rise to the Maxwellian part of the magnetic field of Earth. The inner core however feeds heat energy to the outer core and has a considerable effect on its dynamics. This conforms with the idea that the inner core serves as a source of heat energy.
8. The temperature $T = 1900 - 2000$ K at the boundary of Martian core corresponds to a thermal energy $\simeq .2$ eV, which is below the metabolic energy quantum .5 eV.

On the recent Earth, one has $T = 3000$ K (.3 eV) at the upper boundary of the outer core and $T = 5000$ K (.5 eV) at the upper boundary of the inner core: the latter is the temperature at the surface of the Sun. These observations lead to the proposal [L15] that the energy feed from the inner and outer cores, realized as dark photons, could have served as a metabolic energy source for the evolving life in the underground oceans. Also the solar photons might have transformed into dark photons propagating along the flux tubes of the monopole part of the Earth's magnetic field to the interior of Earth. One can even ask whether this mechanism might be still at work at polar regions of Earth covered by ice or at the bottom of oceans.

There is now a considerable evidence that ancient Mars has had lakes or even oceans (see this). Could underground life in Mars bursted much earlier to the surface of Mars?

1. The average density of Mars is near to that of recent Earth (mass is .1 Earth masses and radius roughly 1/2 of the Earth radius R_E). This leads to the question of whether Mars has already experienced a similar transition increasing its radius by 1/2 and density by factor 8? This would have brought the possible underground water to the surface. Later the water would have been lost. Mars would not have been as lucky as Earth.
2. The objection is that Mars has no plate tectonics, which should have emerged in the rapid expansion. The alternative option that I have discussed earlier is that Mars is still waiting for the expansion to take place. Intriguingly, Mars has the same radius as Earth before the Cambrian explosion. The ancient presence of oceans/large lakes does not support this view. One might however think that the water from underground oceans leaks to the surface and forms lakes and even shallow oceans.

3.3 Some questions related to the pre-Cambrian period

Pre-Cambrian period Cambrian explosion (CE) are not well-understood and there are several paradoxical aspects involved.

3.3.1 Why so few multicellular fossils from pre-Cambrian period?

There is very little evidence for multicellulars from pre-Cambrian period whereas fossils of unicellulars (cyanobacteria) exist. The oldest multicellular fossils are 1.2 billion years old. Multicellular fossils with an age of 600 million years and thus preceding CE, which began 541 million years ago have been also found (<https://cutt.ly/d2jAgzb>).

The multicellular organisms preceding CE are much simpler than plants and animals that emerged in CE. The standard explanation for the nearly complete absence of multicellular fossils is that they have disappeared almost completely because they had no hard parts. What is however very strange is that suddenly the fossils of rather complex multicellulars emerge suddenly in CE [112].

3.3.2 Questions related to energetics

The first questions relate to energetics.

1. Faint Sun paradox means that solar radiation was only 70 per cent from the recent temperature. The Earth surface was however warm since liquid water existed. Could Earth's interior have served as a source of heat energy? This seems to require new physics.
2. In the TGD framework one can ask whether a quantum phase transition proceeding at the level of MB (MB) of Earth could have induced ordinary phase transition at the level of ordinary matter, say solid-liquid phase transition of the Fe-Ni at its boundary giving rise to the outer core. This would have made convective charged currents generating the Maxwellian part of the magnetic field of Earth providing a shield against cosmic rays. This kind of induction of ordinary phase transitions by quantum phase transitions could be crucial for the evolution of life [L14, L13].
3. Could the energy liberated in the phase transition of the MB increasing the thickness of the flux tubes have driven this solid-liquid phase transition and have also feeded heat energy to the surface of Earth increasing the temperature. Also the increase of the gravitational potential energy in the increase of the radius by factor 2 requires a lot of energy.

Note that nuclear physics cannot be involved with the energy production. Basic stable isotopes of Fe and Ni have atomic numbers 56 and 58. Ni has also $A = 60, 61, 62, 64$ as stable isotopes. Fe has stable isotopes with $A = 54, 57, 58$ besides $A = 56$. Ordinary nuclear processes cannot transform Ni nuclei Fe nuclei and new physics is needed.

3.3.3 Did large oceans really exist?

The existence of oceans covering most of the surface of Earth is assumed in the standard view of pre-Cambrian period. But did Earth have oceans at its surface before CE?

1. Could it be that the situation before CE was like in the recent Mars. This plus the assumption that the radius of Earth was only $1/2$ of the recent radius would totally change the views about pre-CE Earth both biologically, geologically, and thermodynamically.
2. The simplest working hypothesis is that plate tectonics was not present before CE. Standard view (<https://cutt.ly/I2zf8v8>) however is that roughly 750 million years ago, the earliest-known supercontinent Rodinia, began to break apart.

Did the breaking apart of Rodinia initiate the expansion of Earth, which gradually accelerated? If so, the area covered by the analogues of oceans between continents was much smaller than on the recent Earth. The narrow fissures would have contained water which had bursted from the underground oceans and brought multicellular life forms.

The analogues of continents would have later recombined to form Pannotia 600–540 Ma. Did the expansion involve a pulsation leading to the formation of Pannotia, which then split again to continents separated by narrow analogs of oceans. The multicellular lifeforms giving rise to 600 Ma old fossils could have emerged from underground oceans in the breaking apart of Pannotia.

3.3.4 Snowball Earth hypothesis from TGD point of view

The assumption is that snowball Earth dating back to 650 Ma, involving several large scale glaciations, preceded CE. The Earth would have suffered a global glaciation but evidence exists only for local glaciations.

What could be the TGD counterpart of the snowball Earth? Milankovitch cycles (<https://cutt.ly/12zkeoF>) with a period of order 100,000 years were present and could have caused local glaciations explaining the observed glaciations. The two large scale glaciations associated with snowball Earth could relate to the expansion of Earth. Did the glaciation take place for the oceans defined by the narrow fissures between the analogues of continents formed in the splitting of Rodinia?

3.3.5 The Great Oxygenation Event

The Great Oxygenation Event oxidized the atmosphere and made oxygen based life possible at the surface of Earth.

1. The oxygen content of the atmosphere and assumed oceans was very low before CE and could not support oxygen based life. During the Great Oxygenation Event (GOE) (see the Wikipedia article (<https://cutt.ly/K2jAxV9>), which started for about 2.45 Gy ago, the oxygenation of atmosphere proceeded very slowly (see the Wikipedia picture <https://cutt.ly/d2jAEd3>).

In CE (CE) the oxygen content of the atmosphere increased very rapidly to the recent level of about 21 per cent. The oxidation is made manifest by the observed layers of oxidized iron. The absence of oxygen for more than 2.45 Gy ago is demonstrated by layers of iron based compounds, which are not oxidized.

2. Usually it is assumed that the oceans were present during the pre-Cambrian period and that the sea water was oxidized gradually. On Mars the situation is however different. There are local sources of oxygenated water such as lakes and rivers but no oceans. There is also evidence for underground life on Mars.

Could the situation on Earth have been like in Mars during the pre-Cambrian Earth? Could the underground oceans, serving as a kind of womb shielding the evolving life from meteoric bombardment and cosmic rays, have made possible the evolution of the photosynthesizing life and multicellulars producing oxygen, which bursted to the surface of Earth in CE?

3. Suppose that the water from the underground oceans, containing highly developed multicellulars, bursted to the surface during the GOE. What could be the time scale of the process? Was GOE implied by this process and did it accelerate during CE? Did this increasing rate of bursts lead to oceans of increasing size so that the scale of glaciations was dramatically increased during the last two glaciations before CE.

3.3.6 Was CE a quantum phase transition in Earth scale?

The TGD view of CE explains the sudden appearance of multicellulars and also implies that the gravitation would have been 4 times stronger before CE so that the possible lifeforms surviving at the surface of Earth are expected to have a flat shape. The reduction of surface gravitation after CE in turn could in turn explain the emergence of giant plants and animals.

In the TGD framework the evolution of life would have involved quantum criticality and quantum phase transitions closely related to ordinary criticality and phase transitions [L13]. CE would be naturally a quantum phase transition in the scale of Earth involving in an essential manner gravitational Planck constant \hbar_{gr} making possible dramatic reduction of gravitational binding energy in the scaling up of Earth radius by factor 2.

In the TGD framework the standard ontology of quantum theory is replaced with zero energy ontology (ZEO) solving the basic paradox of the quantum measurement theory [L5, L9, L8, L12].

Interesting questions relate to the proposal that a pair of "big" state function reductions (BSFRs) in astrophysical scales (counterpart for quantum tunnelling) changing the arrow of time was in question. Could one interpret the time reversed classical time evolution after the first BSFR, that is GOE, as a classical correlate for the pair of BSFR making the process effectively to look like deterministic and smooth classical process [L4] [L4]?

3.3.7 Oxygenation of oceans did not precede Cambrian Explosion as TGD strongly suggests and empirical study concludes

There was an interesting article in Futurism about the findings challenging the basic assumption related to the origin of life (<https://futurism.com/the-byte/textbooks-wrong-origin-life>). The research article "Widespread seafloor anoxia during generation of the Ediacaran Shuram carbon isotope excursion" [I4] can be found at [urlhttps://onlinelibrary.wiley.com/doi/10.1111/gbi.12557](https://onlinelibrary.wiley.com/doi/10.1111/gbi.12557).

On the basis of empirical evidence it is claimed that the view of the gradual oxygenation of oceans is wrong. The abstract of the article explains the findings.

Reconstructing the oxygenation history of Earth's oceans during the Ediacaran period (635 to 539 million years ago) has been challenging, and this has led to a polarizing debate about the environmental conditions that played host to the rise of animals. One focal point of this debate is the largest negative inorganic C-isotope excursion recognized in the geologic record, the Shuram excursion, and whether this relic tracks the global-scale oxygenation of Earth's deep oceans.

To help inform this debate, we conducted a detailed geochemical investigation of two siliciclastic-dominated successions from Oman deposited through the Shuram Formation. Iron speciation data from both successions indicate formation beneath an intermittently anoxic local water column. Authigenic thallium (Tl) isotopic compositions leached from both successions are indistinguishable from bulk upper continental crust ($\epsilon^{205}\text{Tl}_A \simeq -2$) and, by analogy with modern equivalents, likely representative of the ancient seawater $\epsilon^{205}\text{Tl}$ value. A crustal seawater $\epsilon^{205}\text{Tl}$ value requires limited manganese (Mn) oxide burial on the ancient seafloor, and by extension widely distributed anoxic sediment porewaters.

This inference is supported by muted redox-sensitive element enrichments (V, Mo, and U) and consistent with some combination of widespread (a) bottom water anoxia and (b) high sedimentary organic matter loading. Contrary to a classical hypothesis, our interpretations place the Shuram excursion, and any coeval animal evolutionary events, in a predominantly anoxic global ocean.

The absence of oxygenation before the Explosion is also the TGD based prediction. The TGD based model predicts that the mysterious Cambrian Explosion in which advanced multicellulars suddenly emerged involved an increase of Earth radius by factor 2 in a geologically short time scale [L3] [L11].

This view conforms with TGD inspired cosmology in which a smooth cosmological expansion is replaced with a series of rapid step-wise expansions, which are essentially quantum phase transitions involving astrophysical quantum coherence scales.

The Earth would have been like Mars now (not much water at surface, even the radius would have been very near to Mars radius) and life would have evolved in underground oceans (being shielded from cosmic rays and meteors), in the womb of Mother Gaia. The evidence for the underground life on Mars is accumulating. The water in underground oceans would have oxygenated by photosynthesizing life and the underground water would have bursted to the surface and give rise to the oxygenated oceans covering most of the Earth's surface since then.

The basic objection is that photosynthesis was not possible. The core of the Earth however produces radiation in the same wavelength range as the Sun and in the TGD framework this could allow the development of photosynthesis [L15]. Now Quanta Magazine article ([rebrand.ly/jbvkwt](https://www.quantamagazine.org/rebrand.ly/jbvkwt)) told that in a research [I7] published last month in Nature Communications, researchers reported that in groundwater reservoirs 200 meters below the fossil fuel fields of Alberta, Canada, they discovered abundant microbes that produce unexpectedly large amounts of oxygen even in the absence of light. Photosynthesis is the standard way to produce oxygen. But how could photosynthesis work underground? This looks like a complete mystery in the standard physics framework.

Whether the interpretation of the article that life evolved in anoxic global ocean is consistent with the TGD view that this oceans was essentially absent, is not of course clear.

The basic objection is that photosynthesis was not possible. The core of the Earth however produces radiation in the same wavelength range as the Sun and in the TGD framework this could allow the development of photosynthesis [L15].

Whether the interpretation of the article that life evolved in anoxic global ocean is consistent with the TGD view that this oceans was essentially absent, is not of course clear.

What objections can one invent against the TGD view? The first objection against the TGD

proposal is that fossils of complex multicellular life forms have been found with age of 600 million years to be compared with the fossils dating back to 550 years ago when Cambrian explosion could have started. The story of their discovery told in the popular article "How 2 Teens Accidentally Solved Charles Darwin's Most Vexing Problem" (rebrand.ly/uk0ntuk) is fascinating.

In 1956, a teenage girl by the name of Tina Negus found a strange looking fossil in Charnwood Forest in Leicestershire, England. The plant fossil should not have been there since the rock was 600 million years old and the Cambrian explosion started roughly 60 million years later. Tina Negus showed a pencil rubbing to his geography teacher but he didn't believe her.

A year later, in 1957, three teenage boys were playing near the same rock face when they, too, noticed the same fossil. One of these teens, fifteen-year-old Roger Mason, found the second fossil. Roger Mason contacted Trevor Ford, a local geologist, who wrote about the finding to the Journal of the Yorkshire Geological Society. The new plant was named *Charnia Masoni*. *Charnia* is a genus of frond-like life forms belonging to the Ediacaran biota. *Charnia* came from Charnwood Forest in Leicestershire, England where Tina Engus found it. *Masoni* is after Roger Mason.

Does the TGD based view survive these findings? Note first that the *Charnia Masoni* does not conform with the conclusion that the oceans that possibly existed at that time did not contain oxygen. Furthermore, *Charnia Masoni* does not change the basic facts: complex multicellular life forms emerged as if from nowhere. The time span from *Charnia Masoni* fossils about 600 billion years to the beginning of Cambrian Explosion about 538.8 million years ago is about 10 per cent.

Can one explain the finding in the TGD view of the Cambrian explosion?

1. If the surface of Earth did not contain much water before the Cambrian Explosion, one can imagine that water leaked from the underground oceans locally at some places but not everywhere. Instead of oceans, there were oxygenated lakes, where multicellular life forms survived.
2. Another possibility is that the expansion of Earth involves periodic oscillation typical of resonances so that bursts of oxygenated water containing the highly evolved life forms emerged to the surface. I have considered this kind of explanation for the periodic oscillations found to be associated with the Cambrian explosion.
3. One can also imagine that the rock containing the fossil has emerged from the older of a sediment rock at the "bottom" underground water reservoir when a crack was formed in the geologically fast expansion.

There is also indirect evidence for the presence of life about 3.85 billion years ago (rebrand.ly/ebz2q92). The interaction of rocks with life forms lowers the C-13 ratio of rocks to a characteristic value serving as a signature for life. Any sedimentary rocks that formed before life appeared on Earth would have the high C-13 ratio of a volcanic origin. The discovery of rocks with the same C-13 ratio that serve as a signature of life support the view that very primitive microbial life existed already at that time. One can consider the possibility that very primitive life forms existed at the surface of Earth. Second possibility is that they leaked from underground oceans. Also in Mars there is evidence for the presence of water and it could have indeed leaked from underground water reservoirs.

3.3.8 Strange correlation between the strength of the Earth's magnetic field and the atmospheric oxygen content

Sabine Hossenfelder (see this) talked about the recently discovered strong correlation between the strength of the Earth's magnetic field and O_2 content of the atmosphere [\(1001\[?\]\(seethis\)\)](https://www.youtube.com/watch?v=1001[?](seethis)).

This is an extremely interesting finding from the TGD point of view. It is known that the changes in the magnetic field strength and its orientation have had strong effects on biology and consciousness. The local weakening of the magnetic field strength and Schumann resonance have been assigned also with disorders of societies by Callahan [I10, I1] (see this and this). TGD suggests a model for the reversal of the magnetic field [?, L23].

As the figure 1 of the article [?] about the correlation between magnetic field strength and O_2 percent of the atmosphere shows, both the oxygen percent in atmosphere and the strength of the magnetic field characterized by virtual geometric axial dipole moment started to grow about 500

My ago and is maximum about 300 My ago. TGD view of Cambrian Explosion suggests that the strengthening of the magnetic field, perhaps by the emergence of monopole flux tubes, implied a rapid evolution of the oxygen based life forms in the oceans created by the rapid expansion and caused oxygenation of the atmosphere.

3.3.9 Anomalously high rates of tectonic plate motion and the TGD view of Cambrian explosion

The popular Arstechnica article (see this) tells that the motion between plates was surprisingly fast. The rate of the tectonic motion as relative rate for the distance increase between plates was surprisingly fast: even about 4 times the recent one.

The TGD view of expanding Earth relies on the prediction of cosmic expansion as a sequence of fast periods of expansion for astrophysical objects.

1. The model predicts that the tectonic plates were created in rather fast radial expansion of Earth: radius increased by a factor 2. Cracks giving rise plates were formed because rock is not flexible material.
2. The model explains the Cambrian explosion: advanced photosynthesising multicellulars emerged from underground oceans as the oxygenated water bursted to the surface. The TGD view of dark matter allows to circumvent the obvious objections against the model and conforms with the recent surprising findings [L21] (see this) and this).
3. The fast radial expansion caused a fast increase of the distances between plates. The velocity v of this recession would have been $v = dR/dt \times \Delta\Phi$, where $\Delta\Phi$ is the angular distance between the plates and dR/dt is the radial expansion rate.

The duration of Cambrian Explosion was roughly $\Delta T = 30$ million years. Using Earth radius $R = 6,371$ km, one obtains the estimate $dR/dt = R/\Delta T \simeq 20$ cm/year. v is obtained from this by multiplying with $\Delta\Phi < \pi$. The largest rate mentioned in the popular article is $v = 64$ cm/year. The order of magnitude is correct and the rate would have been higher than the average during the fastest periods.

The estimate for v must involve a large enough angle $\Delta\Phi$ and a long enough time period so that $\Delta\Phi$ is expected to be a considerable fraction of π . For $\Delta\Phi$ slightly below π , the estimate is exact but this is probably an accident.

4. Note that the predicted contribution to v is always positive and could provide a test for the TGD view.

A fascinating, and admittedly frightening, question, which just now occurred to me, is whether the Cambrian explosion was gravitational expansion analogous to cosmic expansion in which the metric (!) distances between points doubled! This would have required the scaling of the induced metric by a factor about 4. Could this make sense or does it kill the basic idea?

1. In zero energy ontology (ZEO), light-cone proper time a serves in the role of the cosmic scale factor of either half-cone of the causal diamond (CD) having interpretation as empty cosmology. "Big" state function reductions (BSFRs), serving as TGD counterparts of the ordinary SFRs, change the arrow of time and a pair of BSFRs would be behind quantum tunnelling in the TGD Universe.

In the TGD framework quantum coherence and BSFRs are possible even in astrophysical scales. Could the increase of the radius of Earth be quantum tunnelling be realized as a pair of BSFRs.

2. Can one imagine a local "mini" Big Bang for the CD inside which Earth's space-time surface belongs and a scaling of light-cone proper time a by a factor 2 in astrophysical quantum quantum tunnelling? The value of the light-cone proper time a , characterizing the cosmotemporal position of Earth in a double BSFR, would have increased by factor 2. The spatial scaling by a factor 2 conforms with the p-adic length scale hypothesis stating that p-adic length scales coming as powers of 2 are of special importance.

3. One can try to form a more quantitative view of the situation. Note that the size scale of the initial CD before explosion would be $T/2$, where T is the distance between the tips of the CD would be about 30 million years. The Cambrian explosion occurred about $T_i = 540$ million years ago. If T_i corresponds to the cm of CD, the future tip of the initial CD would be at 570 million years. CD size would be scaled by factor 2 and the end of the cm of CD would correspond to $T_f = 570$ million years.

The quantum average space-time surface would be replaced by a new one in double BSFR and would be modified already 60 million years before T_f . This time would correspond to 630 million years. As explained, some multicellular plant fossils have been found with an age of about 600 million years. Could this replacement of geometric past explain them?

3.3.10 Does the existence of an underground ocean floor at the mantle-core boundary relate to underground life?

The popular article published in Futurism discussed an unexpected observation by the group led by Samantha Hansen published in Science [D1]. The mantle-core boundary in the Earth's interior contains a layer that looks like the crust of Earth in the sense that the seismic perturbations propagating through it have an ultralow velocity. There are mountains many times higher than Himalaya! How is this possible? Is this possible in standard physics?

The answer of the article to this question is based on the idea that subduction for continental plates implies that part of them sinks down because they are denser than the surrounding material and gradually gather to form a second sea floor at the mantle-core boundary. To me this idea looks rather plausible but need not be correct.

My first reaction was the question whether this second sea floor could be a genuine seafloor, the seafloor of an underground ocean! Could new physics predicted by TGD make this possible?

1. The basic prediction of the Expanding Earth hypothesis [L21] explaining Cambrian Explosion is that life evolved in underground oceans and bursted to the surface as the radius of Earth increased by factor 2 in a rapid expansion lasting about 30 million years (cosmic expansion would occurred as rapid jerks for astrophysical objects). During the last weeks several strange findings removing the most obvious objections against this vision have emerged.
2. Could these mountains at the core-mantle boundary correspond to mountains of underground ocean floor?

Could the underground oceans have existed and carried life? Could they reside even in the extremely hostile environment at the mantle-core boundary?

1. Underground oceans near the mantle-core boundary boundary could be imagined as a porous structure having water inside pores. Such structures are very common and if Earth's crust is formed from meteorites the water would be present from the beginning. Even biological matter is analogous to porous structure. When stone is heated it becomes a porous structure. Maybe the enormous heat flux from the core could cause porosity. In accordance with the standard vision of self-organization, this could be understood as complexity developing induced by a constant heat flux. Self-organization takes place at boundaries.
2. It is known that huge reservoirs of water exist underground. The boundary between upper and lower mantle at a depth of about 500 km contains a porous structure carrying water (see this). If the size of the pores is large enough, considerably above cell size, advanced multicellulars could evolve in the underground oceans.

It is easy to invent lethal objections in the standard physics framework.

1. The temperature and pressure increase as one goes towards the core. The temperature of pores should be around 40 C for life to survive. Also the pressure should be normal.
2. Consider the crust first. The temperature reaches the values in the range 100-600 C at the crust-mantle boundary. The temperature increase is about 30 C per kilometer in the upper part of the crust and would be about 30 C at the depth of 1 km if it is 0 C at the surface. The underground water reservoirs should not be at depths much larger than 1 km if the standard physics applies and the largest depths would be possible near the poles.

3. In the underground ocean at the boundary of the upper and lower mantle at a depth of about 500 km, the temperature and pressure are quite too high. Temperature of the surrounding solid material varies from 500 K at the lower boundary of the crust to 1200 K at the boundary of upper and lower mantle. Densities would be several times higher than the normal density of water.
4. Temperature at the mantle-core boundary is about 3000 -4500 K and pressure 1.3 trillion times the atmospheric pressure. The density of mantle is by factor about 5-6 higher than the density of crust so that the pressure is really huge since water and solid matter are almost incompressible. Water in ordinary form cannot exist in this kind of environment if standard physics applies.

Could underground oceans allow some exotic phase of water at physiological temperature around 40 C and normal pressure? This is not possible for the water of standard physics. But the water in living matter is not normal!

1. The phase of water discovered by Pollack [?, ?] called fifth phase of water by Pollack himself (also the term "ordered water" is used). Pollack proposed it to be fundamental for life. Gel phases would represent a basic example of this water.

This phase of water plays a key role in the TGD based model of living matter. The model identifies dark matter based as phases of ordinary matter with non-standard values of Planck constant. The gravitational Planck constant indeed has huge values.

2. The underground life faces the same problem as the biological cell at the surface of Earth: how to isolate itself from the environment. The high temperatures and pressures make the problem orders of magnitudes more challenging. The fifth phase of water surrounding the system could provide the solution in the case of cell membrane and DNA double strand: develop a layer consisting of the fifth phase of water which thermally shields the volume of the ordinary water from the environment at a different temperature.
3. The negatively charged exclusion zones (EZs) generated in the Pollack effect as protons go to magnetic flux tubes as dark protons realizing the dark genetic code, are also able to get rid of various impurities in their interiors. This violates second law in its standard form and suggests that the arrow of time at the magnetic body of EZ controlling its dynamics has a non-standard arrow of time and induces an effective change of the arrow of time for ordinary biomater for long periods of time. This would isolate the system also chemically from the environment.
4. As a matter of fact, it has been discovered that ordinary water in air develops a thin molecular layer at its surface. This layer is neither water or ice and the identification as the fifth phase of water would be suggestive [L16]. This layer could also work at nanoscales and reduce the freezing temperature of the lattice water in materials like concrete to about -70 C. The mechanism could be essentially thermal isolation.

Could thermal isolation work also in high temperature environments, where underground life had to survive?

1. Could the darkness of the ordered water make possible a situation in which the interactions of the water inside pores with the hot high pressure environment are very weak and heat and matter are not transferred between the solid environment and water. Thermal equilibrium would be established very slowly and the temperature could and pressure could be much lower than otherwise for very long periods.
2. Magnetic bodies would carry the dark matter relevant for the biocontrol and would be shielded from the hot environment. They would be gradually heated and this would lead to biological death as it does in ordinary biology according to TGD. Zero energy ontology would however come in rescue and the change of arrow of time would reverse heating to cooling!

3. The unpaired and their chemically non-inert valence electrons of biologically important ions should be dark and reside at the flux tubes associated with very long dark valence bonds. This would generate long range quantum coherence. This would explain why living matter contains these ions although thermal ionization is not possible at physiological temperatures. Also the protons of hydrogen bonds would be dark. Only the chemically inert full electron shells would remain and the system would remain and since be effectively thermally isolated from the hot environment.

As a matter of fact, electrolytes involve ions and the mechanism of ionization is not actually understood and TGD suggest a mechanism of ionization based on the generation of dark valence electrons and dark protons [L17, L10].

While preparing this text, I learned that the standard view of Cambrian explosion has a problem with the Cambrian ocean temperature.

1. If the oceans existed (not clear in the TGD framework before the Cambrian explosion!), their temperatures should have been around 60 C. Marine invertebrates do not however survive above 38 C.
2. Isotopic estimates for Cambrian phosphatic brachiopods [I8] (see) assuming no post-Cambrian O^{18} isotopic depletion relative to the recent concentration suggests that the temperatures of Cambrian oceans were in the range 35-41 C. This range is above the recent range 27-35 C. Assuming a O^{18} depletion of -3 promille of the early Cambrian sea water relative today, one can get Cambrian temperatures around 30 degrees.
3. The problem now is that the underground oceans should have been very near to the surface (unless one assumes the TGD inspired thermal insulation hypothesis). Faint Sun paradox comes to rescue here. The luminosity of the Sun should have been 30 per cent lower than now so that the temperature at the surface of Earth should have been below the freezing point. This has motivated the snowball Earth hypothesis. The paradox disappears if life evolved in the underground oceans in a warmer environment. The temperature deduced from the O^{18} fraction of fossils should be that for the underground ocean. Furthermore, the faint Sun makes it possible for the underground oceans to be located below 1 km depth without assuming the thermal insulation hypothesis.
4. What could have caused the O^{18} depletion of the Cambrian phosphatic brachiopods? A possible explanation is that there were non-oxygenated water reservoirs (presumably in a form of ice) at the surface of Earth and the oxygenated underground water was mixed with this water. Also the surface of Mars, to which the surface of Earth before the Cambrian explosion is analogous, contains some water.

To conclude, I am not suggesting that life developed at the mantle-core boundary: this might be quite too science-fictional an idea. Pole regions of the crust are the most conservative candidate for the seat of underground oceans. It is quite enough for the purposes of the Expanding Earth model that it developed in underground water reservoirs at depths of a few kilometers. Also in this case the thermal and chemical isolation from the environment could have played a key role. An interesting question is whether the critical temperature range 30-40 C of life could fix the depth for the underground oceans in which life most probably evolved.

3.3.11 What are the mysterious structures observed in the lower mantle?

I learned of very interesting results related to geology. The Dailymail popular article (see this) tells about massive structures in the Earth's deep mantle below Pacific Ocean near the mantle-core boundary. The article "Full-waveform inversion reveals diverse origins of lower mantle positive wave speed anomalies" of Schouten et al published in Scientific reports [?] (see this) describes the findings.

There are cold regions deep within the Earth where seismic waves behave in unexpected ways. These regions, located 900 to 1,200 kilometers beneath the Pacific Ocean, defy expectations based on conventional plate tectonics theories. These kinds of structures can result from the subduction of

continental plates leading to the sinking of a plate to the mantle. There are however no subduction records in the Ocean regions so that the mechanism must be different.

It seems that the recent view of the dynamics of the Earth's mantles is in a need of a profound updating. It has been proposed that the structures could be the remnants of ancient, silica-rich materials from the early days of the Earth when the mantle was formed billions of years ago. Alternatively, they may be areas where iron-rich rocks have accumulated over time due to the constant movement of the mantle. However, researchers are still unsure about the exact composition of these deep Earth structures.

Here is the abstract of the article of Schouten et al.

it Determining Earth's structure is paramount to unravel its interior dynamics. Seismic tomography reveals positive wave speed anomalies throughout the mantle that spatially correlate with the expected locations of subducted slabs. This correlation has been widely applied in plate reconstructions and geodynamic modelling. However, global travel-time tomography typically incorporates only a limited number of easily identifiable body wave phases and is therefore strongly dependent on the source-receiver geometry.

Here, we show how global full-waveform inversion is less sensitive to source-receiver geometry and reveals numerous previously undetected positive wave speed anomalies in the lower mantle. Many of these previously undetected anomalies are situated below major oceans and continental interiors, with no geologic record of subduction, such as beneath the western Pacific Ocean. Moreover, we find no statistically significant correlation positive anomalies as imaged using full-waveform inversion and past subduction. These findings suggest more diverse origins for these anomalies in Earth's lower mantle, unlocking full-waveform inversion as an indispensable tool for mantle exploration.

Here some terminology is perhaps in order. Seismic waves are acoustic waves and their propagation in the mantle is studied. Positive speed anomaly means that sound speed is higher than expected. The lowering of temperature or increase of density such as presence of iron can cause this kind of anomalies. The Pacific ocean and the interior regions of plates do not have any subduction history so that the slabs cannot be pieces of continental plates, which have sunk to the mantle.

I have not earlier considered what happened in the lower mantle in the sudden expansion of Earth increasing its radius by factor 2 and giving rise to the Cambrian Explosion. Did these kinds of cracks occur also in the mantle-core boundary and lead to the formation of the recently observed structures also below regions where there is no geologic record for subduction? Could at least some regions which are believed to be caused by the sinking of parts of continental plates have such structure?

Could the Cambrian explosion be a mini Big Bang that happened in the lower mantle and forced the motion of the upper layers leading to the increase of the radius of Earth? The longstanding problem has been the identification of the energy needed to overcome the gravitational force. The order of magnitude of the gravitational binding energy per nucleon is about 1 eV at the surface of the Earth and decreases like $M(R)/M_E)/R \propto R^2$ below it. How did the matter above the monopole flux tube layers get this energy?

1. Since the monopole flux tubes are the key actors, a natural first guess is that there was a layer of dark protons at monopole flux tubes in the lower mantle, say above the core, and that the gravitational energy is compensated by the cyclotron energy of dark proton with gravitational Planck constant $\hbar_{gr}(M(below))$ at monopole flux tube carrying a magnetic field of order of magnitude of endogenous magnetic field. The value of B_{end} need not be the same as its value $B_{end} = .2$ Tesla at the surface of the Earth.
2. If the monopole flux behaves like $1/R^3$, as the dipole character of the Earth's magnetic field suggests, and the mass appearing in the gravitational Planck constant is the mass $M(R) = (R/R_E)^3 M_E$ below the monopole flux tube layer is used, the cyclotron energy is the same as at the surface of the Earth. In the explosion, the value of \hbar_{gr} would be reduced dramatically, perhaps to \hbar and the cyclotron energy would be liberated.

In the interior of the Earth, the gravitational potential energy for mass m is of form $E_{gr} = GM_E m V_{gr}(R)$, $V_{gr}(R) = R^2/2R_E^3 - (3/2)/R_E$ and approaches in the center of the Earth the value $-(3/2)GM_E m/R_E$ and at the surface of the Earth to the value $-GM_E m/R_E$.

3. All nuclei must receive the cyclotron energy compensating the gravitational binding energy and a larger fraction should therefore be dark before the explosion. The gravitational Planck constant $\hbar_{gr} = GMm/\beta_0$ of a nucleus is proportional to its mass number so that the cyclotron energy $\propto ZeB/m$ does not depend on the mass number A of the ion of mass $m \simeq Am_p$. For $1/\beta_0 = 1$, the extreme option is that the entire Earth's interior contains gravitationally dark nuclei meaning that there is a large negatively charged exclusion zone created in the Pollack effect, perhaps giving rise to the electric body assignable to the Earth. Can this be consistent with what is known about the Earth's history? For $1/\beta_0 = 2^{-11}$ assignable with the magnetic body of the Sun-planet system, the value of cyclotron energy would be about 10 keV, which happens to be the energy scale of "cold fusion" identified as dark fusion in the TGD framework [L29]. Could the formation of dark nuclei with nucleon radius of order electron Compton length and with a dark nuclear binding energy of order 10 keV involve the formation of the monopole flux tubes with this dark cyclotron energy?

3.4 A possible mechanism of radiative energy transfer from the Earth's core to underground oceans near the surface of Earth

The recent observations [D1] strongly suggest that there is an ancient seabed on top of the Earth's core, and there are also mountains with a height of about 10 km. The proposed model, in which convection moves the sea floor to the region above the mantle, is probably correct.

This finding combined with the discovery of the so-called superionic ice [D2], which could exist above the Earth's core, allows to develop a proposal for a mechanism of metabolic energy transfer from the Earth's core to the underground oceans near the surface of Earth. This would make possible the development of photosynthesizing life forms in underground oceans. The generalization of the Pollack effect [I9, L1, I13, I11] would play a key role in the mechanism.

3.4.1 Ultralow velocity zones

The following abstract of the article (see this) published by a group led by Dr. Samantha Hansen [D1] gives an overall view of what has been observed.

Ultralow velocity zones (ULVZs) are the most anomalous structures within the Earth's interior; however, given the wide range of associated characteristics (thickness and composition) reported by previous studies, the origins of ULVZs have been debated for decades. Using a recently developed seismic analysis approach, we find widespread, variable ULVZs along the core-mantle boundary (CMB) beneath a largely unsampled portion of the Southern Hemisphere. Our study region is not beneath current or recent subduction zones, but our mantle convection simulations demonstrate how heterogeneous accumulations of previously subducted materials could form on the CMB and explain our seismic observations. We further show that subducted materials can be globally distributed throughout the lowermost mantle with variable concentrations. These subducted materials, advected along the CMB, can provide an explanation for the distribution and range of reported ULVZ properties.

So called S waves (see this) are transversal acoustic waves caused by the shear force parallel to the propagation. This force is proportional to viscosity and is negligible in liquids but much larger in solid phase waves reflected at mantle-core boundary. The core of Earth is in a liquid phase. Therefore sound waves from the surface of Earth are reflected back at the mantle-core boundary.

This makes it possible to deduce information from the structure of the mantle-core boundary and it has turned out that it has a highly complex structure. First of all, these waves propagate very slowly. This allows us to conclude that there is a relatively thin layer with a high density, which could consist of the same material as the seabed. This layer contains mountains with heights of order 10 km.

The TGD inspired view of the evolution life, inspired by the Cambrian Explosion and TGD based view of cosmology, is that photosynthesizing life evolved in underground oceans and that

the expansion of the Earth radius by about factor 2 bursted these oceans to the surface of Earth in Cambrian Explosion [L3, L11, L7, L21, L20].

Could the ancient seabed above the mantle be assigned to the ocean immediately above the mantle? This is not possible. The existence of an underground ocean immediately above the mantle is impossible due to the high pressure and temperature so that the convection remains the natural explanation for the presence of seabed.

The second objection is that life in the underground oceans is not possible because solar energy needed by photosynthesis is not available. How could photosynthesis have developed in the underground oceans? The key observation is that energies of the photons of thermal radiation coming from the core are of the same order as the metabolic energy currency with nominal value of .5 eV: could this radiation have served as a source of metabolic energy.

How would this energy be transferred? The Pollack effect [I9, L1, I13, I11] and its reversal, whose TGD based understanding [L1, L19, L24, L26, L25] has increased considerably during this year, could provide a fast energy transfer mechanism, but in its standard form the Pollack effect requires liquid water. Could the so-called superionic ice [D2], which has been speculated to be found even near the mantle of Earth, make possible the analogy of the Pollack effect?

3.4.2 Ordinary water cannot survive near mantle

Although it is obvious that ordinary liquid water cannot exist at temperatures and pressures prevailing near the mantle, it is useful to look at the situation more quantitatively.

In mechanical equilibrium, pressure gradient and the gravitational force, expressible in terms of the gradient of gravitational potential, cancel each other in good approximation. One can estimate the change of pressure as $\Delta p = \rho \Delta \Phi_{gr} = \rho G M \Delta(\frac{1}{R})$. The equation of state allows an estimate for ΔT .

Pressure is estimated to increase from 100 MPa at the surface of the Earth to 139 GPa above the mantle, that is by a factor 1000. Temperature, converted to thermal energy $E = kT$, is estimated to increase from .03 eV to 0.42 eV. The increase is by a factor of 10. Ordinary water cannot survive in this kind of environment so that underground water is possible only sufficiently near to the surface of Earth.

Could one imagine a phase of water allowing the analog of Pollack effect so that the transformation of protons to dark protons at the gravitational MB could make it possible to transfer metabolic energy to the higher heights, where underground liquid water can exist. This would have made possible the development of photosynthesizing life and would also solve the "faint Sun" paradox (<https://rb.gy/mfhavz>) [L21] meaning that the solar energy feed was not enough for the metabolic needs of life at the surface of Earth.

3.4.3 Pollack effect for superionic water and metabolic energy feed from the core of Earth

Superionic ice [D2] (see this and this) existing at extreme pressures. The density of superionic ice is slightly less than 4 times the density of ordinary ice. In superionic ice O^{2-} ions form a lattice whereas H^+ ions float freely. This phase is conductor with H^+ ions serving as charge carriers. Superionic ice is proposed to appear in the mantles of giant planets such as Uranus and Neptune and in [?] the possibility that it could occur in the Earth's mantle was considered.

Could water appear as superionic ice above the Earth's core and allow Pollack effect and its reversal so that gravitational flux tubes would carry dark protons. Could dark photons emitted in the reverse Pollack effect transfer the energy along gravitational flux tubes to the underground oceans near the surface of the Earth?

Let's assume that there exists superionic ice above the mantle.

1. Could the radiation from the core kick part of the protons of the superionic water to the gravitational magnetic body? The gravitational binding energy of protons at the surface of Earth is about .5 eV and now roughly by a factor 4 larger, that is 2 eV, at the top of the mantle. At the gravitational magnetic flux tubes the reduction of gravitational binding energy is therefore below 2 eV. The temperature of the core corresponds to the metabolic energy currency of about .4 eV so that the radiation could have played the same role as the solar radiation in photosynthesis.

2. If the reverse Pollack effect occurs, dark photons are emitted and they propagate to the MBs of water volumes near the surface of Earth and could provide energy for photosynthesis. Also time reversal can occur for the water near the surface of Earth and the proton can gain the energy required by darkness by emitting a negative energy dark photon propagating to the MB near the mantle. I have called this mechanism remote metabolism or quantum credit card and asked whether it could play a key role also in the ordinary biology.
3. If the temperatures of the lower part of the mantle and the core are the same, the energy input from the core could feed protons to gravitational MB, maintain the superionic water phase and compensate for the energy loss due to the reverse Pollack effect. The transfer of energy near the earth's surface would take place at the speed of light and dissipation would be very small.
4. The number of ordinary-to-dark transitions of protons per unit time determines the energy flow to the MB and the energy flow to the uppermost layers of the mantle. In a steady state, this flow must be the same as the radiative heat flow from the core. This transfer rate is determined by the rate for the photon absorptions kicking protons to the MB. The energy flow of energy coming as radiation is proportional to T^4 .

3.4.4 A mechanism of photosynthesis which does not involve biomolecules

Standard biology teaches us that photosynthesis is needed to produce oxygen, which is the basic prerequisite of life. Besides complex biological apparatus this requires photons, which provide the needed energy. At the bottom of the ocean there is very dark and this might form a bottleneck for the evolution of life. Now it has been found that at the bottom of ocean mineral deposits known as polymetallic nodules can generate oxygen in absence of photons (see). They contain combinations of cobalt, copper, lithium, and manganese and the size of the nodule can be that of a human hand. The initiation of electrolysis splitting water to hydrogen and oxygen needs only 1.5 eV voltage in seawater. This means that one has a battery. It was found that the nodules involve voltage as high as .95 eV.

The nodules could make possible electrolysis and splitting of water. They could make it possible to overcome the hen and egg problem due the fact that a complex biomolecular apparatus is needed for photosynthesis but this apparatus cannot exist in primordial biology.

In the TGD Universe, multicellular life would have evolved in underground oceans and bursted to the surface in the Cambrian explosion for about 450 million years ago [L21], which in the TGD Universe was caused by the expansion of the Earth radius by a factor 2 in a rather short period of time. TGD indeed predicts that the cosmic expansion of astrophysical objects occurs as short bursts. This explains why the astrophysical objects comove in expansion but do not expand themselves .

A heavy objection against this vision is that there are no photons in underground oceans so that photosynthesis is not possible. I have proposed that the light arriving as dark photons - ordinary photons but with a large value of effective Planck constant h_{eff} - from the Earth's core (the temperature is nearly the same as in the solar corona) could have provided the metabolic energy. Also solar photons arriving as dark photons along monopole flux tubes could have provided the energy.

It seems that also the polymetallic nodules could generate photons and make possible the splitting of water. What could be the mechanism making this possible? It must be added that also electrolysis, thought to represent ancient physics, is not a well-understood phenomenon. Remarkably, "cold fusion" was discovered in electrolytes [L2, L6]. The voltages used in electrolysis are in eV range and in atomic physics length scales they correspond to ridiculously weak electric fields. How can they cause the ionization essential for electrolysis?

1. In the Pollack effect [I9, L1, I13, I11], the irradiation of water in the presence of the gel phase generates a voltage, and therefore produces a battery. This battery also makes possible electrolysis and the splitting of water producing oxygen. Pollack effect is not understood in the framework of standard chemistry.
2. The TGD explanation is that in the Pollack effect one fourth of the protons of water are transformed to dark photons and kicked to monopole flux tubes. This creates a negatively

charged region called exclusion zone (EZ). This would generate a charge separation giving rise to the voltage. Photons would provide the needed energy to transform ordinary protons to dark protons with a larger value of h_{eff} and therefore larger energy.

It has become gradually clear that what matters is energy. Therefore the Pollack effect can be realized in several ways. In particular, the formation of molecules as bound states of atoms can provide the needed energy: no photons would be needed [L18, L28].

In particular, the reverse Pollack effect, that is dropping of dark protons from the monopole flux tubes back to ordinary protons, is also possible and would liberate ordinary photons needed in the splitting of water. This could also provide the photonic energy needed in photosynthesis and could provide a temporary storage of metabolic energy needed in photosynthesis and in the storage of energy to ATP [L19](see [this < /A >](https://tgdtheory.fi/public_html/articles/photosynthesis.html)).

3. If this can happen in the nodules, the photosynthesis could have evolved in underground oceans via the fusion of atoms to molecules and completely without external light source.

3.5 Did Earth have a ring before the Cambrian Explosion and did the rapidly expanding Earth catch the ring?

I encountered a link to a very interesting popular article "Did Earth have a ring like Saturnus?" (see this) telling about the article "Evidence suggesting that earth had a ring in the Ordovician" of Tomkins et al published in Earth and Planetary Science Letters [?].

The proposal is that the ring would have formed as a large asteroid was caught by the Earth. The tidal forces of Earth would have destroyed the asteroid so that it became a ring along the equator of the Earth. The ring created a shadow. If it formed along the equator, it could have initiated global cooling about 465 years ago: the so-called Hirnantian Icehouse followed 20 million years later. There are as many as 21 meteor strikes along the equator and this is very implausible if the meteors would have arrived from random directions.

This is a highly interesting finding from the point of view of the Expanding Earth hypothesis inspired by TGD [L21, L20]. About 524 million years ago the so-called Cambrian Explosion occurred. Highly evolved multicellular life forms suddenly emerged. A possible explanation of this mystery could be a fast expansion of the Earth: radius would have increased by about factor 2: these fast expansions could be the TGD counterpart of smooth cosmic expansion. This would have led to the bursting of underground oceans containing the multicellular life to the surface of the Earth. It is not difficult to invent objections against the idea but the new physics predicted by TGD allows to circumvent them and the model explains a large number of anomalies to the evolution of Earth.

The ring would have formed about 60 million years later and existed for a time measured 10 million years as a natural unit. Could one think that Earth had already before this time a ring and the Expanding Earth caught the ring? This could explain why the ring was along the equator, something not obvious if the ring was formed by the asteroid rotating around the Earth. This would have produced the 21 meteor strikes along the equator, a phenomenon which is extremely implausible if the meteors did not originate from the same source. The expansion of the Earth would have gradually increased the width of the shadow and the collision with the ring would have generated dust in the atmosphere and caused an additional shadowing effect causing the cooling of the climate.

3.6 Could the proposed Expansion of Earth during Cambrian Explosion have lead to the formation of Moon?

The discussions related to the Expanding Earth hypothesis stimulated interesting questions. How planets and Moons would have been born in TGD-based astrophysics and could the formation of Moon relate to the Expanding Earth hypothesis explaining also Cambrian Explosion [L21] and whether the gradual growth of Moon's orbit might relate to it?

During the last year, I wrote two articles about the birth of stars and planets and also Moons in TGD Universe last year [L22, L23].

The first basic idea is the fractality of TGD-based cosmology, which follows from the TGD view of space-time as a 4-D surface in $H = M^4 \times CP_2$. Another key idea is the replacement of a smooth continuous expansion with a sequence of fast explosions.

1. The scaled down versions of Big Bang would occur on different scales. For example, a star would produce shells of mass ejected in an explosion that would condense into planets.
2. The planets could also do the same and this would lead to the birth of shells, from these the rings would be born and from these the Moons would be born.

There are many theories about the formation of the Moon. One of them is that for about 4.5 Gy years ago a planet christened as Theia (see this) and having a mass about the mass of Mars collided with Earth and was evaporated in the process, and that the vapour condensed to form Moon (see this). 10-30 percent of Theia's mass formed Moon which is roughly 1 per cent of the mass of Earth and 70-90 per cent of mass contributed to the mass of Earth: this makes less than 10 per cent of the mass of the Earth. This would have led to the increase of the radius of the Earth and the rough estimate gives that the radius increased by a factor 2. Expanding Earth hypothesis in turn proposes that the radius of Earth increased in Cambrian explosion by factor 2.

A possibility suggested by TGD is that Moon was created by the same mechanism as a planets, that is by an explosion creating a spherical layer, which condensed to form a Moon. The condition $4\Delta R/R_E \simeq M_{Moon}/M_E$ gives $\Delta R \simeq 22$ km. Intriguingly, the group led by Weigang Liang has presented strong evidence that the Moon has turned inside out [?]. The heavy elements, which should be in the core are at the surface. For a popular summary see this. Can this model explain this mysterious looking finding? During the condensation of the spherical layer to the Moon, the gravitational acceleration experienced by the outer parts of the shell was stronger than that experienced by the inner parts. Therefore turning inside out of the shell before the condensation to Moon.

A more precise calculation shows that the turning inside out is suggestive even if the shell has a constant density.

1. Let the outer and inner radii of the spherical shell be r_{out} and r_{in} respectively. The $r_{out} - r_{in} = \Delta_{out}$ gives the thickness of the shell. Since the shell is thin, one can write $r = r_{in} + \Delta$ and perform a Taylor approximation. One can write in a good approximation for the mass of the part of the shell extending from r_{in} to $r = r_{in} + \Delta$ as

$$M_S(r = r_{in} + \Delta) = \rho 4\pi r_{in}^2 \Delta \simeq \frac{3(M_E - M_{shell})}{R_E^3} .$$

where the approximation

$$\rho = \frac{3(M_E - M_{shell})}{4\pi R_E^3} .$$

has been used.

2. The total gravitational mass affecting a particle at distance r is the sum of that caused by Earth without the shell and the portion of the shell below it and given by the sum of $M_E - M_{shell} = M_E - M(\Delta_{out})$ and $M(r)$.
3. One can write the gravitational potential as $V_{gr} = GV(r)$, where $V(r)$ is given by

$$V(r) = \frac{M_E - M_{shell} + M_S(r)}{r}$$

By expanding $V(r)$ as second order Taylor polynomial, one obtains

$$V(r) = \left(\frac{M_E - M_{shell}}{r_{in}} + \left[-\frac{M_E}{r_{in}^2} + \frac{3(M_E - M_{shell})}{R_E^3} \right] (r_{in}\Delta - \Delta^2) \right) .$$

4. The radial gravitational acceleration is given by

$$\frac{a(r)}{G} = \frac{dV}{dr} = -\left[\frac{M_E}{r_{in}^2} - \frac{3(M_E - M_{shell})}{R_E^3}\right](r_{in} + 2\Delta)$$

The first term gives constant acceleration, which cannot cause inversion. The second term gives inwards directed acceleration and can force the inversion even in the case that the density of the shell is constant.

The the radial gravitational acceleration was proportional to $d(M(r)/r)/dr = -M(r)/r^2 + (dM(r)/dr)/2 = M(-(r/d)^3 - 1)/r^2 + (4\pi(r/d)^2)/r$

In condensation the spherical shell developed a hole and contracted along the surfae of the shell to form Moon. This preserved the reverted decomposition. The outer parts containing originally lighter stuff went to the core and the heavier stuff on the inner boundary of the shell remained on the surface.

The Cambrian explosion is also an explosion. The composition of the Moon is known to be the same as that of the Earth (see this). The crazy question that comes to mind (I can already hear my colleague's laughter in my ears) is whether the Moon was born this way in the Cambrian explosion about .5 Gy ago (instead of 4.5 Gy ago). This of course does not exclude the possibility that Moon was formed in a similar explosion for 4.5 billion years ago.

Can the Cambrian option be ruled out by comparing the ages of the Earth's Moon? Radiometric age determinations give the matter making up the Earth and the Moon (not the Earth or the Moon itself!!) age estimates of 4.543 Gy and 4.46 Gy. The age difference is 80 million years.

1. The age of the material composing Moon has been deduced from the radioactive decay of Zirconium and in the latest determination the estimate increased by 40 million years (see this). This inaccuracy is of the same order as the difference in the ages of the substances! So can the Moon consist of matter, which has the same radio-active age as the Earth?

One can also critically ask why the Moon's and Earth's matter would be of different ages when the composition is the same? The most natural explanation would that the substance is the same and therefore of the same age.

2. Radioactive age determinations would therefore not rule out the hypothesis of the formation of the Moon in the Cambrian explosion. In such an explosion, a layer with a thickness of about 6 km would have been thrown out and taken with it both the life on the surface and the fossils if there were any! .5 billion years old fossils would be products of underground life!

Is there any empirical evidence that the age of the Moon *cannot* be on the order of .5 billion years. Is there any evidence for the explosive origin of the Moon? Could one compare Theia hypothesis and the two variants of TGD proposal? Could the dynamics of the Moon-Earth system help here?

1. It is known that the distance of the Moon from the Earth increases slowly: $v = 3.78$ cm per year (see this). Could the recent rate for the increase of the orbital radius be interpreted in terms of cosmic expansion? The Hubble constant is about $H = 70$ km/sMpc, where parsec (pc) is 3.26 ly. This gives for the cosmic recession velocity of Moon $v(now) = HR \simeq 2.8$ cm/y. This is 74 per cent of the observed velocity of increase for the orbital radius. This suggests that the velocity due to the explosion has gradually decreased and is approaching the cosmic recession velocity (, which increases linearly with the distance: this effect has been observed but surprisingly, has not been interpreted in terms of the cosmic recession velocity!).

Could the deviation $v - v(now)$ be a remnant of the rapid increase in the orbital radius associated with the Cambrian explosion?

2. If Moon was born in about .5 billion years ago and the velocity would have been *constant* $v = 3.78$ cm/y, the Moon would have reached a distance of about 1.9×10^7 m, which is about $2.97R_E$ (three Earth radii) from the Earth and conserably smaller than $R = 60R_E$ so that the speed should have been significantly faster at the beginning.

3. If the Moon was born in such an explosion 4.5 Gy ago, the same rough estimate assuming constant velocity $v = 3.78$ cm/y would give for the distance of the Moon $R = 26.7R_E$, $R_E = 6,357$ km. This is roughly by a factor $1/2$ smaller than the recent distance $R = 60R_E$ of the Moon. This option looks more reasonable than the Cambrian option.

Cosmic expansion cannot explain the increase of the Moon's orbital radius. One would have $dR/dt = HR$ giving the estimate $R(t) = R_E \exp(v(now)t/R)$ and $R(now) = eR_E$, which is much smaller than $R = 60R_E$.

4. Could Theia hypothesis explain the growth of the distance of the Moon's to its recent in terms of the recoil momentum gained by the evaporated fragment giving rise to the Moon? This should have made the orbit elliptic. The orbit of the Moon is slightly elliptic: the eccentricity is .055 (see this). One should also understand the mechanism, which distributed the remaining matter evenly along the surface of the Earth.

What is intriguing from the TGD point of view is that the radius of Earth could have increased by a factor 2 in the collision with Theia. This would explain the findings motivating the Expanding Earth hypothesis if the continents were formed already in the collision with the Theia.

How quickly did the ejected crust condense to form the Moon? Surprisingly, this can happen very quickly. It is estimated that the Moon could have been born within hours after the impact of Thea (see this)!

4 Some mysteries of the biological evolution from the TGD point of view

Biological evolution involves several deep mysteries. In this article 3 such mysteries are discussed.

4.1 Three mysteries

4.1.1 The mystery of life's origin deepens

Sabine Hossenfelder told about new study, which deepens the mystery of life's origin (see this). The key notion is LUCA, life's universal common ancestor, whose genome should be common to all life forms, which in the most general case involves both archaea, prokaryotes (bacteria), and eucaryotes (plants, fungi and animals).

The newest study gives a considerably larger number than the previous estimates.

1. LUCA would have 2,657 genes. Luca would have had 2.7 million bps to be compared with about 3 billion bps of humans. LUCA would have lived about 4.2 billion years ago.
2. The proteins coded by the genes of LUCA suggest that hydrogen was important in the metabolism of LUCA. Presumably LUCA lived near volcanoes. LUCA also had a rather complex metabolic circuitry and the genome suggests that it was a part of an ecosystem. The size of LUCA is $10 \mu\text{m}$ in size, which is also the size of cell nucleus, and it has a genome but no nucleus.
3. An interesting side observation is that 2,657 is prime and forms a twin prime together with 2659. Maybe number theory is deeply involved with the genome.
4. The earlier estimate for the gene number of LUCA by Bill Martin's team (see this) left only 355 genes from the original 11,000 candidates, and they argue that these 355 definitely belonged to LUCA and can tell us something about how LUCA lived.

The problem is that there are two widely different candidates for the LUCA and the new candidate seems to be too complex if one assumes a single evolutionary tree.

4.1.2 The mystery of Cambrian Explosion

Cambrian Explosion represents a long standing mystery [I12] of evolutionary biology. The basic mystery is that highly evolved multicellular life forms emerged suddenly in the Cambrian explosion about .5 billion years ago. There are much older fossils of monocellular life forms archaea and prokaryotes and they would have lived at the surface of Earth as separate evolutionary lineages.

The TGD based solution of the mystery of Cambrian Explosion does not involve ETs bringing multicellular life to the Earth [L3] [L11, L7, L21].

1. In the TGD Universe, quantum gravitation is possible in arbitrarily long scales and cosmic expansion is replaced by a sequence of quantum phase transitions occurring in astrophysical scales as very rapid local expansions between which there is no expansion.
2. The life on Earth could have evolved in two ways and as three separate evolutionary trees. Multicellular life forms possible for sexually reproducing eukaryotes would have evolved in the underground oceans, where they were shielded from meteor bombardments and cosmic rays. There are indications that underground oceans and underground life are present on Mars and possibly also some other places in the solar system.
3. In the Cambrian Explosion, identified as a short lasting rapid local cosmic expansion, the radius of Earth would have increased by a factor of two. This hypothesis was originally inspired by the observation of Adams [?] that the continents seem to fit nicely together if the radius of Earth is taken to be 1/2 of its recent radius. This hypothesis would generalize the continental drift theory of Wegener. Rather highly developed photosynthesizing multicellular life forms would have bursted to the surface of Earth from underground oceans and oceans were formed [L3] [L11, L7, L21].

The TGD proposal for the solution of the LUCA mystery relies on the solution of the mystery of the Cambrian explosion. Bacteria and archaea would have evolved at the surface of the Earth and eukaryotes having a cell nucleus and reproducing sexually in the underground oceans. Bacteria and archaea would have evolved from a counterpart of LUCA having a much smaller genome and eukaryotes would have evolved from an archaea with maximum size, which became the nucleus of the first eukaryote, LUCA.

4.1.3 Perplexing findings about asteroid Ryugu

Anton Petrov told in in Youtube video (see this) "Shocking Discovery of Earth Bacteria Inside Ryugu Asteroid Samples + Other Updates" of highly interesting recent discoveries, which might provide very strong direct evidence for the TGD view of quantum biology. Ryugu was found to contain basic amino acids and also RNA and microorganisms bacteria and microfossils resembling those living at Earth were found.

The motivation for studying asteroids is that they could have been very important in the planetary formation. The Panspermia hypothesis suggests that asteroids and similar objects could have also brought life to the Earth and the findings about Ryugu could at first be seen as a support for this hypothesis.

One question raised by the study is whether the microorganisms living at the Earth managed somehow to colonize the Ryugu asteroid sample. In standard biology this should not be possible. The alternative possibility is that the evolution in Earth and Ryugu has been very similar: Ryugu indeed orbits the Sun at an orbit for which the distance from the Sun is smaller than the orbital radius of Mars.

This would however require that the biological evolution is dictated by physics in a much longer length scale than the Earth size scale. This is just what TGD predicts. These findings provide a test for the TGD view of life which suggests a very general basic mechanism for the emergence of life [L13, L18, L27].

4.2 LUCA from the TGD perspective

In the following the explanation for the perplexing findings about LUCA are discussed from the TGD point of view.

4.2.1 Some basic facts about evolution

Let us relate the proposed identification of LUCA with 2657 genes to basic facts about biology.

1. It is known that parasitic bacteria have 500–1200 genes, free-living bacteria have 1500–7500 genes, and archaea have 1500–2700 genes. Prokaryotes and archaea are monocellular organism and do not have cell nucleus. Eukaryotes differ from Archaea and prokaryotes in that they can sexually reproduce: this means huge evolutionary step.
2. Archaea (see this) are often thought to be more primitive than prokaryotes (see this) and eukaryotes (see this but it seems that they resemble eukaryotes more than prokaryotes. It should be noticed that mitochondria (see this) responsible for the metabolism have their own genome having only 37 genes to that one could (just for fun) argue that mitochondria are a natural identification of LUCA!
3. The lower bound for the gene number of free living bacteria and archaea is 1500 and smaller than 2657. It is however known that ancient bacteria had more genes than their recent forms. This might be due to the fusion of the genes to longer genes. They could be also predecessors of LUCA in some sense.

4.2.2 The number of genes for LUCA is upper bound for the genes of archaea: what does this mean?

Intriguingly, 2657 is near the upper limit 2700 for the gene number of archaea. This raises questions.

1. Could the branches of the evolutionary tree have genuine dead ends, and are archaea and bacteria such dead ends? Could prokaryotes and archaea correspond to separate lineages of the evolutionary tree and did eukaryotes evolve from archaea as the cell nucleus emerged?
2. Was the gene number 2657 a critical gene number distinguishing between archaea and higher life forms? Could this be the critical gene number above which the cell nucleus possessed by eukaryotes becomes possible and makes possible sexual reproduction and explosive evolution of new life forms. Did the maximally complex archaea with 2657 genes become the first eukaryote nucleus, LUCA?

If this was the case, did prokaryotes really share the genes of LUCA? Also prokaryotes and archaea with shorter genomes exist. Are these predecessors of the nucleus of LUCA? Did prokaryotes with gene number larger than 2657 evolve from prokaryotes separately?

3. Do prokaryotes and archaea have a common predecessor analogous to LUCA having much lower number of genes and is it the previous candidate for the LUCA having 355 genes.

4.2.3 Two mysteries with a common solution: Cambrian Explosion and the complexity of LUCA

The complexity of the genome of LUCA looks like a mystery as also the existence of two widely different candidates for LUCA. One possible solution of the mystery relies on the Panspermia hypothesis. Most life forms would have arrived to the Earth from elsewhere and only sufficiently complex organisms, whose genomes contained the LUCA genome, survived in the new environment. The TGD based solution of the mystery does not involve ETs but relies on the TGD inspired solution of the mystery of Cambrian Explosion [L3] [L11, L7, L21].

In this framework, one can imagine three separate evolutionary lineages.

1. Archaea and prokaryotes such as cyanobacteria, about which there are fossils much before the Cambrian Explosion, would have evolved at the surface of Earth. Note that cyanobacteria have about 559 core-genes (see this).
2. More complex multicellular eukaryotes, having a cell nucleus and reproducing sexually, would have evolved in the underground ocean oceans. The basic evolutionary step would have been the emergence of the cell nucleus, perhaps as a fusion of two monocellulars. This step could have led from LUCA as a maximally complex archaea to eukaryotes. For some reason,

eukaryotes would have survived only in underground oceans. Cosmic rays causing damage to the genes inside it might have made sexual reproduction too risky.

3. Eukaryotes would have had the proposed LUCA as a common ancestor as the maximally complex archaea whereas the earlier candidate for LUCA would be the common ancestor of archaea and prokaryotes.
4. Either all the organisms of the underground oceans contained the maximally complex archaea genome or only the organisms of the underground oceans having a genome not shorter than the LUCA genome were sufficiently complex to survive besides the prokaryotes present from the beginning at the surface of Earth.

LUCA as the first eukaryote, having the maximally evolved archaea as cell nucleus, would have had just the minimum number of genes needed to survive at the surface of the Earth. The archaea and bacteria having a genome shorter than LUCA genome would have emerged in the evolution at the surface of the Earth before the Cambrian Explosion.

5. A fusion of two unicellulars took place in the formation of LUCA and outer cell membrane emerged. An archaea with the genes of LUCA became the nucleus of the first eukaryote cell. The minimal choice for the second unicellular is mitochondria with 37 genes: this would give 2694 genes, almost the upper bound of 2700 for archaea. If cell membranes are an outcome of self-organization rather than being coded, this system might have been able to code itself.

4.3 The perplexing findings about the asteroid Ryugu from the TGD perspective

The TGD perspective about Ryugu is based on the notion of long range quantum coherence predicting the notions of gravitational and electric magnetic body carrying phases of ordinary matter with a very large value of Planck constant [L14, L13, L18, L27]. These field bodies with sizes in astrophysical scales are predicted to be fundamental for life and its evolution and imply that the evolution of the chemical life is predicted to be induced by the evolution at the level of the field bodies. In this framework one can understand why Ryugu can contain life forms resembling those on Earth.

4.3.1 Some facts about Ryugu

Some basic facts about Ryugu are in order. Consider first the origin of Ryugu.

1. The surface of Ryugu is very young and has an age of 8.9 ± 2.5 million years. The composition of Ryugu shows that its material has been at a rather high temperature about 1000 C and presumably near the Sun. Eventually Rygu would have left the inner solar system and its composition suggests that it has been very near to the Kuiper belt with distance 30-55 AU.
2. The asteroid that arrived near the Earth from outer space must have been for a long period in complete darkness. The object giving rise to Ryugu could have originated far from Jupiter, possibly near the Kuiper belt. Some compounds in Ryugu can only form near the Kuiper belt. A larger object of radius about 100 km could have suffered a collision near Earth and produced Ryugu with a size of 10 km near Earth.
3. Recently Ryugu orbits the Sun at a distance of 0.96-1.41 AU once every 16 months (474 days (16 months); semi-major axis of 1.19 AU). Note that the distance of Mars from the Sun is about 1.5 AU. Its orbit has an eccentricity of 0.19 and an inclination of 6 degrees with respect to the ecliptic.

The circumstances at Ryugu are favorable for life.

1. The highest temperature on the Ryugu asteroid reaches 100 degrees C, while the coldest regions sit at about room temperature. Temperatures also change depending on the solar distance of the asteroid, lowering as Ryugu moves further away from the Sun. This would

mean that the circumstances at Ryugu become favourable for life as it passes Earth. The lowering of the temperature at a large distance would not be fatal.

Hydration is essential for life. The required range of dehydration reaction temperature decreases with increasing substitution of the hydroxy-containing carbon: Primary alcohols: 170–180 degrees C; secondary alcohols: 100–140 degrees C; tertiary alcohols: 25 degrees–80 degrees C. Primary/secondary/tertiary refers to the position of -OH substitution in Carbon atom.

2. Ryugu contains liquid water and also carbonated water. Coral-like inorganic crystals are present. The sample contained carbon rich molecules, amino acids and components of RNA and hydrated compounds! Ammonium.
3. It has also been found that Ryugu contains phosphorus rich samples. Phosphorus plays a central role in metabolism and in the "dark" realization of the genetic code in TGD. The abstract of the article [I5] summarizes the findings.

Parent bodies of C-type asteroids may have brought key volatile and organic-rich compounds to the terrestrial planets in the early stages of the Solar System. At the end of 2020, the JAXA Hayabusa2 mission successfully returned samples from Ryugu, providing access to a primitive matter that has not suffered terrestrial alteration. Here we report the discovery of a peculiar class of grains, up to a few hundreds of micrometres in size, that have a hydrated ammonium–magnesium–phosphorus (HAMP)-rich composition. Their specific chemical and physical properties point towards an origin in the outer Solar System, beyond most snow lines, and their preservation along Ryugu history. These phosphorus-rich grains, embedded within an organic-rich phyllosilicate matrix, may have played a major role when immersed in primitive terrestrial water reservoirs. In particular, in contrast to poorly soluble calcium-rich phosphates, HAMP grains favour the release of phosphorus-rich and nitrogen-rich ionic species, to enter chemical reactions. HAMP grains may have thus critically contributed to the reaction pathways of organic matter towards a biochemical evolution.

4.3.2 Objections against the Panspermia hypothesis as explanation of the findings about Ryugu

The panspermia hypothesis states that Ryugu and similar objects could have served as a source of life on Earth.

1. Overpopulation problem is the theoretical objection against the Panspermia hypothesis. No new forms of life are possible since no niches are left untouched.
2. There is also a second objection against the panspermia hypothesis as an explanation of these findings about Ryugu. It has been claimed that the Ryugu sample was contaminated by terrestrial microorganisms [I3] (see this). Nitrogen dioxide NO_2 is used in sterilization meant to remove, kill, or deactivate all forms of life present in fluid or on a specific surface. Life forms of Earth should not be able to colonize samples under extremely sterile conditions. If contamination occurred, its mechanism is unknown.

The Ryugu samples contained terrestrial microbes and they evolved with time. Their DNA has not yet been identified. They resemble bacilles, which are everywhere on the Earth.

3. Microfossils have been found in meteorites [I6]. They have been found also in Ryugu but only at the surface of Ryugu and were reported to be new fossils. The reason could be that microbes have survived only at the surface of Ryugu where they receive solar light necessary for photosynthesis. The proposal of [I3] is that terrestrial organisms might by some unknown mechanism have contaminated the surface of Ryugu and produced the microfossils.

4.3.3 The TGD view of Ryugu

Neither panspermia hypothesis nor contamination look plausible in the TGD framework. Life would have evolved by the same basic mechanism both at the Earth and the asteroids and other similar objects.

1. Ryugu stays relatively near the Earth at its orbit. This could have also made possible the generation of organic matter inside the sample during the period that Ryugu has spent at its orbit around the Sun. This requires a model for how this happens and standard physics does not provide such a model.
2. The notion of the field body is central in the TGD inspired quantum biology and would act as controller of the biological body [L13, L18]. Ordinary genetic code is proposed to be accompanied by its dark variant realized at the field body for ordinary particles at it having a very large value of effective Planck constant and behaving like dark matter. Could the field body of the Earth and Sun have induced the generation of organic molecules and even bacterial life forms in the same way as they did this at the Earth?
3. The notion of the gravitational magnetic body, characterized by gravitational Planck constant introduced by Nottale [?], containing protons behaving like dark matter, represents new quantum physics relevant to the TGD inspired quantum biology. OH-O^- + dark proton qubits and their generalizations based on biologically important ions formed by salts would be the key element of life [L27] suggesting besides chemical life also other forms of life.

Any cold plasma (plasmoids as life forms) and even quartz crystals could give rise to these qubits at temperatures near the room temperature around which the flips of these qubits are possible. The difference of OH bonding energy and O^- binding energy determines the relevant energy. Its nominal value is .33 eV and is near the metabolic energy quantum of about .5 eV and near to the thermal energy .15 eV at physiological temperatures.

4. These qubits would make the matter living and life in this sense is universal. Dark genetic code is predicted and corresponds to the ordinary chemical genetic code. Basic biomolecules would give rise to analogs of topological quantum computers.

The flipping of these qubits would make quantum computation like information processing possible? Pollack effect by photon absorption can induce $\text{OH} \rightarrow \text{O}^-$ + dark proton transition and the reversal of this process and the reversal of this process can take place spontaneously. If O^- + dark proton has a lower energy than OH, it can be also induced by a presence of electric field or absorption of photons by O^- so that OH becomes the minimum energy state.

Could one understand the findings about Ryugu in this framework?

1. The presence of gravitational magnetic bodies of Earth and Sun could have induced the formation of OH-O^- qubits and more general qubits, not only at the Earth but also at Ryugu. The presence of OH bonds requires hydration and hydration is indeed possible at Ryugu.

Therefore the same mechanism could have led to the emergence of the basic organic molecules at the Earth, at Mars and inside the Ryugu asteroid and meteorites. Since the minimal distance of the Earth and Ryugu from the Sun is nearly the same, the temperature of Ryugu is near its maximal value when it is near the Earth so that the temperature would never get too hot.

2. Ryugu is under the influence of the gravitational bodies of both the Earth and the Sun. Ryugu passes near the Earth repeatedly with a period of 4 years. The organic molecules and various hydrated compounds could have gradually formed during about 10 million years as it passed near the Earth. Also bacterial life could have emerged in this way. Therefore contamination need not be in question.

Acknowledgements: I want to thank my friend Heikki Haila for enlightening discussions concerning glaciations.

REFERENCES

Theoretical Physics

- [B1] Minev ZK et al. To catch and reverse a quantum jump mid-flight, 2019. Available at: <https://arxiv.org/abs/1803.00545>.

Condensed Matter Physics

- [D1] Hansen SE et al. Globally distributed subducted materials along the Earth's core-mantle boundary: Implications for ultralow velocity zones. *Science*, 9(14), 2023. Available at: <https://www.science.org/doi/10.1126/sciadv.add4838>.
- [D2] Prakapenka VB et al. Structure and properties of two superionic ice phases. *Nature Physics*, 17(11):1–6, 2021. Available at: https://www.researchgate.net/publication/355212207_Structure_and_properties_of_two_superionic_ice_phases.

Biology

- [I1] Earth's magnetic field regions of weakness correlated to sites of political unrest and war: the paradigm quaking measurements of professor Phil Callahan. Available at: <https://www.acacialand.com/Callahan.html>.
- [I2] The Fourth Phase of Water: Dr. Gerald Pollack at TEDxGuelphU, 2014. Available at: <https://www.youtube.com/watch?v=i-T7tCMUDXU>.
- [I3] Genege MJ et al. Rapid colonization of a space-returned Ryugyu sample by terrestrial microorganisms. *Meteorites & Planetary Sciences*, 2024. Available at: <https://doi.org/10.1111/maps.14288>.
- [I4] Ostrander CM et al. Widespread seafloor anoxia during generation of the Ediacaran Shuram carbon isotope excursion. *gebiology*, 2023. Available at: <https://onlinelibrary.wiley.com/doi/10.1111/gbi.12557>.
- [I5] Pilorget C et al. Phosphorus-rich grains in Ryugu samples with major biochemical potential. *Nature Astronomy*, 2024. Available at: <https://www.nature.com/articles/s41550-024-02366-w>.
- [I6] Rozanov AYu et al. New findings of Microfossils in the Orgueil Meteorite. *Palentological Journal*, 55:111–115, 2021. Available at: <https://link.springer.com/article/10.1134/S0031030121010111>.
- [I7] Ruff SE et al. Hydrogen and dark oxygen drive microbial productivity in diverse groundwater ecosystems. *Nature Communications*, 14(3194), 2023. Available at: <https://www.nature.com/articles/s41467-023-38523-4>.
- [I8] Wotte T et al. Isotopic evidence for temperate oceans during the Cambrian Explosion. *Sci Rep.*, 9(6330), 2019. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6474879/>.
- [I9] Pollack G. *Cells, Gels and the Engines of Life*. Ebner and Sons, 2000. Available at: <https://www.cellsandgels.com/>.
- [I10] Callahan P. *Paramagnetism–Rediscovering Nature's Secret Force of Growth*, volume 2003. Acres U.S.A.. Available at: <https://www.nexusmagazine.com.>, 1995.
- [I11] Zhao Q Pollack GH, Figueroa X. Molecules, water, and radiant energy: new clues for the origin of life. *Int J Mol Sci*, 10:1419–1429, 2009. Available at: <https://tinyurl.com/ntkfhlc>.

- [I12] Gould SJ. *Wonderful Life*. Penguin Books, 1991.
- [I13] Pollack GH Zheng J-M. Long-range forces extending from polymer-gel surfaces. *Phys Rev E*, 68:031408–, 2003. Available at: <https://tinyurl.com/ntkfhlc>.

Books related to TGD

- [K1] Pitkänen M. Expanding Earth Model and Pre-Cambrian Evolution of Continents, Climate, and Life. In *Physics in Many-Sheeted Space-Time: Part I*. <https://tgdtheory.fi/tgdhtml/Btgdc1.html>. Available at: <https://tgdtheory.fi/pdfpool/expearth.pdf>, 2023.

Articles about TGD

- [L1] Pitkänen M. Pollack's Findings about Fourth phase of Water : TGD View. Available at: https://tgdtheory.fi/public_html/articles/PollackYoutube.pdf, 2014.
- [L2] Pitkänen M. Cold fusion, low energy nuclear reactions, or dark nuclear synthesis? Available at: https://tgdtheory.fi/public_html/articles/krivit.pdf, 2017.
- [L3] Pitkänen M. Expanding Earth Model and Pre-Cambrian Evolution of Continents, Climate, and Life. Available at: https://tgdtheory.fi/public_html/articles/expearth.pdf, 2018.
- [L4] Pitkänen M. Copenhagen interpretation dead: long live ZEO based quantum measurement theory! Available at: https://tgdtheory.fi/public_html/articles/Bohrdead.pdf, 2019.
- [L5] Pitkänen M. Some comments related to Zero Energy Ontology (ZEO). Available at: https://tgdtheory.fi/public_html/articles/zeoquestions.pdf, 2019.
- [L6] Pitkänen M. Could TGD provide new solutions to the energy problem? Available at: https://tgdtheory.fi/public_html/articles/proposal.pdf, 2020.
- [L7] Pitkänen M. Empirical support for the Expanding Earth Model and TGD view about classical gauge fields. https://tgdtheory.fi/public_html/articles/expearthnewest.pdf, 2021.
- [L8] Pitkänen M. Negentropy Maximization Principle and Second Law. Available at: https://tgdtheory.fi/public_html/articles/nmpsecondlaw.pdf, 2021.
- [L9] Pitkänen M. Some questions concerning zero energy ontology. https://tgdtheory.fi/public_html/articles/zeonew.pdf, 2021.
- [L10] Pitkänen M. TGD and Condensed Matter. https://tgdtheory.fi/public_html/articles/TGDcondmatshort.pdf, 2021.
- [L11] Pitkänen M. Updated version of Expanding Earth model. https://tgdtheory.fi/public_html/articles/expearth2021.pdf, 2021.
- [L12] Pitkänen M. About the number theoretic aspects of zero energy ontology. https://tgdtheory.fi/public_html/articles/ZEOnumber.pdf, 2022.
- [L13] Pitkänen M. Comparison of Orch-OR hypothesis with the TGD point of view. https://tgdtheory.fi/public_html/articles/penrose.pdf, 2022.
- [L14] Pitkänen M. How animals without brain can behave as if they had brain. https://tgdtheory.fi/public_html/articles/precns.pdf, 2022.
- [L15] Pitkänen M. Krebs cycle from TGD point of view. https://tgdtheory.fi/public_html/articles/krebs.pdf, 2022.

- [L16] Pitkänen M. TGD inspired model for freezing in nano scales. https://tgdtheory.fi/public_html/articles/freezing.pdf, 2022.
- [L17] Pitkänen M. A revolution in lithium-sulphur battery technology? https://tgdtheory.fi/public_html/articles/LiS.pdf, 2023.
- [L18] Pitkänen M. About long range electromagnetic quantum coherence in TGD Universe. https://tgdtheory.fi/public_html/articles/hem.pdf, 2023.
- [L19] Pitkänen M. About the mechanism of the energy transfer in photosynthesis. https://tgdtheory.fi/public_html/articles/photosynth.pdf, 2023.
- [L20] Pitkänen M. Empirical Support for Expanding Earth Hypothesis. https://tgdtheory.fi/public_html/articles/preCEagain.pdf, 2023.
- [L21] Pitkänen M. Expanding Earth Hypothesis and Pre-Cambrian Earth. https://tgdtheory.fi/public_html/articles/preCE.pdf, 2023.
- [L22] Pitkänen M. Magnetic Bubbles in TGD Universe: Part I. https://tgdtheory.fi/public_html/articles/magnbubble1.pdf, 2023.
- [L23] Pitkänen M. Magnetic Bubbles in TGD Universe: Part II. https://tgdtheory.fi/public_html/articles/magnbubble2.pdf, 2023.
- [L24] Pitkänen M. Pollack Effect and Some Anomalies of Water. https://tgdtheory.fi/public_html/articles/pollackwater.pdf, 2023.
- [L25] Pitkänen M. Pollack effect, lightnings and ball lightnings. https://tgdtheory.fi/public_html/articles/balllightning.pdf, 2023.
- [L26] Pitkänen M. Some new aspects of the TGD inspired model of the nerve pulse . https://tgdtheory.fi/public_html/articles/np2023.pdf, 2023.
- [L27] Pitkänen M. Quartz crystals as a life form and ordinary computers as an interface between quartz life and ordinary life? https://tgdtheory.fi/public_html/articles/QCs.pdf, 2024.
- [L28] Pitkänen M. Taos hum, stochastic resonance, and sensory perception. https://tgdtheory.fi/public_html/articles/taoshum.pdf, 2024.
- [L29] Pitkänen M. A new experimental demonstration for the occurrence of low energy nuclear reactions. https://tgdtheory.fi/public_html/articles/LENRagain.pdf, 2025.