

Pollack's Findings about Fourth phase of Water: TGD View

M. Pitkänen
Email: matpitka6@gmail.com.
<http://tgdtheory.com/>.

June 20, 2019

Abstract

The discovery of negatively charged exclusion zone formed in water bounded by gel phase has led Pollack to propose the notion of gel like fourth phase of water. In this article this notion is discussed in TGD framework. The proposal is that the fourth phase corresponds to negatively charged regions - exclusion zones - with size up to 100-200 microns generated when energy is fed into the water - say as radiation, in particular solar radiation. The stoichiometry of the exclusion zone is $H_{1.5}O$ and can be understood if every fourth proton is dark proton residing at the flux tubes of the magnetic body assignable to the exclusion zone and outside it. This leads to a model for prebiotic cell as exclusion zone. Dark protons are proposed to form dark nuclei whose states can be grouped to groups corresponding to DNA, RNA, amino-acids, and tRNA and for which vertebrate genetic code is realized in a natural manner. The voltage associated with the system defines the analog of membrane potential, and serves as a source of metabolic energy as in the case of ordinary metabolism. The energy is liberated in a reverse phase transition in which dark protons transform to ordinary ones. Dark proton strings serve as analogs of basic biopolymers and one can imagine analog of bio-catalysis with enzymes replaced with their dark analogs. The recent discovery that metabolic cycles emerge spontaneously in absence of cell support this view.

Contents

1	Introduction	1
1.1	Basic findings	2
1.2	Summary of TGD inspired model for the findings	2
2	Dark nuclei and Pollack's findings	2
2.1	Model for the formation of exclusion zones	2
2.2	Minimal metabolic energy consumption and the value of membrane potential	3
3	Fourth phase of water and pre-biotic life in TGD Universe	4
3.1	Metabolism and fourth phase of water	4
3.2	Pollack's findings in relation to osmosis and model for cell membrane and EEG	5
3.2.1	Simplest Josephson junction model for cell membrane	5
3.2.2	The counterpart of chemical potential in TGD description	6
3.3	Is time reversal involved with Pollack effect?	7
3.4	Which came first: metabolism or cell membrane?	9

1 Introduction

The discovery of negatively charged exclusion zone formed in water bounded by gel phase has led Pollack to propose the notion of gel like fourth phase of water. One can find a biographical sketch [I1] (<http://tinyurl.com/ycqtuchp>) giving a list of publications containing items related to the notions of exclusion zone and fourth phase of water discussed in the talk.

1.1 Basic findings

I list below some basic experimental findings about fourth gel like phase of water made in the laboratory led by Gerald Pollack [I2].

1. In water bounded by a gel a layer of thickness up to 100-200 microns is formed. All impurities in this layer are taken outside the layer. This motivates the term "exclusion zone". The layer consists of layers of molecular thickness and in these layers the stoichiometry is $H_{1.5}O$. The layer is negatively charged. The outside region carries compensating positive charge. This kind of blobs are formed in living matter. Also in the splitting of water producing Brown's gas negatively charged regions are reported to emerge [H2, H1].
2. The process requires energy and irradiation by visible light or thermal radiation generates the layer. Even the radiation on skin can induce the phase transition. For instance, the blood flow in narrow surface veins requires metabolic energy and irradiation forces the blood to flow.
3. The layer can serve as a battery: Pollack talks about a form of free energy deriving basically from solar radiation. The particles in the layer are taken to the outside region, and this makes possible disinfection and separation of salt from sea water. One can even understand how clouds are formed and mysteries related to the surface tension of water as being due the presence of the layer formed by $H_{1.5}O$.
4. In the splitting of water producing Brown's gas [H2, H1] having a natural identification as Pollack's fourth phase of water the needed energy can come from several alternative sources: cavitation, electric field, etc...

1.2 Summary of TGD inspired model for the findings

In the following this notion is discussed in TGD framework. The proposal is that the fourth phase corresponds to negatively charged regions - exclusion zones - with size up to 100-200 microns generated when energy is fed into the water - say as radiation, in particular solar radiation. The stoichiometry of the exclusion zone is $H_{1.5}O$ and can be understood if every fourth proton is dark proton residing at the flux tubes of the magnetic body assignable to the exclusion zone and outside it.

This leads to a model for prebiotic cell as exclusion zone. Dark protons are proposed to form dark nuclei whose states can be grouped to groups corresponding to DNA, RNA, amino-acids, and tRNA and for which vertebrate genetic code is realized in a natural manner. The voltage associated with the system defines the analog of membrane potential, and serves as a source of metabolic energy as in the case of ordinary metabolism. The energy is liberated in a reverse phase transition in which dark protons transform to ordinary ones. Dark proton strings serve as analogs of basic biopolymers and one can imagine analog of bio-catalysis with enzymes replaced with their dark analogs. The recent discovery that metabolic cycles emerge spontaneously in absence of cell support this view.

2 Dark nuclei and Pollack's findings

While listening the lecture of Pollack I realized that a model for dark water in term of dark proton sequences is enough to explain the properties of the exotic water according to experiments done in the laboratory of Pollack. There is no need to assume sequences of half-dark water molecules containing one dark proton each.

2.1 Model for the formation of exclusion zones

The data about formation of exclusion zones allows to construct a more detailed model for what might happen in the formation of exclusion zones.

1. The dark proton sequences with dark proton having size of order atomic nucleus would reside at the flux tubes of dark magnetic field which is dipole like field in the first approximation and defines the magnetic body of the negatively charged water blob. This explains the charge separation if the flux tubes have length considerably longer than the size scale of the blob which is given by size of small cell. In the model inspired by Moray B. King's lectures charge separation is poorly understood.
2. An interesting question is whether the magnetic body is created by the electronic currents or whether it consists of flux tubes carrying monopole flux: in the latter case no currents would be needed. This is obviously purely TGD based possibility and due to the topology of CP_2 .
3. This means that in the model inspired by the lectures of Moray B. King discussed above, one just replaces the sequences of partially dark water molecules with sequences of dark protons at the magnetic body of the $H1.5O$ blob. The model for the proto-variants of photosynthesis and metabolism remain as such. Also now genetic code would be realized [K2, K3].
4. The transfer of impurities from the exclusion zone could be interpreted as a transfer of them to the magnetic flux tubes outside the exclusion zone as dark matter.

These primitive forms of photosynthesis and metabolism form could be key parts of their higher level chemical variants. Photosynthesis by irradiation would induce a phase transition generating dark magnetic flux tubes (or transforming ordinary flux tubes to dark ones) and the dark proton sequences at them. Metabolism would mean burning of the resulting blobs of dark water to ordinary water leading to the loss of charge separation. This process would be analogous to the catabolism of organic polymers liberating energy. Also organic polymers in living matter carry their metabolic energy as dark proton sequences: the layer could also prevent their hydration. That these molecules are typically negatively charged would conform with the idea that dark protons at magnetic flux tubes carry the metabolic energy.

The liberation of energy would involve increase of the p-adic prime characterizing the flux tubes and reduction of Planck constant so that the thickness of the flux tubes remains the same but the intensity of the magnetic field is reduced. The cyclotron energy of dark protons is liberated in coherent fashion and in good approximation the frequencies of the radiation corresponds to multiplies of cyclotron frequency: this prediction is consistent with that in the original model for the findings of Blackman and others [J1].

The phase transition generating dark magnetic flux tubes containing dark proton sequences would be the fundamental step transforming inanimate matter to living matter and the fundamental purpose of metabolism would be to make this possible.

2.2 Minimal metabolic energy consumption and the value of membrane potential

This picture raises a question relating to the possible problems with physiological temperature.

1. The Josephson radiation generated by cell membrane has photon energies coming as multiples of ZeV , where V is membrane potential about .06 V and $Z = 2$ is the charge of electron Cooper pair. This gives $E = .12$ eV.
2. There is a danger that thermal radiation masks Josephson radiation. The energy for photons at the maximum of the energy density of blackbody radiation as function of frequency is given as the maximum of function $x^3/(e^x - 1)$, $x = E/T$ given by $e^{-x} + x/3 - 1 = 0$. The maximum is given approximately by $x = 3$ and thus $E_{max} \simeq 3T$ (in units $c = 1, k_B = 1$). At physiological temperature $T = 310$ K (37 C) this gives .1 eV, which is slightly below Josephson energy: living matter seems to have minimized the value of Josephson energy - presumably to minimize metabolic costs. Note however that for the thermal energy density as function of *wavelength* the maximum is at $E \simeq 5T$ corresponding to 1.55 eV which is larger than Josephson energy. The situation is clearly critical.
3. One can ask whether also a local reduction of temperature around cell membrane in the fourth phase of water is needed.

- (a) "Electric expansion" of water giving rise to charge separation and presumably creating fourth phase of water is reported to occur [H2, H1].
- (b) Could the electric expansion/phase transition to dark phase be adiabatic involving therefore no heat transfer between the expanding water and environment? If so, it would transform some thermal energy of expanding water to work and reduce its temperature. The formula for the adiabatic expansion of ideal gas with f degrees of freedom for particle ($f = 3$ if there are no other than translational degrees of freedom) is $(T/T_0) = (V/V_0)^{-\gamma}$, $\gamma = (f + 2)/f$. This gives some idea about how large reduction of temperature might be involved. If p-adic scaling for water volume by a power of two takes place, the reduction of temperature can be quite large and it does not look realistic.
- (c) The electric expansion of water need not however involve the increase of Planck constant for water volume. Only the Planck constant for flux tubes must increase and would allow the formation of dark proton sequences and the generation of cyclotron Bose-Einstein condensates or their dark analog in which fermions (electrons in particular) effectively behave as bosons (the anti-symmetrization of wave function would occur in dark degrees of freedom corresponding to multi-sheeted covering formed in the process).

3 Fourth phase of water and pre-biotic life in TGD Universe

3.1 Metabolism and fourth phase of water

If the fourth phase of water defines pre-biotic life form then the phase transition generating fourth phase of water and its reversal are expected to be fundamental elements of the ordinary metabolism, which would have developed from the pre-biotic metabolism. The following arguments conforms with this expectation.

1. Cell interiors, in particular the interior of the inner mitochondrial membrane are negatively charged as the regions formed in Pollack's experiments. Furthermore, the citric acid cycle (<http://tinyurl.com/y8ubjgnc>), which forms the basic element of both photosynthesis (<http://tinyurl.com/yauwzkho>) and cellular respiration (<http://tinyurl.com/ybeefxmb>), involves electron transport chain (<http://tinyurl.com/yat3m4vk>) in which electron loses gradually its energy via production of NADP and proton at given step. Protons are pumped to the other side of the membrane and generates proton gradient serving as metabolic energy storage just like battery. The interpretation for the electron transport chain in terms of Pollack's experiment would be in terms of generation of dark protons at the other side of the membrane.
2. When ATP is generated from ADP three protons per ATP flow back along the channel formed by the ATP synthase molecule (<http://tinyurl.com/yd5ndcyk>) (perhaps Josephson junction) and rotate the shaft of a "motor" acting as a catalyst generating three ATP molecules per turn by phosphorylating ADP. The TGD based interpretation is that dark protons are transformed back to ordinary ones and possible negentropic entanglement is lost.
3. ATP is generated also in glycolysis (<http://tinyurl.com/ybzgdgve>), which is ten-step process occurring in cytosol so that membrane like structure need not be involved. Glycolysis involves also generation of two NADH molecules and protons. An open question (to me) is whether the protons are transferred through an endoplasmic reticulum or from a region of ordered water (fourth phase of water) to its exterior so that it would contribute to potential gradient and could go to magnetic flux tubes as dark proton. This would be natural since glycolysis is realized for nearly all organisms and electron transport chain is preceded by glycolysis and uses as input the output of glycolysis (two pyruvate molecules (<http://tinyurl.com/y8v7aq9s>)).
4. Biopolymers - including DNA and ATP - are typically negatively charged. They could thus be surrounded by fourth phase of water and neutralizing protons would reside at the magnetic bodies. This kind of picture would conform with the idea that the fourth phase (as also

3.2 Pollack's findings in relation to osmosis and model for cell membrane and EEG

magnetic body) is fractal like. In phosphorylation the metabolic energy stored to a potential difference is transferred to shorter length scales (from cell membrane scale to molecular scale).

In glycolysis (<http://tinyurl.com/ybzgdgve>) the net reaction $C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2(g) + 6H_2O(l) + \text{heat}$ takes place. The Gibbs free energy change is $\Delta G = -2880$ kJ per mole of $C_6H_{12}O_6$ and is negative so that the process takes place spontaneously. Single glucose molecule is theoretized to produce $N = 38$ ATP molecules in optimal situation but there are various energy losses involved and the actual value is estimated to be 29-30. From $Joule = 6.84 \times 10^{18}$ eV and $mol = 6.02 \times 10^{23}$ and for $N = 38$ one would obtain the energy yield.86 eV per single ATP. The nominal value that I have used.5 eV. This is roughly 5 to 8 times higher than $E = ZeV, Z = 2$, which varies in the range.1-.16 eV so that the metabolic energy gain cannot be solely due to the electrostatic energy which would actually give only a small contribution.

In the thermodynamical approach to metabolism the additional contribution would be due to the difference of the chemical potential μ for cell exterior and interior, which is added to the membrane potential as effective potential energy. The discrepancy is however rather large and this forces the question the feasibility of the model. This forces to reconsider the model of osmosis in the light of Pollack's findings.

3.2 Pollack's findings in relation to osmosis and model for cell membrane and EEG

Osmosis (<http://tinyurl.com/gz1ck>) has remained to me poorly understood phenomenon. Osmosis means that solvent molecules move through a semipermeable membrane to another side of the membrane if the concentration of solute is higher at that side. Solute can be water or more general liquid, supercritical liquid, and even gas.

Osmosis is not diffusion: it can occur also towards a higher concentration of water. Water molecules are not attracted by solute molecules. A force is required and the Wikipedia explanation is that solute molecules approaching pores from outside experience repulsion and gain momentum which is transferred to the water molecules.

The findings of Pollack inspire the question whether the formation of exclusion zone could relate to osmosis and be understood in terms of the fourth phase of water using genuine quantal description.

In the thermodynamical model for ionic concentrations one adds to the membrane resting potential a contribution from the difference of chemical potentials μ_i at the two sides of the membrane. Chemical potentials for the ions parametrize the properties of the cell membrane reducing basically to the properties of the channels and pumps (free diffusion and membrane potential do not entirely determine the outcome).

If the transfer of ions - now protons - through cell membrane is quantal process and through Josephson junctions defined by transmembrane proteins, then the thermodynamical model can at best be a phenomenological parameterization of the situation. One should find the quantum counterpart of thermodynamical description, and here the identification of quantum TGD as square root of thermodynamics in Zero Energy Ontology (ZEO) suggests itself. In this approach thermodynamical distributions are replaced by probability amplitudes at single particle level such that their moduli squared give Boltzmann weights.

3.2.1 Simplest Josephson junction model for cell membrane

The first guess is that quantum description is achieved by a generalization of the Josephson junction model allowing different values of Planck constant at magnetic flux tubes carrying dark matter.

1. Josephson junctions correspond microscopically to transmembrane proteins defining channels and pumps. In rougher description entire cell membrane is described as Josephson junction.
2. The magnetic field strength at flux tube can differ at the opposite side of the membrane and even the values of h_{eff} could in principle be different. The earlier modelling attempts suggest that $h_{eff}/h = n = 2^k A$, where A is the atomic weight of ion, is a starting assumption deserving testing. This would mean that each ion resides at its own flux tubes.

The phase transitions changing the value of h_{eff} could induce ionic flows through cell membrane, say that occurring during nerve pulse since the energy difference defining the ratio of square roots of Boltzmann weights at the two sides of the membrane would change. Also the change of the local value of the magnetic field could do the same.

Consider first the simplest model taking into account only membrane potential.

1. The simplest model for Josephson junction defined by the transmembrane protein is as a two state system (Ψ_1, Ψ_2) obeying Schrödinger equation.

$$i\hbar_1 \frac{\partial \Psi_1}{\partial t} = ZeV\Psi_1 + k_1\Psi_2 \quad ,$$

$$i\hbar_2 \frac{\partial \Psi_2}{\partial t} = k_2\Psi_1 \quad .$$

One can use the decomposition $\Psi_i = R_i \exp(i\Phi(t))$ to express the equations in a more concrete form. The basic condition is that the total probability defined as sum of moduli squared equals to one: $R_1^2 + R_2^2 = 1$. This is guaranteed if the hermiticity condition $k_1/\hbar_1 = \overline{k_2}/\hbar_2$ holds true. Equations reduce to those for an ordinary Josephson junction except that the frequency for the oscillating Josephson current is scaled down by $1/h_{eff}$.

2. One can solve for R_2 assuming $\Phi_1 = eVt/\hbar_{eff}$. This gives

$$R_2(t) = \sin(\Phi_0) + \frac{k_1}{\hbar_1} \sin\left(\frac{eVt}{\hbar_1}\right) \quad .$$

R_2 oscillates around $\sin(\Phi_0)$ and the concentration difference is coded by Φ_0 taking the role of chemical potential as a phenomenological parameter.

3. The counterparts of Boltzmann weights would be apart from a phase factor square roots of ordinary Boltzmann weights defined by the exponent of Coulomb energy:

$$R = \sin(\phi_0) = \exp\left(\frac{ZeV(t)}{2T}\right) \quad .$$

Temperature would appear as a parameter in single particle wave function and the interpretation would be that thermodynamical distribution is replaced by its square root in quantum theory. In ZEO density matrix is replaced by its hermitian square root multiplied by density matrix.

3.2.2 The counterpart of chemical potential in TGD description

This model is not as such physically realistic since the counterpart of chemical potential is lacking. The most straightforward generalization of the thermodynamical model is obtained by the addition of an ion dependent chemical potential term to the membrane potential: $ZeV \rightarrow ZeV + \mu_I$. This would however require a concrete physical interpretation.

1. The most obvious possibility is that also the chemical potential actually correspond to an interaction energy - most naturally the cyclotron energy $E_c = \hbar_{eff}ZeB_{end}/m$ of ion - in this case proton - at the magnetic flux tube. Cyclotron energy is proportional to h_{eff} and can be rather large as assumed in the model for the effects of ELF em fields on brain.
2. This model would predict the dependence of the effective chemical potential on the mass and charge of ion for a fixed value of on h_{eff} and B_{end} . The scales of ionic chemical potential and ion concentrations would also depend on value of h_{eff} .
3. The model would provide a different interpretation for the energy scale of bio-photons, which is in visible range rather than infrared as suggested by the value of membrane potential.

The earlier proposal [K1] was that cell membrane can be in near vacuum extremal configuration in which classical Z^0 field contributes to the membrane potential and gives a large contribution for ions. The problematic aspect of the model was the necessity to assume

Weinberg angle in this phase to have much smaller value than usually. Furthermore, for proton the Z^0 contribution is negligible in good approximation so that this model does not explain the high value of the metabolic energy currency.

4. The simplest model the communications to magnetic body rely on Josephson radiation whose fundamental frequency f_J is at resonance identical with the cyclotron frequency $f_c(MB)$ at particular part of the flux tube of the magnetic body: $(f_c(MB) = f_J)$. $f_c(MB)$ corresponds to EEG frequency in the case of brain and biophotons are produced from dark EEG photons as ordinary photons in phase transition reducing $h_{eff} = n \times h$ to h .

In the modified model the sum $f_c + f_{J,n}$ ($f_{J,n} = E_J/n \times h$) of h_{eff} -independent cyclotron frequency and Josephson frequency proportional to $1/h_{eff}$ equals to cyclotron frequency $f_c(MB)$ at "personal" magnetic body varying slowly along the flux tube: $f_c + f_{J,n} = f_c(MB)$. If also the variation of f_J assignable to the action potential is included, the total variation of membrane potential gives rise to a frequency band with width roughly

$$\frac{\Delta f}{f} \simeq \frac{2f_{J,n}}{f_c + f_{J,n}} = \frac{2f_{J,1}}{nf_c + f_{J,1}} .$$

If dark photons correspond to biophotons the energy of cyclotron photon is in visible and UV range one has $nf_c = E_{bio}$ and

$$\frac{\Delta f}{f} \simeq \frac{2ZeV}{E_{bio} + ZeV} .$$

The prediction is scale invariant and same for all ions and also electron unless E_{bio} depends on ion. For $eV = .05$ eV, $Z = 1$, and $E_{bio} = 2$ eV ($f \simeq 5 \times 10^{14}$ Hz) one has $\Delta f/f \sim .1$ giving 10 per cent width for EEG bands assumed in the simpler model.

If this vision is on the correct track, the fundamental description of osmosis would be in terms of a phase transition to the fourth phase of water involving generation of dark matter transferred to the magnetic flux tubes. For instance, the swelling of cell by an in-flow of water in presence of higher concentration inside cell could be interpreted as a phase transition extending exclusion zone as a process accompanied by a phase transition increasing the value of h_{eff} so that the lengths of the flux tube portions inside the cell increase and the size of the exclusion zone increases. In general case the phase transitions changing h_{eff} and B_{end} by power of two factor are possible. This description should bring magnetic body as part of bio-chemistry and allow understanding of both equilibrium distributions, generation of nerve pulse, and basic metabolic processes leading to the generation of ATP.

3.3 Is time reversal involved with Pollack effect?

EZs have the very strange property that the impurities are spontaneously removed from them. This seems to be in conflict with the second law of thermodynamics according to which both temperature and concentration gradients should tend to disappear. Could one understand this as being due to a reversal of the arrow of time?

Indeed, TGD inspired theory of consciousness relying on zero energy ontology (ZEO) predicts the possibility of time reversed selves [?]. When conscious entity - self - dies, it reincarnates as a self with opposite arrow of geometric time.

1. In ZEO zero energy states replace ordinary quantum states assigned with time=constant snapshots of time evolution in space-time. Zero energy states are pairs of ordinary quantum states at opposite light-like boundaries of causal diamond (CD) identifiable as counterparts of initial and final states of a physical event. Conservation quantum numbers translates to a mathematical statement that the quantum numbers associated with the members of pairs are opposite. One can also say that zero energy state is analogous to a deterministic computer program or a behavioral mode. The act of free will replaces this program/behavior with a new one so that one avoids the paradox between the non-determinism of free will and determinism of physics.

2. Causal diamond (CD) defines the imbedding space correlate of self. One can assign to the opposite light-like boundaries the attributes active and passive. During the sequence of analogs of “small” state function reductions analogous to weak quantum measurements (resembling classical measurements) the passive boundary remains unaffected as also the members of state pairs defining zero energy states associated with it. Active boundary recedes farther away from the passive boundary and the members of state pairs at it change. The size of CD thus increases and gives rise flow of geometric time as an increase of the temporal distance between the tips of CD.
3. Eventually the first state function reduction to the opposite boundary of CD must occur, and active and passive boundary change their roles. Self dies and re-incarnates as a self with opposite arrow of geometric time: the formerly passive boundary of CD becomes now active and moves in opposite time direction reduction by reduction. In the next re-incarnation self continues almost from the moment of geometric time at which it died. It might be that we die repeatedly without noticing it at all!
4. The many-sheeted space-time approximated with slightly curved regions of Minkowski space would certainly tend to mask the time reversals in given length scale. In elementary particle length scales the state function reductions would indeed change the arrow of time but this would occur so often that there would be no arrow of time in statistical sense: one would speak of microscopic reversibility. In time scales considerably longer than those of human consciousness the observed arrow of time would correspond to that associated with selves with very large CDs and with lifetime much longer than ours. The change of the arrow of time could be detectable in time scales relevant to living matter and human consciousness and just these scales are the scales where the anomalies occur!

Could the ghostly space-time regions - time reversed selves - have some physical signatures making possible to prove their existence empirically?

1. Second law would still hold true but in opposite direction of geometric time for the space-time sheets with non-standard arrow of time. The effects implied by second law would be present as their reversals. The observer with standard direction of geometric time would see temperature and density gradients to develop spontaneously. Also parameters describing dissipation rates such as Ohmic resistance and viscosity could have in some situations negative values.

This indeed seems to take place in living matter. For instance, the building bricks of molecules spontaneously arrange to molecules: DNA replication, transcription and translation of RNA to proteins are basic examples about this. The development of concentration gradients is also clear in the strange ability of EZs to get rid of impurities. Also the charge separation creating EZs could be seen as disappearance of charged separation in reversed direction of time. Healing of living organism could be a basic example of the process in which the arrow of time changes temporarily at some level of hierarchy of space-time sheets.

2. The generation of temperature gradients would be a clear signature for the reversal of the arrow of time. Water is the basic stuff of life, and the thermodynamics of water involves numerous anomalies summarized at Martin Chaplin’s homepage “Water structure and science” (see <http://tinyurl.com/ye77f7d>). TGD based explanation could be naturally in terms of dark variants of protons at magnetic flux tubes and possible change of the arrow of geometric time.
3. There is a lot of anecdotal evidence for the effects challenging our beliefs about standard arrow of time. A spontaneous generation of temperature differences is basic example. There is a nice popular document about this boundary region of science by Phie Ambo (see <http://tinyurl.com/yaog8h39>), which even skeptic might enjoy as art experience.

It was a great surprise for me that one of the key personalities in the document is Holger B Nielsen, one of the pioneers of string models. I have had the honor to have intense discussions with him in past: he is one of the very few colleagues who has shown keen interest on the basic ideas of TGD. The document discusses strange phenomena associated with the physics

of water possibly having interpretation in terms of time reversal and formation of EZs. From the document one also learns that in Denmark physics professionals are beginning to take these anomalies seriously.

Unfortunately, the people who claim having discovered this kind of effects - often not science professionals - are labelled as crackpots. The laws of science also tell what we are allowed to observe (and think), at least if we want to be called scientists!

4. The ghost stories might also reflect something real - this real need of course not be ghost but something deep about consciousness. Could it be that it is sometimes possible to consciously experience the presence of a space-time region - self - with an opposite arrow of geometric time? Ghost stories typically involve a claim about the reduction of temperature of environment in presence of ghost: could this be something real and a signature for the reversal of time at some level of dark matter hierarchy affecting also dark matter? As a matter of fact, in TGD Universe our conscious experience could involve routinely sub-selves (mental images) with non-standard arrow of time [?]: motor actions could be identified as sensory mental images with opposite arrow of time.

3.4 Which came first: metabolism or cell membrane?

One of the basic questions of biology is whether metabolism preceded basic biopolymers or vice versa. RNA world scenario assumes that RNA and perhaps also genetic code was first.

1. The above view suggests that both approaches are correct to some degree in TGD Universe. Both metabolism and genetic code realized in terms of dark proton sequences would have emerged simultaneously and bio-chemistry self-organized around them. Dark proton sequences defining analogs of amino-acid sequences could have defined analogs of protein catalysts and played a key role in the evolution of the metabolic pathways from the primitive pathways involving only the phase transition between ordinary water and fourth phase of water.
2. There is very interesting article (see <http://tinyurl.com/ycdhd4fd> [?]) reporting that complex metabolic pathways are generated spontaneously in laboratory environments mimicking hot thermal vents. Glycolysis and pentose phosphate pathway were detected. The proposal is that these pathways are catalyzed by metals rather than protein catalysts.
3. In standard biology these findings would mean that these metabolic pathways emerged before basic biopolymers and that genetic code is not needed to code for the metabolic pathways during this period. In TGD framework dark genetic code [K2, K3] would be there, and could code for the dark pathways. Dark proton strings in one-one correspondence with the amino-acid sequences could be responsible for catalysts appearing in the pathways. Only later these catalysts would have transformed to their chemical counterparts and might be accompanied by their dark templates. One cannot even exclude the possibility that the chemical realization of the DNA-aminoacid correspondence involves its dark analog in an essential manner.

REFERENCES

Fringe Physics

- [H1] M. B. King. Free-energy devices. Available at: <http://www.free-energy-info.co.uk/MorayKing.pdf>.
- [H2] King MB. Water Electrolyzers and the Zero-Point Energy. *Phys Procedia* . Available at: <http://www.sciencedirect.com/science/journal/18753892>, 20:335–445, 2011.

Biology

- [I1] Pollack Laboratory- Biographical Sketch. Available at: <http://faculty.washington.edu/ghp/cv/>.
- [I2] The Fourth Phase of Water : Dr. Gerald Pollack at TEDxGuelphU. Available at: <https://www.youtube.com/watch?v=i-T7tCMUDXU>, 2014.

Neuroscience and Consciousness

- [J1] Blackman CF. *Effect of Electrical and Magnetic Fields on the Nervous System*, pages 331–355. Plenum, New York, 1994.

Books related to TGD

- [K1] Pitkänen M. General Theory of Qualia. In *Bio-Systems as Conscious Holograms*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/holography.html#qualia>, 2006.
- [K2] Pitkänen M. Homeopathy in Many-Sheeted Space-Time. In *Bio-Systems as Conscious Holograms*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/holography.html#homeoc>, 2006.
- [K3] Pitkänen M. Nuclear String Hypothesis. In *Hyper-finite Factors and Dark Matter Hierarchy*. Online book. Available at: <http://www.tgdtheory.fi/tgdhtml/neuplanck.html#nuclstring>, 2006.