

# Holography=holomorphy vision in relation to quantum criticality, hierarchy of Planck constants, and $M^8 - H$ duality

June 28, 2024

Matti Pitkänen

orcid:0000-0002-8051-4364.

email: matpitka6@gmail.com,

url: [http://tgdtheory.com/public\\_html/](http://tgdtheory.com/public_html/),

address: Rinnekatu 2-4 A 8, 03620, Karkkila, Finland.

## Abstract

Holography = holomorphy vision generalizes the realization of quantum criticality in terms of conformal invariance. Holography = holomorphy vision provides a general explicit solution to the field equations determining space-time surfaces as minimal surfaces  $X^4 \subset H = M^4 \times CP_2$ . For the first option the space-time surfaces are roots of two generalized analytic functions  $P_1, P_2$  defined in  $H$ . For the second option single analytic generalized analytic function defines  $X^4$  as its root and as the base space of 6-D twistor twistor-surface  $X^6$  in the twistor bundle  $T(H) = T(M^4) \times TCP_2$  identified as a zero section

By holography, the space-time surfaces correspond to not completely deterministic orbits of particles as 3-surfaces and are thus analogous to Bohr orbits. This implies zero energy ontology (ZEO) and to the view of quantum TGD as wave mechanics in the space of these Bohr orbits located inside a causal diamond (CD), which form a causal hierarchy. Also the construction of vertices for particle reactions has evolved dramatically during the last year and one can assign the vertices to partonic 2-surfaces.

$M^8 - H$  duality is a second key principle of TGD.  $M^8 - H$  duality can be seen a number theoretic analog for momentum-position duality and brings in mind Langlands duality.  $M^8$  can be identified as octonions when the number-theoretic Minkowski norm is defined as  $Re(o^2)$ . The quaternionic normal space  $N(y)$  of  $y \in Y^4 \subset M^8$  having a 2-D commutative complex sub-space is mapped to a point of  $CP_2$ .  $Y^4$  has Euclidian signature with respect to  $Re(o^2)$ . The points  $y \in Y^4$  are lifted by a multiplication with a co-quaternionic unit to points of the quaternionic normal space  $N(y)$  and mapped to  $M^4 \subset H$  inversion.

This article discusses the relationship of the holography = holomorphy vision with the number theoretic vision predicting a hierarchy  $h_{eff} = nh_0$  of effective Planck constants such that  $n$  corresponds to the dimension for an extension rationals (or extension  $F$  of rationals). How could this hierarchy follow from the recent view of  $M^8 - H$  duality? Both realizations of holography = holomorphy vision assume that the polynomials involved have coefficients in an extension  $F$  of rationals Partonic 2-surfaces would represent a stronger form of quantum criticality than the generalized holomorphy: one could say islands of algebraic extensions  $F$  from the ocean of complex numbers are selected. For the  $P$  option, the fermionic lines would be roots of  $P$  and  $dP/dz$  inducing an extension of  $F$  in the twistor sphere. Adelic physics would emerge at quantum criticality and scattering amplitudes would become number-theoretically universal. In particular, the hierarchy of Planck constants and the identification of p-adic primes as ramified primes would emerge as a prediction.

Also a generalization of the theory of analytic functions to the 4-D situation is suggestive. The poles of cuts of analytic functions would correspond to the 2-D partonic surfaces as vertices at which holomorphy fails and 2-D string worlds sheets could correspond to the cuts. This

provides a general view of the breaking of the generalized conformal symmetries and their super counterparts as a necessary condition for the non-triviality of the scattering amplitudes.

## Contents

|          |   |          |
|----------|---|----------|
| <b>1</b> | <b>Introduction</b>   | <b>2</b> |
| <b>2</b> | <b>About the evolution of the concept of <math>M^8 - H</math> duality</b>   | <b>4</b> |
| 2.1      | The problems with the original forms of $M^8 - H$ duality . . . . .   | 4        |
| 2.2      | The recent view of $M^8 - H$ duality . . . . .  | 4        |
| <b>3</b> | <b>Holography = holomorphy vision</b>   | <b>6</b> |
| 3.1      | About the geometric description of fermion pair creation vertex in the framework of catastrophe theory . . . . .      | 6        |
| 3.2      | The tension between the holography = holomorphy vision and number-theoretic vision . . . . .                          | 8        |
| 3.2.1    | Criticism of the $(P_1, P_2) = (0, 0)$ option . . . . .   | 8        |
| 3.2.2    | Does $P = 0$ option for holomorphy = holography option solve the problems? . . . . .                                  | 9        |
| 3.2.3    | $(P_1, P_2) = (0, 0)$ option and $P = 0$ option need not be inconsistent with each other . . . . .                    | 11       |
| 3.3      | Catastrophe theoretic vision of TGD . . . . .   | 11       |
| 3.4      | Holography = holomorphy vision as a realization of quantum criticality and the theory of analytic functions . . . . . | 12       |

## 1 Introduction

Quantum criticality is realized in terms of conformal invariance in string models and conformal field theories. Holography = holomorphy vision [L15] generalizes the realization of vision to the TGD framework. This view has developed considerably during the last years and provides a general explicit solution to the field equations determining space-time surfaces as minimal surfaces  $X^4 \subset H = M^4 \times CP_2$  obeying generalized holomorphy.

By holography, the space-time surfaces correspond to not completely deterministic orbits of particles as 3-surfaces and are thus analogous to Bohr orbits. This leads to zero energy ontology (ZEO) and to the view of TGD as wave mechanics in the space of these Bohr orbits inside causal diamonds (CD), which form a scale hierarchy. The construction of vertices for particle reactions identified as these singular lower-D belonging to the light-like 3-surfaces as interfaces of Minkowskian and Euclidean space-time regions defining partonic orbits has also evolved dramatically during the last year and one can assign the vertices to the partonic 2-surfaces thanks to the understanding of the modified Dirac equation [L12, L17].

There are two guesses for the vertices. Either as partonic 2-surfaces or as point-like singularities at which the monopole flux tube representing a particle splits to two. The latter interpretation looks more plausible. Fermion lines at light-like partonic orbits would correspond to poles of analytic functions, which are not vertices. At them the minimal surface property would fail. String world sheets having partonic lines as boundaries would be the analogs of cuts as singularities.

$M^8 - H$  duality is a second key principle of TGD. It has developed to its recent form [L16] rather slowly during the years via several wrong tracks [L4, L5, L14].  $M^8 - H$  duality can be seen as a number theoretic analog for the momentum-position duality and also brings in mind Langlands duality [A2, A1] [L10]. In the recent view, it is essential that  $M^8$  can be identified as octonions when its number-theoretic Minkowski norm is defined as  $Re(o^2)$ . The quaternionic normal space  $N(y)$  of  $y \in Y^4 \subset M^8$ , having a 2-D commutative complex sub-space, is mapped to a point of  $CP_2$ .  $Y^4$  has Euclidean signature with respect to  $Re(o^2)$ . The points  $y \in Y^4$  are lifted by a multiplication with a co-quaternionic unit to points of the quaternionic normal space  $N(y)$  and mapped to  $M^4 \subset H$  inversion.

This article discusses the relationship of the holography = holomorphy vision developed in [L13, L15] with the number theoretic vision predicting a hierarchy  $h_{eff} = nh_0$  of effective Planck constants such that  $n$  corresponds to the dimension of an extension rationals. The question is

whether and how this hierarchy follows from the recent view of  $M^8 - H$  duality differing considerably from the earlier view. Also the consistency with the number theoretic vision is discussed.

There are two realizations for the holography = holomorphy vision which need not be mutually exclusive.

1. For the first option, the space-time surfaces are roots of two generalized analytic functions  $f_1, f_2$  of hypercomplex coordinate (having light-like coordinate curves in  $M^4$ ) and 3 complex coordinates of  $H$ . As a special case  $f_1, f_2$  reduces to a pair  $(P_1, P_2)$  of polynomials. The minimal surface property holds true for any general coordinate invariant action constructible in terms of the induced geometry and fails only at lower-dimension singularities. This vision has tensions with the number theoretical vision. The roots  $P_1 = 0$  and  $P_2 = 0$  define 6-surfaces, whose intersection is  $X^4$ . A possible interpretation is as analogs of twistor spaces of  $M^4$  and  $CP_2$ .
2. There is also a variant of this picture in which the space-time surface is identified as a holomorphic zero section of the twistor surface  $X^6 \subset T(M^4) \times T(CP_2)$  defined as a root of single polynomial  $P$ . This option makes the nice predictions of the number theoretic vision relying strongly on polynomials with coefficients, which are rational or in an extension  $F$  of rationals.

The islands of rationals and their algebraic extensions would be selected by quantum criticality from the ocean of complex numbers. In music, the rational ratios for the frequencies of the Pythagorean scale would represent a similar phenomenon [L1, L7, L8, L11]. In this way quantum criticality defining a hierarchy of number theoretical discretization of WCW would make possible adelic physics and scattering amplitudes would become number-theoretically universal. For the  $P = 0$  option, the hierarchy of Planck constants and Galois groups as number theoretical symmetry groups would emerge as predictions. Also p-adic primes essential for p-adic mass calculations would correspond to ramified primes of  $P$ . The  $P = 0$  option seems to be however consistent with the  $(P_1, P_2) = (0, 0)$  option.

The generalization of the theory of analytic functions to the 4-D situation is suggestive.

1. The poles and cuts of analytic functions determine them as an analog of holographic data. Poles could correspond to light-like fermion lines at which the holomorphy fails and the 2-D string worlds sheets connecting partonic orbits of different monopole flux tubes could correspond to the cuts. This failure of analyticity would make non-trivial scattering amplitudes possible, and also lead to the breaking of various conformal symmetries and their super counterparts and also to the generation of an analog of Higgs vacuum expectation [L17].
2. These singularities would define at least a part of the data defining holography allowing to determine the space-time surface (also the 3-surfaces at the ends of space-time surfaces at the boundaries of CD might be needed). This picture brings in mind electrostatics in 2-dimensions where poles and cuts define point charges and line charges as sources. This is nothing but an analog of holography. The light-like fermionic lines at the light-like partonic orbits are the most natural counterparts of poles of analytic function and string world sheets define the analogs of cuts. This would mean a strong analogy with the ordinary complex analysis
3. The singularities would realize a stronger form of quantum criticality than the generalized holomorphy, and there is a rather precise geometric analogy with catastrophe theory predicting a hierarchy of critical surfaces at which some number of roots coincide (cusp catastrophe represents a simple example).
4. The splitting of a closed monopole flux tube representing a particle reaction vertex rise to a fundamental vertex for particle creation involving a creation of a fermion pair which does not seem to correspond to these singularities. The splitting process starts as a touching of the flux tube portions at parallel Minkowskian space-time sheets and generates a wormhole contact, which then splits to two wormhole contacts identifiable as "ends" of final state monopole flux tubes. This involves a splitting of the wormhole throats at a point in which a fermion pair can be created. These two touching points represent genuinely 0-dimensional

singularities defining analogs of QFT vertices having no counterpart in 2-D complex analysis. A connection with exotic smooth structures [A5, A6, A4], uniquely distinguishing 4-manifolds, and identifiable as standard smooth structure with defects, proposed to be identifiable in terms of the 2-D singularities, is highly suggestive [L9, L17].

## 2 About the evolution of the concept of $M^8 - H$ duality

$M^8 - H$  duality [L4, L5, L14] can be seen as a number-theoretic analog of momentum position duality.  $M^8 - H$  duality also relates geometric and number theoretic vision of quantum TGD and could serve as a physical realization of Langlands duality [L10]. Note that all single-fermion states are predicted to be massless in the 8-D sense and therefore their 8-D momenta are expected to reside at the 7-D light-cone of  $M^8$ .

The development of  $M^8 - H$  duality has been a rather tortuous process and has involved several wrong tracks.

### 2.1 The problems with the original forms of $M^8 - H$ duality

The very first form of  $M^8 - H$  duality mapped  $M^4 \subset H$  to a quaternionic tangent space of  $M^4 \subset M^8$ : octonions are metrically  $M^8$  with respect to the number theoretic norm  $RE(o^2)$ . The cold shower [L4, L5] was that the distribution of quaternionic tangent spaces is not in general integrable to a 4-surface  $Y^4$  and it seems that only trivial associative surfaces ( $M^4$ ) are possible.

This forced the conclusion that the normal spaces must be quaternionic, one could call this co-associativity [L4, L5]. The distribution of normal spaces is indeed known to be associative always. This however forced complexification of  $M^8$  in order to get the metric signature correctly and this led to a multitude of problems.

The form of  $M^8 - H$  duality discussed in [L4, L5] assumed the algebraic continuation of polynomials  $P(z)$  of complex variable  $z$  for which the imaginary unit commutes with octonionic imaginary units to polynomials of complexified octonion.

1.  $P(z)$  was assumed to have rational or even integer coefficients smaller than the degree. This led to the proposal that the roots of  $P(t)$  correspond to mass squared values  $m^2$  defining mass shells in  $M^4 \subset M^8$  as  $p_0^2 - \sum p_i^2 = m^2$ . This assumption had very nice implications concerning the number theoretic vision. This led to the notion of associative holography. One fixes the 3-surfaces defining the boundary data of the holography to be mass shells  $H^3 \subset M^4 \subset M^8$  and continues this data to associative 4-surfaces. The problem was that these holographic data look quite too simple. Deformations in normal direction would seem to be necessary.
2. Perhaps the most serious problem of the approach was that complex roots correspond to complex masses so that one must complexify  $M^8$ . This led to rather complicated constructions [L4, L5, L14]. The assumption that the roots are real could solve this problem but looks somewhat ad hoc.
3. The image of  $M^4 \subset H$  in  $M^8$  was identified as a co-quaternionic sub-space of complexified octonions  $M_c^8$ . The octonionic number theoretic norm is defined without the conjugation with respect to  $i$ . There are many choices for the subspace for which the octonionic coordinates are real or purely imaginary and all signatures of the number theoretic norm defined by  $o^2$  are possible. This also led to interpretational problems: to what particular  $M^4 \subset M^8$  does  $M^4 \subset H$  correspond to? Does this choice have a physical significance?

### 2.2 The recent view of $M^8 - H$ duality

In [L16], I introduced a dramatic simplification of the earlier version of the  $M^8 - H$  duality [L4, L5, L14] allowing to get rid of the complexification of  $M^8$  interpreted as an 8-D momentum space.

The trick is to define the number theoretic norm, not as  $o\bar{o}$ , but as the real part  $RE(o^2)$  of  $o^2$ : the real part here means the part of  $o^2$  proportional to the octonionic real unit. This

definition applies also to complex numbers and quaternions as subspaces of octonions. This norm is Minkowskian and allows the identification of octonions as  $M^8$  in the metric sense.

In this interpretation, the projections of the points of  $H$  to  $M^4 \subset H = M^4 \times CP_2$  correspond by  $M^8 - H$  duality to the points of  $Y^4 \subset M^8 = M^4 \times E^4$  such that the  $M^4$  projection  $m$  of the point of  $X^4$  associated with  $CP_2$  point  $s$  is mapped by inversion  $I : m^k \rightarrow \hbar_{eff} m^k / m_l m^l$  to a point of the quaternionic normal space  $N(y(s))$  as its preferred point and naturally projects to the corresponding point of  $Y^4$ .  $Y^4$  is Euclidian with respect to the number theoretic norm already described.

Number theoretic holography would be realized by requiring that the mass squared value assignable to the  $M^4$  point of the normal space of point of  $Y^4$  has the same Euclidean mass squared value as the point corresponds to so that  $Y^4$  belongs to the 7-D light-cone of octonions with Minkowskian number-theoretic norm. This correspondence would define an analogue of Wick rotation and pose a constraint on the number theoretic holography. The boundary data given at these Euclidian mass shells  $S^3$  and associativity (quaternionic normal space) would serve as a dynamical principle.

Consider first a simplified view of  $M^8 - H$  duality as I understand it now, assuming that the  $M^4$  projection of  $X^4$  is 4-D.

1. The idea is that the points of  $X^4 \subset H$  are mapped to the points of  $Y^4 \subset M^8$ . The  $M^4$  projection of  $X^4$  is mapped to the normal Minkowskian normal space of Euclidian surface  $E^4$ . The  $CP_2$  projection associated with a given point of  $Y^4$  parametrizes the normal spaces for it. The assumption that the normal spaces are quaternionic and contain a Minkowskian 2-D commutative space implies that they can be parametrized by  $CP_2$ . The normal spaces and 2-D subspaces form integrable distributions and determine  $Y^4$  by number theoretic holography using the integrable distribution of Minkowskian normal spaces as holographic data located at 3-D surfaces. The tangent spaces of  $Y^4$  have an Euclidian number theoretic metric. The points of  $Y^4$  are in 1-1 correspondence with those of  $CP_2$  except at the singularities at which the normal space is not unique.
2. The 4-surface  $Y^4 \subset M^8$  is Euclidean. All points of  $Y^4$  have as quaternionic normal space isomorphic to  $M^4 \subset M^8$  containing  $M^2$ , which is complex and commuting sub-space of octonions. The normal space  $N(y)$  of  $y \in Y^4$  as  $M^4$  is not unique but this has a geometric interpretation and does not lead to the problem of non-uniqueness of  $M^4 \subset M^8$  as the previous view of  $M^8 - H$  duality.

The normal space contains the  $M^4$  point associated with the point of  $CP_2$  as a preferred point having an interpretation as 4-momentum. This point is "active" if there is a fermion at the corresponding point of  $X^4$ . One has a geometric representation of the many-fermion state in  $Y^4$  mapped to the points of  $M^4$  projection of  $X^4$  by  $M^8 - H$  duality.

3. The new form of  $M^8 - H$  duality requires a lift of the point  $y \in Y^4$  to a point of its quaternionic and Minkowskian normal space  $N(y)$ , which is mapped by inversion to  $M^4 \subset H$ . Multiplication with an imaginary unit  $e$  of  $Y^4$  would perform the lift but can one choose  $e$  uniquely?

Is the uniqueness really necessary? The choice of the basis of quaternionic units for the quaternionic normal space is fixed up to local  $U(2)$  rotation. The same is true for the basis of the complement. Could the interpretation be in terms of the  $M^8$  counterpart of the electroweak gauge group? Color symmetries would correspond to the  $SU(3) \subset G_2$  of octonionic automorphisms so that the standard model symmetries would be realized number-theoretically. The inverse of the lift would allow to map the points  $M^4 \subset H$  to the points of  $Y^4$  and realize the correspondence between points of  $M^4$  and  $CP_2$  for the surfaces representable as graphs of a map  $M^4 \rightarrow CP_2$ .

4. The spinor modes of fermion in  $CP_2$ , can be mapped to the spinor modes in  $Y^4$  for each fermionic momentum. Number theoretical vision suggests that the momentum components are real algebraic integers of the extensions of rationals associated with the space-time surface in question.

This picture applies when the  $M^4$  projection of  $X^4$  is 4-D. If this is not the case the situation is more complex.

1. The ideal  $CP_2$  extremal for which  $M^4$  coordinates are constant, corresponds to a singularity, where the normal space at the point of  $M^8$  is not unique: the normal open spaces span entire  $CP_2$  one has what algebraic geometers call blow up and it occurs often for algebraic surfaces. The vertex of a cone is the basic example: in this case tangent and normals spaces are not unique.

For deformed  $CP_2$  extremals  $M^4$  projection is a 1-D light-like geodesic line in  $M^4 \subset H$  and also in  $M^8$ . Along this curve, the normal spaces form give rise to a 3-dimensional surfaces of  $CP_2$ .

2. Cosmic strings also correspond to such a singularity: in a 2-D string world sheet  $X^2 \subset M^4$ , the normal spaces at a given point form a 2-D complex manifold of  $CP_2$ .
3. At singularities at which the normal space of  $Y^4$  is not unique, there are additional conditions on the allowed spinor modes since the spinor mode must have the same value for all normal spaces involved. The vanishing of the allowed spinor modes at these points would allow to satisfy these condition.
4. The number-theoretic quantization of  $M^4$  momenta requires that the momentum components are real integers in the algebraic extension of the rationals related to the region of  $X^4$  considered. The momentum unit is determined by the size scale of the causal diamond (CD).

What could be the physical interpretation of  $Y^4$ ?

1.  $Y^4$  can be sliced by the images of  $r = \text{constant}$  3-spheres  $S^3 \subset CP_2$ . Could the time evolution in  $X^4$  with respect to the light-cone proper time of  $M^4 \subset H$  have as an analog the evolution of the  $CP_2$  projection with respect to the radial coordinate  $r$  of  $CP_2$  defining a slicing of  $Y^4$  by  $M^8 - H$  duality? Or can one speak of time evolution below the scale of causal diamond CD, which implies temporal non-locality below its scale.
2. Positive and negative energy states at the half-cones of  $CD$  would be mapped to the opposite light-cones in the Minkowskian normal space of  $Y^4$ . This brings in mind Wick rotation as the Euclidization trick used in quantum field theories. Could  $M^8 - H$  duality define  $Y^4$  as a kind of Euclidization of  $X^4$ ? If so, one would have both  $M^4$  and  $CP_2$  perspective of the dynamics and also mixed perspectives (cosmic strings). The failure of  $M^4$ - and/or  $CP_2$  projection to be 4-D would force mixed perspective.

### 3 Holography = holomorphy vision

In this section holography = holomorphy vision and its relation to the theory of analytic functions, to the number theoretic vision and to  $M^8 - H$  duality will be discussed. It must be confessed that during the preparation of this section, the view of the realization of the holography = holomorphy vision simplified dramatically and a considerable portion of the text became obsolete.

#### 3.1 About the geometric description of fermion pair creation vertex in the framework of catastrophe theory

It is good to start from a physical picture. The intuitive idea is that vertices somehow correspond to partonic 2-surfaces  $X^2$  on one hand and to the points of  $X^2$  at which fermion lines turn back in time. Is the fundamental vertex a 2-D parton surface or is it point-like or are both aspects involved? This is the basic question. One must clarify what happens topologically in the creation of a fermion pair. A non-vanishing monopole flux means homological non-triviality of the partonic 2-surfaces.

1. The catastrophe theory picture implies that a given catastrophe involves a hierarchy of singularities such that the number of co-inciding roots and criticality increases. In the case of

cuspid catastrophe, the tip of the cusp would correspond to the situation in which all 3 roots of the third order catastrophe polynomial coincide.

There are good reasons to expect that also now this is the generic situation. This would suggest that a point-like vertex initiates a process leading to a creation of a pair of wormhole contacts with opposite monopole fluxes.

2. What happens that a single closed monopole flux tube splits two two monopole flux tubes? This flux tube has portions at parallel Minkowskian space-time sheets with distance of order  $CP_2$  radius and they carry opposite monopole fluxes. Topologically this process corresponds to a reconnection when the flux tubes are idealized with strings. One must generalize this to that for 3-D objects.

At the point of touch, the signature of the induced metric must be between Minkowskian and Euclidean signatures so that the induced 4 metric is degenerate. The touching point starts to grow to an Euclidean 3-D region and gives gradually rise to a wormhole contact with a vanishing total monopole flux between the throats. It has opposite wormhole throats with a vanishing total magnetic flux having a 2-D interface with Minkowskian regions. After this both wormhole throats are pinched to two parts with one common point, pinch, between them. Splitting occurs and one has two separate closed flux tubes having the wormhole contacts as their second "end".

3. This process could be also visualized in terms of the 2-D  $CP_2$  projections  $X^2$  and  $Y^2$  of two parallel flux tube portions in  $M^4$ , which fuse, evolve to a flux tube portion in  $CP_2$  and split again to 2 wormhole contacts.  $X^2$  and  $Y^2$  touch and fuse to form an intermediate wormhole contact, which evolves to two pieces with opposite monopole fluxes, which are side by side and in  $CP_2$ . The two flux tubes separate and develop a pinch, which then splits.
4. The fermion pair would be naturally created at the pinch formed in the final step. For this pinched partonic 2-surface, a global generalized complex structure is not possible since the generalized complex structures at the two parts differ by the conjugation of the hyper hypercomplex coordinate (note that also the conjugation of also  $CP_2$  coordinates might be involved). The holomorphy fails at the pinch at least. The other option is that one just does not have global smooth generalized complex coordinates. The situation would be similar also before the pinch and the holomorphy would fail at the 3-D pieces of the orbits of the throats of the intermediate worm contact.
5. Here one must ask what one means as one speaks of singularity at which the minimal surface property fails. The evolution associated with the splitting of the monopole flux tube involves a development of the opposite light-like wormhole throats as a boundary of a growing 2-D Euclidean region. This evolution defines a 3-dimensional light-like region. Does the trace of the second fundamental form have a 3-D delta function singularity?: this is in conflict with the guess that the singularities are 2-D [L17]. The final state just before the splitting is a pair of touching wormhole throats. Does this define the 2-D delta function singularity or does the touching point correspond to a point-like delta function singularity analogous to the tip of the cusp catastrophe?

It is good to make a more explicit comparison with the catastrophe theory of Rene Thom [A3].

1. One considers gradient dynamics  $dx^i/dt = \partial_i V$  and one considers possible equilibria as extrema of a potential function  $V(\{x^i\}, \{a_j\})$ . A deep result is that, although the number of the behaviour variables  $x^i$  can be large, it is possible to choose a single relevant variable, call it  $x$ . Furthermore, in suitable coordinates the potential function can be chosen to be a polynomial. A possible physical identification for the control variables is as 4 space-time coordinates or a subset of them.
2. If the number of control variables  $a_j$  is not larger than 4 there are 7 elementary catastrophes and the catastrophes for  $m$  behavior variables can be engineered from the catastrophes for  $m - 1$  behavior variables.

3. The number of sheets of the catastrophe manifold in the space spanned by the behavior variable  $x$  and control variables  $a_i$  corresponds to the number of roots to the condition  $dV/dx = 0$ . The catastrophe manifold is a  $k$ -dimensional surface for  $k$  control variables.

Different roots correspond to sheets of the catastrophe manifold and meet at lower-dimensional surfaces at which the roots are degenerate. This means criticality in which a dropping to another stable sheet driven by the rapid dynamics can occur. The critical manifolds form a hierarchy and at the tip of the many-sheeted catastrophe region all roots are identical.

### 3.2 The tension between the holography = holomorphy vision and number-theoretic vision

The relationship between holography = holomorphy vision and number theory vision involving the hierarchy of effective Planck constants and  $M^8 - H$  duality should be understood better. The updated view  $M^8 - H$  duality [L16] modifies radically the original vision [L4, L5, L14] based on polynomials  $P(o)$  of complexified octonions with rational (or equivalently integer -) coefficients obtained as a continuation of ordinary complex polynomials  $P(z)$  of variable  $z = x + iy$  with imaginary unit  $i$  commuting with the octonionic imaginary units. The earlier vision was consistent with several general ideas such as prediction of the hierarchy of Planck constants and the identification of p-adic primes as ramified primes for these kinds of the polynomials  $P$ . It also supported the notion of Galois confinement. These parts of the earlier vision should survive.

The hope is that holography = holomorphy vision allows to get naturally the roots of polynomials and corresponding algebraic extensions and even polynomials with rational or integer coefficients. This would also give ramified primes.

One needs a guiding principle and number theoretical quantum criticality is such a principle. It states that rationals and algebraic numbers correspond to islands in the ocean of complex continuum unstable under perturbations selected by quantum criticality which is the basic principle of TGD. This already implies holography = holomorphy principle but does not fix its details completely. p-Adic primes would characterize elementary particles rather than space-time regions. This suggests that the number theoretic quantum criticality is reduced to single fermion level and allows to identify light-like fermion lines at the light-like orbits of partonic 2-surfaces and assign the ramified primes and  $h_{eff} = nh_0$  to them.

#### 3.2.1 Criticism of the $(P_1, P_2) = (0, 0)$ option

There are many tensions to be resolved. The holography = holomorphy vision forces us to reconsider both the earlier view of  $M^8 - H$  duality and the number theoretical vision. The existing number theoretical vision in turn challenges the detailed realization of the holography = holomorphy vision based on  $(P_1, P_2) = (0, 0)$  option.

1. In the earlier version of  $M^8 - H$  duality [L4, L5, L14] a single polynomial  $P$  of single complex variable  $z$  with coefficients in the field of rationals (or its extension), continued to a polynomial in a complexification of octonions, defined the holographic data in turn defining the space-time surface.

Although this approach had shortcomings it also had very nice features. The dimension of the algebraic extension determined by the roots of  $P$  defined effective Planck constant and the spectrum of ramified primes of  $P$  as factors of its discriminant had interpretation as p-adic primes. The applications of TGD rely on these notions.

2. For the most obvious realization of holography = holomorphy vision a pair  $(P_1, P_2)$  of polynomials replaces a single polynomial  $P(o)$ . Is it possible to reduce the conditions  $(P_1, P_2) = (0, 0)$  to a single condition  $P(z) = 0$  for some choice of  $P$  and  $z$ ? In the recent case  $z$  would correspond to a complex coordinate at the light-like partonic 2-surface as a slice of the partonic orbit and there are very many choices.

Putting the lightlike-coordinate  $u$  to zero (restriction of fermion line at a light-like partonic orbit) one has polynomials  $P_i$  of  $w$ ,  $\xi^1$ , and  $\xi^2$  and one can choose any  $w$ ,  $\xi^1$ , or  $\xi^2$  as



dependent variable  $z$ . The degree of  $P_i$  as a polynomial of  $z$  depends on the choice of  $z$ . One can find the common 6-D roots of  $P_i$  for each choice and they correspond to the intersection of 4-D surfaces  $P_1 = 0$  and  $P_2 = 0$ . If the argument  $w, \xi^1$ , or  $\xi^2$  is an algebraic number, the roots are algebraic numbers. This leaves a lot of freedom and it is very difficult to figure out the general picture!

Therefore it is far from clear how to identify fermionic lines represented as points of  $X^2$  such that they are roots of a polynomial with coefficients in some extension of rationals. However, if can identify a unique extension  $E$  of rationals, and a unique polynomial  $P(z)$  of a highly unique variable  $z$  independent of the variables  $w, \xi^1$ , and  $\xi^2$ , its ramified primes would determine the spectrum of p-adic length scales and  $h_{eff} = nh_0$  would corresponds the degree of its Galois group.

### 3.2.2 Does $P = 0$ option for holomorphy = holography option solve the problems?

How could one solve the problems of the  $(P_1, P_2)$  option?

1. The quantum criticality of TGD suggests that there is a catastrophe theoretic hierarchy of criticalities corresponding to the surfaces  $P(z) = 0$  giving the space-time surface and as special case partonic 2-surface  $X^2$  as  $v = 0$  constant section of the partonic orbit  $X^3$ . Criticality corresponds to the coincidence of two roots so that one would have  $P(z) = 0$  and  $dP/dz = 0$  at criticality. The roots would give the intersections of the fermionic lines with the partonic 2-surface. The roots of  $P$  would define an extension of the coefficient field  $F$  of  $P$  as an extension of rationals and the ramified primes of  $P$  belonging to  $F$ .
2. The twistor lift suggests a natural identification of the coordinate  $z$ . Twistor lift replaces the space-time surface  $X^4$  with a twistor space  $X^6$  as a  $S^2$  bundle over  $X^4$ .  $X^6$  would be determined by the 6-D Kähler action.  $z$  could be identified as a complex coordinate of  $S^2$  determined up to holomorphy. Suppose that  $X^6$  is known. A natural identification of  $X^4$  is a section of twistor bundle  $X^6$  can be identifiable as a root of a polynomial  $P_{u,t}(z)$ , where  $t$  can be take to be one of the coordinates  $(w, \xi^1, \xi^2)$  (note that  $u = constant$  at  $X^3$ ). The conditions  $P = 0$  and  $dP/dz = 0$  at fermionic lines would fix the value of  $z$  and if  $t$  belongs to  $F$ , one obtains an algebraic extension of  $F$ .

The fermion line would be identified sufficiently uniquely if the choice  $P$  defining the section is sufficiently unique. In fact, different sections could define different physics. The optimistic expectation is that there is a finite number of sections or at least a finite-dimensional moduli space of sections for a given for a given twistor-surface  $X^6$ .

There are 3 obvious choices for the coordinate  $t$  corresponding to the set  $\{t_1, t_2, t_3\} \equiv \{w, \xi^1, \xi^2\}$ . Can one identify the complex coordinate  $t$  uniquely or does one obtain 3 kinds of roots also now and what could this mean?

1. If the partonic 2-surface is regarded as a Riemann surface, the natural local coordinate is  $t_k$ , and the polynomials  $P_k(z, u, t_k)$  are uniquely determined. The choice of  $t_k$  is determined apart from holomorphic bijection and Hamilton-Jacobi structure [L13] dictates the choice of  $w$  a high degree and in  $CP_2$  Eguchi-Hanson coordinates, favoured by their group theoretical properties, are natural.
2. If all 3 choices of  $t_k$  are possible, one obtains 3 kinds of roots. If the roots  $z_{k_1, i}$  and  $z_{k_2, j}$  coincide, they can correspond to the same point of  $X^4$  but need not do so.

The partonic 2-surface is many-sheeted with respect to both  $CP_2$  and  $M^4$ . Different  $w$ -roots would correspond to the multi-sheetedness with respect to  $CP_2$  and different sheets could be assigned with parallel monopole flux tubes going through the  $w$ -plane. The physical intuition suggests that, due to the small size of  $CP_2$ , the number of roots in  $CP_2$  direction is small  $CP_2$  for a given  $M^4$  point whereas in the direction of  $M^4$  the number of  $w$ -roots can be very large giving rise to a large value of  $h_{eff}/h_0 = n$ .

3. In the case of the standard twistor bundle over  $M^4$ ,  $S^2$  represents the directions of light-like geodesics emanating from a point of  $M^4$ . The twistor fibers  $S^2$  at different  $M^4$  points have

a common point if there is a light-like geodesic connecting them. This is expected to be a reasonable guess also now.

Could the  $w$ -roots correspond to intersecting twistor spheres for which the points with a different  $w$  coordinate are connected by a light-like geodesic of  $M^4$ ? Since the light-like coordinate  $u$  is constant and  $v$  is fixed, this is not plausible. This would suggest that the complex coordinate for the  $S^2$  as the cross section of the light cone boundary defines a preferred coordinate identifiable as the coordinate of the twistor sphere.

4. The same question can be posed in the case of  $CP_2$  for which the light-like geodesics are replaced with radial geodesics from the origin of Eguchi-Hanson coordinates directed to the homologically non-trivial geodesic sphere at  $r = \infty$ . The points pairs at these geodesics would have intersecting twistor spheres. Could the  $CP_2$  points associated with  $w$ -roots in the radial direction be located along these radial geodesics of  $CP_2$  so that there could be a large number of  $w$ -roots per  $CP_2$  type roots? The complex coordinate of the homologically non-trivial sphere of  $CP_2$  could serve as the analog of the coordinate of the twistor sphere.

One can argue that there is a problem with number theoretic general coordinate invariance (GCI) since the form of the  $P$  can change in a generalized holomorphism of  $H$  expected to have no physical effect. Is there a unique choice of coordinates allowing to avoid the problem?

1. For the  $(P_1, P_2)$  option, the  $X^4$  is identified as an intersection  $X_1^6 \cap X_2^6$  of 6-surfaces  $X_i^6$  as roots of  $P_i$ . Could  $X_i^6$  be identified as twistor surfaces as counterparts of the twistor spaces  $T(M^4)$  and  $T(CP_2)$  with different twistor spheres but the same base space?

If so, the complex coordinates of the twistor spheres of  $T(M^4)$  and  $T(CP_2)$  should correspond to the complex coordinates of the twistor sphere of light-like geodesics of the light-cone boundary and of radial geodesics of  $CP_2$  directed from origin to homologically nontrivial sphere of  $CP_2$  "at infinity".

2. The construction of  $X^6$  as an extremal of the 6-D Kähler action for  $X^6 \subset T(M^4) \times T(CP_2)$  [?] identifies the twistor spheres of  $T(M^4)$  and  $T(CP_2)$ . Does this mean  $X_i^6$  as twistor bundles are related by the mapping of the space of light-like geodesics of light-cone boundary and/or light-like partonic orbit to the space of radial geodesics of  $CP_2$ ?
3. This would make the situation highly unique. Holomorphies for a given choice of  $w$  *resp.*  $\xi^1$  or  $\xi^2$  would correspond to  $SO(3)$  and  $U(2)$  acting linearly on the complex coordinate. These groups reduce to  $SO(2)$  and  $U(1) \times U(1)$  by the choice of the quantization axes. The coordinate  $w$  would reduce to the complex coordinate of the twistor sphere of  $T(M^4)$  at the light-cone boundary (at least). At partonic orbits the complex coordinate  $\xi$  of the geodesic sphere of  $CP_2$  and  $w$  would be related by a map characterized by a winding number.

Some comments on the physical interpretation are in order.

1. p-Adic primes are rather large,  $M_{127} = 2^{127} - 1$  for electrons. I have proposed that one could pose constraints on the size of the polynomial coefficients, say that they are smaller than the degree of the polynomial. In this case it is not clear how to obtain such large ramified primes unless the degree of  $P$  is very large. The degree of polynomial increases exponentially in repeated iteration giving rise to an analog for the approach to chaos [L6]. This would increase the dimension of the extension.
2. As found, the polynomial  $P$  can be identified as a polynomial of Minkowski-coordinate  $w$ , or of  $CP_2$  coordinate  $\xi^1$  or  $\xi^2$ .  $CP_2$  is rather small and one expects that in  $CP_2$  directions the number of sheets is rather small so that  $P$  as a polynomial of  $\xi^1$  or  $\xi^2$  should have a rather small degree and corresponding  $h_{eff}/h_0$  should be rather small.

In  $M^4$  there is a lot of room and the degree of  $P$  as a polynomial of  $w$  can be rather large and therefore also the value of  $h_{eff}/h_0$  for these fermion lines is large and their number can be large. Therefore the corresponding ramified primes and associated p-adic length scales can be rather large in this case. It would seem that the p-adic length scale is naturally assignable to  $P(t, u, w)$ . If p-adic length scale is assignable to  $P(t, \xi^i)$ , it should be smaller than  $CP_2$  scale and could correspond to excitations of superconformal and supersymplectic algebras with mass scales which are higher than  $CP_2$  mass scale.

$P = 0$  option reduces the pair of polynomials  $(P_1, P_2)$  to single polynomial  $P$ , allows to interpret the space-time surface  $X^4$  as a section of its twistor space  $X^6$  determined by 6-D Kähler action and to identify fermion lines as surfaces ( $P = 0, dP/dz = 0$ ). This view implies the notions of effective Planck constant and ramified primes, and allows to understand number theoretical evolution as the increase of algebraic complexity in two ways: as a collective evolution of the extensions of rationals  $F$  appearing as the coefficient field of  $P$  and as the evolution at the single particle level for the polynomial  $P$ . If the polynomial  $P$  can be iterated, a connection with chaos theory [L6] emerges. Also a complexification of catastrophe theory emerges and the space-time surface is analogous to a complexification of the cusp catastrophe.

### 3.2.3 $(P_1, P_2) = (0, 0)$ option and $P = 0$ option need not be inconsistent with each other

$P = 0$  option starts from the twistor surface  $X^6$  as a known entity and determines  $X^4$  as its section whereas the  $(P_1, P_2) = (0, 0)$  option represents  $X^6$  as a solution of field equations. Therefore these options need *not* be inconsistent with each other.

1. One can start from a problem. One can argue that the  $(P_1, P_2) = (0, 0)$  option cannot represent the twistor surface  $X^6$ . However, the condition  $P_1 = 0$  or  $P_2 = 0$  defines a 6-D surface  $X_i^6$  as a solution. Could these 2 6-surfaces have interpretation as concrete representations for the  $H$  projections of two twistor surfaces  $X_i^6 \in T(M^4) \times T(CP_2)$  having a common base space  $X^4$ ?
2. How could  $X_i^6$  correspond to  $S^2$  bundle over  $X^4$ ? Could the two disjoint twistor spheres  $S^2$  be counterparts for the twistor spheres of  $T(M^4)$  and  $T(CP_2)$  having as base-spaces  $M^4$  and  $CP_2$ . Could one say that these base spaces are replaced with the projections of  $X^4$  to  $M^4$  and  $CP_2$  to get  $X^6$ . Geometrically these two spheres would correspond to the space of light-like rays emanating and to the space of  $CP_2$  radial geodesics of  $CP_2$  emanating from the point  $(m, s) \in X^4 \supset H$  and would now be concretely represented as "heavenly spheres". These spheres need not be metric spheres. Note however that the light-cone boundary as the space of light-like geodesics is metrically a 2-sphere.

In ZEO the non-trivial homology of  $M^4$  twistor sphere could be represented by the effectively missing tips of  $cd \subset M^4$ , by the axis connecting the tips of the  $cd$ , or by the light-like boundaries of  $cd$  indeed representing the space of light-rays analogs to the moment of Big Bang and Big Crunch.

### 3.3 Catastrophe theoretic vision of TGD

The analog catastrophe theoretic vision emerges in the TGD framework. One can consider this vision at the level of the space-time surface and at the level of  $H$  and they correspond to the representations of the spacetime surface as a root of two polynomials  $P_1, P_2$  and as a root of single polynomial  $P$ .

For the  $(P_1, P_2) = (0, 0)$  option, which is mathematically correct but for which the relationship to the basic number theoretical ideas is unclear, one has the following picture.

1. The vanishing of the gradient of  $V$  is replaced by the vanishing two generalized complex functions  $f_1$  and  $f_2$ , which can be also taken as polynomials  $P_i$  with coefficients, which are rational or in an extension  $F$  of rationals. The polynomial coefficients of  $P_i$  depend on the 4 generalized complex coordinates of  $H$ .
2. General coordinate invariance however reduces their number to 2 when one chooses as space-time coordinates a subset of 2 generalized complex coordinates  $(z^1, z^2)$  (hypercomplex and complex coordinate in Minkowskian regions and 2 complex coordinates in Euclidean regions). Space-time coordinates become the analogs of complex control variables and the remaining 2 generalized complex  $H$  coordinates  $(w^1, w^2)$  take the role of complex behavior variables. Instead of  $dV/dx = 0$ , one has  $P_1 = 0$  and  $P_2 = 0$ .

3. Space-time surface as the counterpart of catastrophe manifold has two generalized complex dimensions. An interesting question is whether it makes sense to regard  $(P_1, P_2)$  as a gradient  $(\partial V/\partial w^1, \partial V/\partial w^2)$  of a complex value potential function with respect to the complex behavior variables. If this is possible, one could have a complexification of the catastrophe theory and the catastrophe could correspond to a complexified cusp with two generalized complex control variables instead of 2 real control variables.

The second option, introduced in this article, is based on a single polynomial with coefficients, which are rational or in an extension  $F$  of rationals and elegantly reproduces the earlier basic ideas related to the number theoretic vision.

The catastrophe manifold is replaced with a space-time surface  $X^4$ . The condition  $dV/dx = 0$  is replaced with the condition  $P_{u,t}(z) = 0$  for generalized complex polynomial  $P$ , which has as the behavior variable  $z$  identified as the complex coordinate of the twistor sphere and the light-like hyper-complex coordinate  $u$  and complex variable  $t$  are control variables. There are 3 basic ways to choose  $t$  as  $t \in \{w, \xi^1, \xi^2\}$ .  $P$  can be interpreted as a complex gradient  $dV/dz$  of  $V = \int Pdz$  with respect to the behavior variable  $z$  so that one has  $dV_{u,t}(z)/dz = 0$ . The conditions  $(P, dP/dz) = 0$  are true for the fermion lines which represent criticality at which two sheets of the catastrophe graph coincide. Fermion lines are analogous to the folds of the cusp catastrophe. Space-time surface as the counterpart of a catastrophe manifold has 2 generalized complex dimensions as also the cusp catastrophe. An interesting question is how unique the function  $P(z)$  defining the space-time surface as a section of the twistor bundle  $X^6$  is.

It is interesting to look for the catastrophe theoretic interpretation of the already discussed model for what would happen in the splitting of a closed monopole flux tube to two monopole fluxes leading to a creation of a fermion pair.

2. In the ordinary catastrophe theory, cusp catastrophe might be enough to describe what happens. The abstract cusp catastrophe corresponds to a 2-D surface in 3-space spanned by a behavior variable and two control variables. The upper and lower sheets are stable whereas the intermediate sheet between them is unstable.
2. The stable sheets would correspond to the initial state with a single monopole flux tube and to the final state with two monopole flux tubes. The intermediate wormhole contact would correspond to the unstable sheet of the cusp. The touching of the flux tubes would lead from the initial stable sheet to the unstable sheet. The intermediate wormhole contact would then move along the unstable sheet and end up to a critical point (formation of pinch) and end up to the second stable sheet.
3. In this description, the state of the entire system corresponds to a point of the 3-D space spanned by the 2 control parameters and 1 behavior variable. Therefore it is not possible to identify the control variables of cusp as a subset of space-time coordinates and the behavior variable as an  $H$ -coordinate orthogonal to the space-time surface. Rather, the evolution of the system between the initial and final states should correspond to a part of a 4-D catastrophe manifold.

### 3.4 Holography = holomorphy vision as a realization of quantum criticality and the theory of analytic functions

A generalized complex structure can be seen as a realization of the quantum criticality of the TGD Universe just like 2-D conformal theories can be seen as a realization of criticality. The generalized complex structure of the space-time surface, or Hamilton-Jacobi structure as I call it, combines hypercomplex and complex structures into a 4-D structure [L13].

Hamilton-Jacobi structure involves an integral distribution of the local tangent space-decompositions  $M^2(x) \times E^2(x)$  allowing to assign a pair  $(u, v)$  of coordinates with light-like coordinate curves to the distribution of  $M^2(x)$  and complex coordinates  $w, \bar{w}$  to the distribution transversal spaces  $E^2(x)$ . This structure generalizes to H by introducing complex coordinates  $(\xi^1, \xi^2)$  for  $CP_2$ .

There are two options, which can be consistent. In the  $(P_1, P_2)$  option determining space-time as 4-surface, the spacetime surfaces are identified as roots of two generalized analytic functions  $f_1(u, w, \xi^1, \xi^2)$  and  $f_2(u, w, \xi^1, \xi^2)$  (polynomials in the special case) defined in  $H$  and are minimal surfaces apart from possible 2-D singularities. This option could also allow the identification of the analogs of twistor spaces of  $M^4$  and  $CP_2$  as 6-D roots of  $P_1$  or  $P_2$  having  $X^4$  as intersection and common base space. In the  $P = 0$  option space-time surface  $X^4$  is identified as a base space of the twistor-surface  $X^6$  in  $T(M^4) \times T(CP_2)$ .  $X^6$  could be determined by the  $(P_1, P_2)$  option.

This leads to the question, whether the theory of analytic functions could generalize from dimension 2 to dimension 4. I am not a mathematician in a technical sense and know almost nothing about the existing mathematical knowledge. However, I can make guesses by using a physicist's intuition.

1. In the theory of analytic functions, one basic question is whether an analytic function can be constructed from its singularities. This is essentially holography.

Electrostatics in dimension  $D = 2$  serves as an instructive physical example. The electric field can be fixed when the point charges and line charges corresponding to the poles and cuts for the analytic function are known. Only poles are obtained for rational functions. Cuts (such as  $z^{1/n}$  and  $\log(z)$ ) and essential singularities ( $\exp(1/z)$ ) are also possible. Does this picture generalize to the 4-D case?

2. The twistor Grassmannian description of the scattering amplitudes leads to the hypothesis that the amplitudes are coded by the lower-dimensional singularities of the amplitudes. Also this situation corresponds to holography in a very general sense.
3. In TGD, the construction of scattering amplitudes with the help of vertices and the associated light-like parton surfaces leads to the same kind of situation.

These observations motivate the question whether and how the theory of analytic functions in dimension 2 could generalize to dimension 4 in the TGD framework.

1. Does, for example, the representation of an analytic function in terms of its singularities generalize? Suppose that 3-D light-like parton surfaces and associated 2-D partonic 2-surfaces identified as vertices, where the generalized holomorphy breaks down, are known? Can the space-time surface be derived as a generalized holomorphic 4-surface in  $H$ , determined by two analytic functions  $f_1$  and  $f_2$  (in the generalized sense) as their roots. Could the partonic 2-surfaces be enough? Or are also string world sheets perhaps identifiable as the counterpart of cuts needed as data. What about the boundary values of the function  $f_i$  at the light-like boundaries of causal diamond CD inside which the space-time surfaces are located as analogs of Bohr orbits.
2. Hilbert's principal value theorem (see this) gives an analytic function determined in the complex plane as a sum of contributions corresponding to poles and cuts. Charges and line charges determine the electric field. In poles, the holographic information is determined by residues, and in cuts by the discontinuity of the real-analytical function across the cut.

What can one say about the singularities? Does the singularity correspond to an entire light-like partonic orbit, to a 2-D partonic surface as a section of partonic orbit, to mere light-like fermion line along it, to string world sheet, or to a point-like particle reaction vertex as a defect of a smooth structure as has been assumed hitherto?

1. The number theoretical considerations related to holography = holomorphy vision realized using  $P = 0$  option imply that the light-like fermion lines along a light-like partonic orbit are theoretically very special and analogous to the lines of criticality for cusp catastrophe.

The fermion line is a static structure and one cannot assign to it acceleration in  $M^4$  degrees of freedom as the trace of the second fundamental form defining a generalized acceleration. What could diverge is the  $CP_2$  part of the generalized acceleration having an interpretation as an analog of the Higgs field.

The light-like coordinate along the light-like fermion line is constant so that metrically it is analogous to a pole of an analytic function. On the other hand, topologically the fermion

line is analogous to a cut and it would form part of a boundary of a string world sheet having interpretation as a generalization of cut. The analogy of the fermion line as the critical line of the catastrophe graph of cusp catastrophe suggests that the trace of the second fundamental form in  $CP_2$  degrees of freedom indeed diverges at it. There would be a fold in the  $CP_2$  direction. A possible interpretation is that the fermion line serves as a source of various gauge fields.

2. There is also another kind of singularity: in a creation of a fermion pair in the splitting of closed monopole flux tube, the point at which the fermion and antifermion lines begin would correspond to singularity as an analog for an edge at which fermion turns backwards in time. This singularity is analogous to the vertex of a cusp catastrophe at which 2 folds meet.

This kind of turning point is analogous to an edge on the path of a Brownian particle. A monopole flux tube would decay to two so that the topological viewpoint of a particle emission would be in question. Also at this singularity one expects the holomorphy and minimal surface property to fail. In fact this could be true inside the entire intermediate wormhole contact formed in the process. The vision about the splitting of monopole flux tubes [L17] leads to the identification of interaction vertices as genuinely point-like singularities at which standard smooth structure has a defect so that an exotic smooth structure is obtained.

3. String world sheets are assumed to connect the 3-D partonic orbits of different monopole flux tubes. The interpretation of string as an analog of a cut is suggestive. For the cuts of the analytic function  $z^{1/n}$ , the real axis is a seat of discontinuity. By introducing the notion of  $n$ -fold covering space of the complex plane one gets rid of the discontinuity. Could the string world sheets correspond to the orbits of stringy singularities such that  $2\pi$  rotation around the singularities in the generalized complex coordinates of  $H$  lead to a point at a different space-time sheet and one obtains  $n$ -fold covering. The analogy with line charges suggests that it makes sense to assign dynamics to these objects.

$n$ -fold covering suggests an  $n$ -fold value of the effective Planck constant and the interpretation as a dark phase as  $n$ -fold covering of  $M^4$  or subspace  $M^2$ . Also  $n$ -fold coverings of  $CP_2$  or its geodesic sphere  $S^2$  are possible and would look like copies of  $M^4$  regions inside which the map from  $CP_2$  to  $M^4$  is 1-valued. Now the sheets of the covering would correspond to different copies, for instance flux tubes.

Consider now what happens in singularities from the perspective of field equations.

1. At singularities defined as loci where the minimal surface property fails, the field equations for the *entire* action are valid, but are not separately true for various parts of the action. Generalized holomorphy breaks down. The fermion lines as singularities are completely analogous to the poles of an analytic function in 2-D case and there is analogy with the 2-D electrostatics, where the poles of analytic function correspond to point charges. The fermion lines are boundaries of string world sheets and these are analogous to cuts as line charges. The cut disappears as a singularity when the complex plane is replaced with its covering and the same occurs at the space-time level.
2. The field equations at the singularity give the TGD counterparts of Einstein's equations, analogs of geodesic equations, and also the analogy Newton's  $F=ma$  (also in  $CP_2$  degrees of freedom). The generalized 8-D acceleration  $H^k$  defined by the trace of the second fundamental form, is localized on the singularities. Singularities can be seen as analogs for the sources of gauge fields. This interpretation could apply to fermion lines analogous to line charges, which correspond to cuts of an analytic function in 2-D electrostatics. At these singularities the  $CP_2$  part of the generalized acceleration  $H^k$  could diverge.

Singularities in a stronger sense can be interpreted as sources for various fermion currents and supercurrents implying their apparent non-conservation at the singularity. This kind of singularity could correspond to a vertex at which a fermion-antifermion pair is created and fermion number conservation is apparently violated since the fermion line ends. Actually

only the separate conservation of fermion and antifermion numbers is broken. Here also the  $M^4$  part of  $H^k$  would diverge.

One can pose several questions.

1. What are the counterparts of the residues, i.e. holographic data? Does the holographic data correspond to the entire 3-D parton trajectory, or only to the 2-D vertices as partonic surfaces and to the wormhole throats defining the interfaces of Minkowskian and Euclidean space-time sheets, or perhaps only light-like fermionic lines at the light-like orbits of the partonic 2-surfaces?

Under what conditions are generalized analytic functions  $f_1(u, w, \xi^1, \xi^2)$  and  $f_2(u, w, \xi^1, \xi^2)$  are determined by this holographic data? It should be possible to determine the Taylor or Laurent coefficients of  $f_i$  from the data. Can one imagine an explicit formula? Under what conditions the functions reduce to polynomials with rational (equivalently integer -) coefficients?

Generalized holomorphy could be seen as a realization of quantum criticality. Physical intuition also suggests that quantum criticality selects extensions of rationals as islands in the sea of real numbers and also makes possible number theoretical physics as adelization of TGD making possible cognitive representations as unique number-theoretical discretizations [L2, L3].

2. Should also the 2-D string world sheets connecting wormhole throats of different monopole flux tubes be included as the physical intuition suggests (these string like entities should not be confused with 4-D cosmic strings  $X^2 \times Y^2 \subset M^4 \times CP_2$  and the monopole flux tubes as their thickenings).

Are the genuinely 2-D string world sheets analogous to hypercomplex cuts? At them, the hypercomplex analytic function, which depends on the coordinate of a light-like curve of  $M^4$ , would be discontinuous. The 4-surface would be either discontinuous along the stringy curve or space-time surface could correspond to a covering of a region of  $M^4$  so that a rotation of  $2\pi$  around the stringy curve leads to another sheet and rotation of  $n2\pi$  rotation is needed in to return to the original point. I have assigned a fractional spin to this kind of situation. In  $CP_2$  degrees of freedom the same phenomenon is possible and fractional charges could be assigned with them.

3. What about the 3-surfaces at the light-like boundaries of the  $CD$  defining the "ends" of  $X^4$ . Should these 3-surfaces be given as data or could they be fixed from holographic data provided by the fermion lines and string world sheets?
4. Is it possible to have a general formula for the functions  $f_1$  and  $f_2$  (or for  $P$ ) determining the space-time surface  $X^4$  analogous to the formula of Hilbert in terms of the holographic data? This formula would be a classical analog for the determination of the scattering amplitudes from their lower-dimensional momentum space singularities in the twistor Grassmannian approach.

## REFERENCES

### Mathematics

- [A1] Frenkel E. Recent Advances in Langlands program. *AMS*, 41(2):151–184, 2004.
- [A2] Frenkel E. Lectures on Langlands program and conformal field theory, 2005. Available at: <https://arxiv.org/abs/hep-th/0512172>.
- [A3] Zeeman EC. *Catastrophe Theory*. Addison-Wessley Publishing Company, 1977.
- [A4] Etesi G. Exotica or the failure of the strong cosmic censorship in four dimensions, 2015. Available at: <http://arxiv.org/abs/1503.04945v4>.

- [A5] Gompf RE. Three exotic  $R^4$ 's and other anomalies. *J. Differential Geometry*, 18:317–328, 1983. Available at: [http://projecteuclid.org/download/pdf\\_1/euclid.jdg/1214437666](http://projecteuclid.org/download/pdf_1/euclid.jdg/1214437666).
- [A6] Asselman-Maluga T and Brans CH. World Scientific. <https://docs.google.com/document/d/1Hp656gzBkZlrHtmWvTUt0A5zsYZ415R3HHIe2eI-tP0/edit.>, 2007.

## Articles about TGD

- [L1] Pitkänen M. Geometric theory of harmony. Available at: [https://tgdtheory.fi/public\\_html/articles/harmonytheory.pdf.](https://tgdtheory.fi/public_html/articles/harmonytheory.pdf.), 2014.
- [L2] Pitkänen M. Philosophy of Adelic Physics. Available at: [https://tgdtheory.fi/public\\_html/articles/adelephysics.pdf.](https://tgdtheory.fi/public_html/articles/adelephysics.pdf.), 2017.
- [L3] Pitkänen M. Philosophy of Adelic Physics. In *Trends and Mathematical Methods in Interdisciplinary Mathematical Sciences*, pages 241–319. Springer. Available at: [https://link.springer.com/chapter/10.1007/978-3-319-55612-3\\_11](https://link.springer.com/chapter/10.1007/978-3-319-55612-3_11), 2017.
- [L4] Pitkänen M. A critical re-examination of  $M^8 - H$  duality hypothesis: part I. Available at: [https://tgdtheory.fi/public\\_html/articles/M8H1.pdf.](https://tgdtheory.fi/public_html/articles/M8H1.pdf.), 2020.
- [L5] Pitkänen M. A critical re-examination of  $M^8 - H$  duality hypothesis: part II. Available at: [https://tgdtheory.fi/public\\_html/articles/M8H2.pdf.](https://tgdtheory.fi/public_html/articles/M8H2.pdf.), 2020.
- [L6] Pitkänen M. Could quantum randomness have something to do with classical chaos? Available at: [https://tgdtheory.fi/public\\_html/articles/chaostgd.pdf.](https://tgdtheory.fi/public_html/articles/chaostgd.pdf.), 2020.
- [L7] Pitkänen M. How to compose beautiful music of light in bio-harmony? [https://tgdtheory.fi/public\\_html/articles/bioharmony2020.pdf.](https://tgdtheory.fi/public_html/articles/bioharmony2020.pdf.), 2020.
- [L8] Pitkänen M. Is genetic code part of fundamental physics in TGD framework? Available at: [https://tgdtheory.fi/public\\_html/articles/TIH.pdf.](https://tgdtheory.fi/public_html/articles/TIH.pdf.), 2021.
- [L9] Pitkänen M. Intersection form for 4-manifolds, knots and 2-knots, smooth exotics, and TGD. [https://tgdtheory.fi/public\\_html/articles/finitefieldsTGD.pdf.](https://tgdtheory.fi/public_html/articles/finitefieldsTGD.pdf.), 2022.
- [L10] Pitkänen M. Some New Ideas Related to Langlands Program *viz.* TGD. [https://tgdtheory.fi/public\\_html/articles/Langlands2022.pdf.](https://tgdtheory.fi/public_html/articles/Langlands2022.pdf.), 2022.
- [L11] Pitkänen M. About tessellations in hyperbolic 3-space and their relation to the genetic code . [https://tgdtheory.fi/public\\_html/articles/tessellationH3.pdf.](https://tgdtheory.fi/public_html/articles/tessellationH3.pdf.), 2023.
- [L12] Pitkänen M. About the Relationships Between Weak and Strong Interactions and Quantum Gravity in the TGD Universe . [https://tgdtheory.fi/public\\_html/articles/SW.pdf.](https://tgdtheory.fi/public_html/articles/SW.pdf.), 2023.
- [L13] Pitkänen M. Holography and Hamilton-Jacobi Structure as 4-D generalization of 2-D complex structure. [https://tgdtheory.fi/public\\_html/articles/HJ.pdf.](https://tgdtheory.fi/public_html/articles/HJ.pdf.), 2023.
- [L14] Pitkänen M. New findings related to the number theoretical view of TGD. [https://tgdtheory.fi/public\\_html/articles/M8Hagain.pdf.](https://tgdtheory.fi/public_html/articles/M8Hagain.pdf.), 2023.
- [L15] Pitkänen M. Symmetries and Geometry of the "World of Classical Worlds" . [https://tgdtheory.fi/public\\_html/articles/wcwsymm.pdf.](https://tgdtheory.fi/public_html/articles/wcwsymm.pdf.), 2023.
- [L16] Pitkänen M. A fresh look at  $M^8H$  duality and Poincare invariance. [https://tgdtheory.fi/public\\_html/articles/TGDcritics.pdf.](https://tgdtheory.fi/public_html/articles/TGDcritics.pdf.), 2024.
- [L17] Pitkänen M. What gravitons are and could one detect them in TGD Universe? [https://tgdtheory.fi/public\\_html/articles/whatgravitons.pdf.](https://tgdtheory.fi/public_html/articles/whatgravitons.pdf.), 2024.